

DAFTAR PUSTAKA

- Bothamley, J., & Boyle, M. (2013). *Patofisiologi dalam kebidanan*. Jakarta: EGC.
- Carthy, D. M. and J. M. (1992). A Framework for Analyzing the Determinants of Maternal Mortality Author (s): James McCarthy and Deborah Maine Source : Studies in Family Planning , Vol . 23 , No . 1 (Jan . - Feb . , 1992), pp . 23-33 Published by : Population Council Stable URL : <http://doi.org/10.2307/231>, 23–33.
- Chen, X. K., Wen, S. W., Smith, G., Yang, Q., & Walker, M. (2006). Pregnancy-induced hypertension is associated with lower infant mortality in preterm singletons. *BJOG: An International Journal of Obstetrics and Gynaecology*, 113(5), 544–551. <http://doi.org/10.1111/j.1471-0528.2006.00898.x>
- Courneya, K. S., & Friedenreich, C. (2011). *Physical Activity and Cancer*. (K. S. C. C. M. Friedenreich, Ed.). Springer. <http://doi.org/10.1007/978-3-642-04231-7>
- Crowther, C. A. (1985). Eclampsia at Harare Maternity Hospital. An epidemiological study. *S Afr Med J*, 68, 927–929.
- Cunningham, F. G., Hauth, J. C., Leveno, K. J., III, L. G., Bloom, S. L., & Wenstrom, K. D. (2005). *Williams Obstetrics*. (D. Rouse, B. Rainey, C. Spong, & G. D. Wendel, Eds.) (22nd ed.). New York: McGraw-Hill Medical. <http://doi.org/10.1017/CBO9781107415324.004>
- Cunningham, F. G., Leveno, K. J., Bloom, S. L., Hauth, J. C., Rouse, D. J., & Spong, C. Y. (2014). *Williams Obstetrics 23rd Edition*. Texas: McGraw-Hill Education.
- Dahlan, M. S. (2010). *Mendiagnosis dan Menatalaksana 13 Penyakit Statistik : Disertai Aplikasi Program Stata*. Jakarta: Sagung Set.
- Dalmaz, C. A. ;Katia G. D. S. (2011). Risk factors for hypertensive disorders of pregnancy in Southern Brazil, 692–696. <http://doi.org/10.1590/S0104-42302011000600018>
- Duckitt, K., & Harrington, D. (2005). Risk factors for pre-eclampsia at antenatal booking: systematic review of controlled studies. *BMJ (Clinical Research Ed.)*, 330(7491), 565. <http://doi.org/10.1136/bmj.38380.674340.E0>
- Fagard, R. H., & Cornelissen, V. A. (2007). Effect of exercise on blood pressure control in hypertensive patients. *European Journal of Cardiovascular Prevention and Rehabilitation : Official Journal of the European Society of Cardiology, Working Groups on Epidemiology & Prevention and Cardiac Rehabilitation and Exercise Physiology*, 14(1), 12–17. Retrieved from <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=med5&NE>

WS=N&AN=17301622

- Gordis, L. (2008). *Epidemiology* (4th ed.). Philadelphia: Elsevier.
- Guðnadóttir, T. A., Bateman, B. T., Hernández-Díaz, S., Luque-Fernandez, M. A., Valdimarsdóttir, U., & Zoega, H. (2016). Body Mass Index, Smoking and Hypertensive Disorders during Pregnancy: A Population Based Case-Control Study. *Plos One*, *11*(3), e0152187. <http://doi.org/10.1371/journal.pone.0152187>
- Hoggatt, K. J. (2002). *Biostatistical Methods in Epidemiology*. New York: John Wiley & Sons Inc.
- Holmes, D., & Baker, P. N. (2011). *Buku Ajar Ilmu Kebidanan.pdf*. Jakarta: EGC.
- Huang, N., Duggan, K., & Harman, J. (2008). Lifestyle management of hypertension. *Australian Prescriber*, *31*(6), 150–153.
- James, P. R., & Nelson-Piercy, C. (2004). Management of hypertension before, during, and after pregnancy. *Heart (British Cardiac Society)*, *90*(12), 1499–1504. <http://doi.org/10.1136/hrt.2004.035444>
- Kaplan, N. M., & Flynn, J. T. (2006). *Kaplan 's Clinical Hypertension*. Lippincott Williams & Wilkins.
- Katz, M. H. (2006). *Multivariable Analysis: A Practical Guide for Clinicians Second Edition* (2nd ed.). Cambridge: Cambridge University Press. <http://doi.org/10.1017/CBO9781107415324.004>
- Kemendes RI. (2013a). *Buku Saku Pelayanan Kesehatan Ibu di Fasilitas Kesehatan Dasar dan Rujukan*. (E. M. Moegni, D. Ocviyanti, G. M. K. Sakti, R. Floranita, L. Lawintono, & S. J. Olam, Eds.) (I). Jakarta: KEMENKES RI.
- Kemendes RI. (2013b). *Pedoman Teknis Penemuan dan Tatalaksana Hipertensi*. <http://doi.org/10.1017/CBO9781107415324.004>
- Kemendes RI. (2013c). *RISSET KESEHATAN DASAR*.
- Kemendes RI. (2014). *Profil kesehatan Indonesia*. Jakarta: Kementerian Kesehatan Republik Indonesia.
- Kemendes RI. (2015a). *Profil Kesehatan Indonesia*. Jakarta. <http://doi.org/10.1017/CBO9781107415324.004>
- Kemendes RI. (2015b). *Rencana Strategis Kementerian Kesehatan Tahun 2015-2019*. Jakarta: Kementerian Kesehatan Republik Indonesia.
- Kiondo, P., Wamuyu-Maina, G., Bimenya, G. S., Tumwesigye, N. M., Wandabwa, J., & Okong, P. (2012). Risk factors for pre-eclampsia in Mulago Hospital, Kampala, Uganda. *Tropical Medicine and International*

Health, 17(4), 480–487. <http://doi.org/10.1111/j.1365-3156.2011.02926.x>

- Leeman, L., & Fontaine, P. (1997). Hypertensive disorders of pregnancy. *Clinical Obstetrics*, 133–154.
- Lemeshow, S., Jr, D. W. H., Klar, J., & Lwanga, S. K. (1990). Sample Size for Case-Control Studies. In S. Lemeshow, D. W. H. Jr, J. Klar, & S. K. Lwanga (Eds.), *Adequacy of Sample Size in Health Studies* (pp. 16–20). Chichester: John Wiley & Sons.
- Lind, J. M., Hennessy, A., & Chiu, C. L. (2015). Association Between a Woman's Age at First Birth and High Blood Pressure. *Medicine (Baltimore)*, 94(16), e697. <http://doi.org/10.1097/md.0000000000000697>
- Luc Dauchet, Emmanuelle Kesse-Guyot, Sébastien Czernichow, Sandrine Bertrais, Carla Estaquio, Sandrine Péneau, Anne-Claire Vergnaud, Stacie Chat-Yung, Katia Castetbon, Valérie Deschamps, Pauline Brindel, and S. H. (2007). Dietary patterns and blood pressure change over 5-y follow-up in the. *Journal of Clinical Nutrition*, 85(3), 1650–1656. <http://doi.org/85/6/1650> [pii]
- Magann, E. F., Evans, S. F., Weitz, B., & Newnham, J. (2002). Antepartum, intrapartum, and neonatal significance of exercise on healthy low-risk pregnant working women. *Obstetrics and Gynecology*. [http://doi.org/10.1016/S0029-7844\(01\)01754-9](http://doi.org/10.1016/S0029-7844(01)01754-9)
- Marfitarini, W. (2009). Hubungan pola konsumsi sayur, buah dan susu dengan hipertensi dalam kehamilan.
- Martaadisoebrata, D., Wirakusumah, F. f., & Effendi, J. S. (2016). *Obstetri patologi ilmu kesehatan reproduksi* (3rd ed.). Jakarta: EGC.
- Murti, B. (2003). *Prinsip dan Metode Riset* (2nd ed.). Yogyakarta: Gadjah Mada University Press.
- NHBPEP. (2000). *National High Blood Pressure Education Working Group Report on High Blood Pressure in Pregnancy*. (R. Gifford, P. August, G. Cunningham, L. Green, M. Lindheimer, & D. McNellis, Eds.) *National Institutes of Health NIH publication no. 00-3029*. NATIONAL INSTITUTES OF HEALTH.
- NICE. (2010). *Hypertension in pregnancy: diagnosis and management*. (J. A. Turner, S. A. Lowe, M. A. Brown, G. A. Dekker, S. Gatt, C. K. McLintock, ... J. M. Moutquin, Eds.) *Am J Obstet Gynecol* (Vol. 77). Manchester: National Institute for Health and Care Excellence. <http://doi.org/10.1111/j.1479-828X.2009.01003.x>
- Nugroho, T., & Utama, B. I. (2014). *Masalah Kesehatan Reproduksi Wanita*. Yogyakarta: Nuha Medika.

- O'Brien, R. M. (2007). A caution regarding rules of thumb for variance inflation factors. *Quality and Quantity*, 41(5), 673–690. <http://doi.org/10.1007/s11135-006-9018-6>
- Owiredu, W. K. B. A., Ahenkorah, L., Turpin, C. A., Amidu, N., & Laing, E. F. (2012). Putative risk factors of pregnancy-induced hypertension among Ghanaian pregnant women. *Journal of Medical and Biomedical Sciences*, 1(3), 62–76. Retrieved from <http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L372721184>
- Pace, B. (2001). High Blood Pressure During Pregnancy. *JAMA: The Journal of the American Medical Association*, 285(12), 1664–1664. <http://doi.org/10.1001/jama.285.12.1664>
- PERKI. (2015). *Pedoman tatalaksana pencegahan penyakit kardiovaskular pada perempuan* (I). Jakarta: Perhimpunan Dokter Spesialis Kardiovaskular Indonesia.
- Phaswana-Mafuya, N., Peltzer, K., Chirinda, W., Musekiwa, A., & Kose, Z. (2013). Sociodemographic predictors of multiple non-communicable disease risk factors among older adults in South Africa. *Global Health Action*, 6(1), 1–8. <http://doi.org/10.3402/gha.v6i0.20680>
- Quasem Iftekhar, Mrunal S. Shetye, Shiney C. Alex, Anjan Kumar Nag, P.S. Sarma, K. R. T. (2001). Prevalence, awareness, treatment and control of hypertension among the elderly in Bangladesh and India: a multicentre study. *Bulletin of the World Health Organization*, 79(6), 490–500. <http://doi.org/10.1291/hypres.27.703>
- RI, K. K. (2014). *Pedoman Gizi Seimbang*.
- Roberts, J. M., Pearson, G., Cutler, J., & Lindheimer, M. (2003). Summary of the NHLBI Working Group on research on hypertension during pregnancy. *Hypertension*, 41(3), 437–445. <http://doi.org/10.1161/01.HYP.0000054981.03589.E9>
- Ros, H. S., Cnattingius, S., & Lipworth, L. (1998). Comparison of risk factors for preeclampsia and gestational hypertension in a population-based cohort study. *American Journal of Epidemiology*, 147(11 SUPPL.), 1062–1070. Retrieved from <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed4&N=EWS=N&AN=1998180154>
- Rudra, P., Basak, S., Patil, D., & Latoo, M. Y. (2011). Recent Advances In Management Of Pre-Eclampsia. *Bjamp*, 4(3).
- Sastrawinata, S., Martaadisoebrata, D., & Wirakusumah, F. F. (2003). *Obstetri Patologi*. (S. Sastrawinata, D. Martaadisoebrata, & F. F. Wirakusumah, Eds.)

(2nd ed.). Jakarta: EGC.

- Sastroasmoro, S., & Ismael, S. (2010). *Dasar-Dasar Metodologi Penelitian Klinis*. (S. Sastroasmoro & S. Ismael, Eds.) (3rd ed.). Jakarta: Sagung Seto.
- SEARO. (2011). *Noncommunicable Diseases in the South-East Asia Region*. <http://doi.org/978-92-9022-413-6>
- Silva, L., Coolman, M., Steegers, E., Jaddoe, V., Moll, H., Hofman, a, ... Raat, H. (2008). Maternal educational level and risk of gestational hypertension: the Generation R Study. *Journal of Human Hypertension*, 22(7), 483–492. <http://doi.org/10.1038/jhh.2008.22>
- Sirait, A. M. (2007). Prevalensi hipertensi pada kehamilan di Indonesia dan berbagai faktor yang berhubungan (Riset Kesehatan Dasar 2007), 103–109.
- SOMANZ. (2014). *Guidelines for the Management of Hypertensive Disorders of Pregnancy*. (L. SA, B. L, L. K, M. Lp, M. Mr, N. Ra, ... S. Jm., Eds.). Society of Obstetric Medicine of Australia and New Zealand. Retrieved from http://www.somanz.org/pdfs/somanz_guidelines_2008.pdf
- Suleiman, A. K. (2014). Risk Factors on Hypertensive Disorders among Jordanian Pregnant Women. *Global Journal of Health Science*, 6(2), 138–144. <http://doi.org/10.5539/gjhs.v6n2p138>
- Taslim, R. W. R., Kundre, R., & Masi, G. (2016). Hubungan Pola Makan dan Stres dengan Kejadian Hipertensi Grade 1 dan 2 Pada Ibu Hamil di Wilayah Kerja Puskesmas Kamonji Kecamatan Palu Barat. *E-Journal Keperawatan*, 4.
- Tebeu, P. M., Foumane, P., Mbu, R., Fosso, G., Biyaga, P. T., & Fomulu, J. N. (2011). Risk factors for hypertensive disorders in pregnancy: a report from the maroua regional hospital, cameroon. *J Reprod Infertil*, 12(3), 227–234.
- Tumbelaka, A. R., Riono, P., Sastroasmoro, S., Wirjodiarjo, M., Pudjiastuti, P., & Firman, K. (2010). Pemilihan uji hipotesis. In S. Sastroasmoro & S. Ismael (Eds.), *Dasar-dasaar Metodologi Penelitian Klinis* (3rd ed., pp. 279–300). Jakarta: Sagung Seto.
- Van Den Hooven, E. H., De Kluizenaar, Y., Pierik, F. H., Hofman, A., Van Ratingen, S. W., Zandveld, P. Y. J., ... Jaddoe, V. W. V. (2011). Air pollution, blood pressure, and the risk of hypertensive complications during pregnancy: The generation r study. *Hypertension*, 57(3), 406–412. <http://doi.org/10.1161/HYPERTENSIONAHA.110.164087>
- Varady, K. a, & Jones, P. J. H. (2005). Combination diet and exercise interventions for the treatment of dyslipidemia: an effective preliminary strategy to lower cholesterol levels? *The Journal of Nutrition*, 135(8), 1829–1835.

- Wagner, L. K. (2004). Diagnosis and management of preeclampsia. *American Family Physician*, 70(12), 2317–2324.
- WHO. (2003). *DIET , NUTRITION AND THE PREVENTION OF Chronic diseases. DIET, NUTRITION AND THE PREVENTION OF Chronic diseases: report of a Joint WHO/FAO Expert Consultation, NUTRITION AND THE PREVENTION OF Chronic diseases: report of a Joint WHO/FAO Expert Consultation*. <http://doi.org/0512-3054>
- WHO. (2008). *Conducting the Survey, Data Entry, Data Analysis and Reporting and Disseminating Results Overview*. Geneva: World Health Organization.
- WHO. (2011). *Prevention and treatment of pre-eclampsia and eclampsia*. World Health. Geneva: World Health Organization.
- Wong, T. Y., Groen, H., Faas, M. M., & Van Pampus, M. G. (2013). Clinical risk factors for gestational hypertensive disorders in pregnant women at high risk for developing preeclampsia. *Pregnancy Hypertension*, 3(4), 248–253. <http://doi.org/10.1016/j.preghy.2013.07.003>
- Wu, W. (2010). Confounding , interaction , and mediation in multivariable / multivariate regression modeling logo. *Cancer Biostatistics*.
- Ye, C., Ruan, Y., Zou, L., Li, G., Li, C., Chen, Y., ... Zhang, W. (2014). The 2011 survey on hypertensive disorders of pregnancy (HDP) in China: Prevalence, risk factors, complications, pregnancy and perinatal outcomes. *PLoS ONE*, 9(6). <http://doi.org/10.1371/journal.pone.0100180>