EXPRESSION OF LUTEINIZING HORMONE RECEPTOR (LHR) IN THE LEYDIG CELL CULTURE OF SPRAGUE DAWLEY RAT
THAT INDUCED BY ADVANCED GLYCATION END-PRODUCT 200MICRO GRAM PER MILI LITER AND GAMMA MANGOSTEEN
5MIKRO MOL
MUHAMMAD HERDHANA ASH SHIDIQI, Dr dr. Dicky Moch. Rizal,Sp. And., AIFM, M.Kes ; dr. Rustamaji,M.Kes
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

CHAPTER VI. REFERENCES

Ames, BN, Shigenaga, MK, Hagen, TM, 1993, Oxidants, antioxidants, and the degenerative diseases of aging. *Proc Natl Acad Sci U S A*, vol.90, no.17, pp.7915-7922.

Basta, G, Schmidt, AM, Caterina, RD 2004, Advanced Glycation End Products and Vascular Inflammation: Implications for Accelerated AtherosclerosisiIn Diabetes. *J Cardiovasc,* vol.63, pp.582- 592.

Bayne, JW, Thorpe, SR 1999, Role of oxydative stress in diabetic complication a new perspective on an old papadigm. *J Diabet,* vol.48, pp.1-9.

Bierhaus, A, Chevion, M, Hofmann, M, Quehenberger, P, Illmer, T, Luther, T, Berentshtein, E, Muller, M, Wahl, P, Ziegler R, Nawroth, 1997, Advanced Glycation End Product-Induced Activation of NF-KB IS Suppressed by a-Lipoic Acid in Cultured Endothelial Cells. *Int med,* vol.46 pp.1481-1489.

EXPRESSION OF LUTEINIZING HORMONE RECEPTOR (LHR) IN THE LEYDIG CELL CULTURE OF SPRAGUE DAWLEY RAT
THAT INDUCED BY ADVANCED GLYCATION END-PRODUCT 200MICRO GRAM PER MILI LITER AND GAMMA MANGOSTEEN
5MIKRO MOL
MUHAMMAD HERDHANA ASH SHIDIQI, Dr dr. Dicky Moch. Rizal,Sp. And., AIFM, M.Kes ; dr. Rustamaji,M.Kes
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Boel, E, Selmer, J, Flodgaard, HJ, Jensen, T 1995, Diabetic late complicatioms: will aldose reductase inhibitor or inhibitors of advanced glysosylation end-product formation hold promise?, *J Diabet Complication*, vol.9, pp.104-129.

Brownlee, M 1995, Advanced protein glycosylation in diabetes and aging. *Ann Rev Med*, vol.46, pp.223–234.

Brownlee, M, Vlassara, H, Cerami, A 1985, Nonenzymatic glycosylation products on collagen covalently trap low-density lipoprotein. *J Diabet*, vol.34, pp.938–941.

Bumrungpert, A, Ruchaneekorn, W, Kalpravidh, Chitchumroonchokchai, C, Chuang, C, West, T, Kennedy A, McIntosh, M 2009. Xanthones from Mangosteen Prevent Lipopolysaccharide-Mediated In ammation and Insulin Resistance in Primary Cultures of Human Adipocytes. *Dep Nutri*, vol.139 pp.1185-1191.

UNIVERSITAS GADJAH MADA
EXPRESSION OF LUTEINIZING HORMONE RECEPTOR (LHR) IN THE LEYDIG CELL CULTURE OF SPRAGUE DAWLEY RAT THAT INDUCED BY ADVANCED GLYCATION END-PRODUCT 200MICRO GRAM PER MILI LITER AND GAMMA MANGOSTEEN 5MIKRO MOL
MUHAMMAD HERDHANA ASH SHIDIQI, Dr dr. Dicky Moch. Rizal,Sp. And., AIFM, M.Kes ; dr. Rustamaji,M.Kes
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Ceriello, A 1999, Hyperglycaemia: the bridge between non-enzymatic glycation and oxidative stress in the pathogenesis of diabetic complications. *Diabet Nutri Met*, vol.12, pp.42-46.

Chang, H, Yang, L 2012. Gamma-Mangostin, a Micronutrient of Mangosteen Fruit, Induces Apoptosis in Human Colon Cancer Cells. *Pharmacol,* vol.17 pp.8010-8021.

Chen, H, Matthew, P, Hardy, Barry, R, Zirkin 2002, Age-Related Decreases in Leydig Cell Testosterone Production Are Not Restored by Exposure to LH in Vitro. *The Endocrinol soc*, vol. 143, no.5, pp.1637-1642.

Chin, YA. Kinghorn, D, 2008, Structural Characterization, Biological Effects, and Synthetic Studies on Xanthones from Mangosteen (Garcinia mangostana), *a Popular Botanical Dietary Supplement.* Rev Org Chem, vol.5, no.4, pp.355-364.

UNIVERSITAS GADJAH MADA
EXPRESSION OF LUTEINIZING HORMONE RECEPTOR (LHR) IN THE LEYDIG CELL CULTURE OF SPRAGUE DAWLEY RAT THAT INDUCED BY ADVANCED GLYCATION END-PRODUCT 200MICRO GRAM PER MILI LITER AND GAMMA MANGOSTEEN 5MIKRO MOL
MUHAMMAD HERDHANA ASH SHIDIQI, Dr dr. Dicky Moch. Rizal,Sp. And., AIFM, M.Kes ; dr. Rustamaji,M.Kes
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Esser, K, Stadtman, ER, Wiley, C, Martin, GM 1995, The status of oxidatively modified proteins as a marker of aging, *Eds. Mol Aspects of Aging*, pp.129-144.

Fu, MX, Wells-Knecht, KJ, Blackledge, JA, Lyons, TJ, Thorpe, SR, Baynes, JW 1994. Glycation, glycoxidation, and cross-linking of collagen by glucose kinetics, mechanisms, and inhibition of late stages of the Maillard reaction. *J Diabet,* vol.43, pp.676-683.

Fulda, S, Gorman, AM, Hori, S, Samali,A 2010, Cellular Stress Responses: Cell Survival and Cell Death. *Inter J Cell Biol*, Vol.2010, Article ID.214074, pp. 23. http://dx.doi.org/10.1155/2010/214074 (accessed on 1 december 2015).

Goldin, A, Beckman, JA, Schmidt, AM, Creager, MA, 2015, Advanced Glycation End Products Sparking the

EXPRESSION OF LUTEINIZING HORMONE RECEPTOR (LHR) IN THE LEYDIG CELL CULTURE OF SPRAGUE DAWLEY RAT
THAT INDUCED BY ADVANCED GLYCATION END-PRODUCT 200MICRO GRAM PER MILI LITER AND GAMMA MANGOSTEEN
5MIKRO MOL
MUHAMMAD HERDHANA ASH SHIDIQI, Dr dr. Dicky Moch. Rizal,Sp. And., AIFM, M.Kes ; dr. Rustamaji,M.Kes
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Development of Diabetic Vascular Injury. *Cardiovasc Division, Boston* pp.4-5.

Goluza, 2014, Macrophages and Leydig Cells in Testicular Biopsies of Azoospermic Men, *Biol Med Research Inter.* http://dx.doi.org/10.1155/2014/828697 (accessed on 1 April 2015)

Huebschmann, AG, Regensteiner, JG, Vlassara, H, Reusch, JEB 2006, Diabetes and Advanced Glycoxidation End Products. *Diabet Care*, vol.29, pp.1420.

Huo, S, Zhong, X, Wu, X, Li, Y 2012, Effect of norepinephrine and acetylcholine on the development of culture leydig cells in mice. *J Biol med and Biotech,* vol.2012, Article ID.503093, pp.8

Irina, G, Tikhonova, Costanzi, S 2009, Unraveling the structure and function of G protein-coupled receptors through NMR spectroscopy. *Lab Biol Mod*, Vol.15, No.35, pp.4003–4016.

UNIVERSITAS GADJAH MADA
EXPRESSION OF LUTEINIZING HORMONE RECEPTOR (LHR) IN THE LEYDIG CELL CULTURE OF SPRAGUE DAWLEY RAT
THAT INDUCED BY ADVANCED GLYCATION END-PRODUCT 200MICRO GRAM PER MILI LITER AND GAMMA MANGOSTEEN
5MIKRO MOL
MUHAMMAD HERDHANA ASH SHIDIQI, Dr dr. Dicky Moch. Rizal,Sp. And., AIFM, M.Kes ; dr. Rustamaji,M.Kes
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Li, J, Schmidt, AM 1997, Characterization and functional analysis of the promoter of RAGE, the receptor for advanced glycation end products. *J Biol Chem* vol.272 pp.16498-16506.

Linn, S, Esser, K, Martin, GM, Chichester, W 1995, DNA repair in mitochondria: How is it limited? What is its function? *Eds. Mol Aspects of Aging*, pp.199-225.

Liu, G, Duranteu, L, Claue, J, Monroe, J, Daniel, A, Doyle, Shenker, A 1999, Leydig-cell Tumors Caused by an Activating Mutation of Gene Encoding The Luteinizing Hormone Receptor. *Brief Report* vol.341, pp.1731-1735.

Liu, J, Yan, L, Niu, R, Tian, L, Zhang, Q, Quan, J, Liu, H, Wei, S, Guo, Q 2013, Protection Effect of Endomorphins on Advanced Glycation End Products Induced Injury in Endothelial Cells. *J Diabet Research,* Vol.2013, Article.ID 105780, pp.9.

EXPRESSION OF LUTEINIZING HORMONE RECEPTOR (LHR) IN THE LEYDIG CELL CULTURE OF SPRAGUE DAWLEY RAT
THAT INDUCED BY ADVANCED GLYCATION END-PRODUCT 200MICRO GRAM PER MILI LITER AND GAMMA MANGOSTEEN
5MIKRO MOL
MUHAMMAD HERDHANA ASH SHIDIQI, Dr dr. Dicky Moch. Rizal,Sp. And., AIFM, M.Kes ; dr. Rustamaji,M.Kes
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Loo AEK, Huang D, 2007, Assay-guided Fractionation study of ά-amylase inhibitors from *Garcia Mangostana* pericarp. *J Agricult Food Chem*, Vol.55, pp.9805-9810.

Morgan, JA, Linch, J, Schuetz, JD 2015, Apotosome activation, an important molecular instigator in 6-mercapturopurine induced leydig cell death. *Dep Pharmacol sci,* vol.5 pp.16488.

Nakatani, K, Nakahata, N, Arakawa, T,Yasuda, H, Ohizumi, Y, 2002, Inhibition of cyclooxygenase and prostaglandin E2 synthesis by -mangostin, a xanthone derivative in mangosteen, in C6 rat glioma cells. *J Biochem Pharmacol*, vol.63, pp.73–79.

Nakatani, K, Yamakuni, T, Kondo, N, Arakawa, T, Oosawa, K, Shimura, S, Inoue, H, Ohizumi, Y 2004, γ–Mangostin inhibits Inhibitor-kB kinase activity and decreases lipopolysaccharide-induced cycloocygenase-2 gene

EXPRESSION OF LUTEINIZING HORMONE RECEPTOR (LHR) IN THE LEYDIG CELL CULTURE OF SPRAGUE DAWLEY RAT
THAT INDUCED BY ADVANCED GLYCATION END-PRODUCT 200MICRO GRAM PER MILI LITER AND GAMMA MANGOSTEEN
5MIKRO MOL
MUHAMMAD HERDHANA ASH SHIDIQI, Dr dr. Dicky Moch. Rizal,Sp. And., AIFM, M.Kes ; dr. Rustamaji,M.Kes
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

expression in C6 rat glioma cells. J *Mol Pharmacol,* vol.66, pp.667-674.

Nieschlag, E, Behre, HM, 2010, *Textbook of Male Reproductive Health and Dysfunction*, 3th edn, Springer, Berlin.

Ogiwara, K, Fujimori, C, Rajapakse, C, Takahashi, T 2013, Characterization of Luteinizing Hormone and Luteinizing Hormone Receptor and Their Indispensable Role in the Ovulatory Process of the Medaka. *PLos ONE,* 8(1), p.e54482.

Rajfer , J 2003, Decreased Testosterone in the Aging Male. *J Rev of Urol*, 5(Suppl 1):S1-S2.

Ran, O, Gu, M, Van Remmen, H 2006. Glutatione peroxidase 4 protects cortical neurons from oxidative injury and amyloid toxicity. *J Neuro Sci Res,* vol.84, pp.202-208.

Rasheed, Z, Akhtar, N, Tariq, Haqqi, M 2011, Advanced glycation end products induce the expression of

EXPRESSION OF LUTEINIZING HORMONE RECEPTOR (LHR) IN THE LEYDIG CELL CULTURE OF SPRAGUE DAWLEY RAT
THAT INDUCED BY ADVANCED GLYCATION END-PRODUCT 200MICRO GRAM PER MILI LITER AND GAMMA MANGOSTEEN
5MIKRO MOL
MUHAMMAD HERDHANA ASH SHIDIQI, Dr dr. Dicky Moch. Rizal,Sp. And., AIFM, M.Kes ; dr. Rustamaji,M.Kes
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

interleukin-6 and interleukin-8 by receptor for advanced glycation end product-mediated activation of mitogen-activated protein kinases and nuclear factor-κB in human osteoarthritis chondrocytes. J *Rheumatol (Oxford)*, vol. 50, no.5 pp. 838–851.

Rizal, DM 2014, Kadar Testosteron, Ekspresi COX-2, Reseptor Advanced Glycation End Products, dan Reseptor Luteinizing Hormone Pada Kultur Sel Leydig Tikus Sprague Dawley yang diinduksi Advanced Glycation End Products dan diberi *Gamma-mangostin*, Proposal Disertasi Doktoral. Universitas Gadjah Mada, Yogyakarta.

Roess, DA, Brady, CJ, Barisas, BG, 2000, Biological function of the LH receptor is associated with slow receptor rotational diffusion. Vol.1464.

Rothstein, M, Lippman, RD, Liss, AR 1983, Lipid peroxidation and metabolism in aging. *Rev Biol Research in Aging*, vol.1, pp.315-342.

UNIVERSITAS GADJAH MADA
EXPRESSION OF LUTEINIZING HORMONE RECEPTOR (LHR) IN THE LEYDIG CELL CULTURE OF SPRAGUE DAWLEY RAT THAT INDUCED BY ADVANCED GLYCATION END-PRODUCT 200MICRO GRAM PER MILI LITER AND GAMMA MANGOSTEEN 5MIKRO MOL
MUHAMMAD HERDHANA ASH SHIDIQI, Dr dr. Dicky Moch. Rizal,Sp. And., AIFM, M.Kes ; dr. Rustamaji,M.Kes
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Sasaki, N, Fukatsu, R, Tsuzuki, K, Hayashi, K, Yoshida, T, Fujii, N, Wakayama, I, Yanagihara, R, Garutto, R, Amano, N, Makita, Z, 1998, Advanced Glycation End-products in Alzheimer's Disease and Other Neurodegenerative Disease. *Am J Pathol*, Vol.153, no.4, pp.1149–1155.

Schmidt AM, Yan SD, Yan SF, Stern, DM 2001, The multiligand receptor RAGE as a progression factor amplifying immune and inflammatory responses. *J Clin Invest*, vol.108, pp.949–955

Schmidt, AM, Mora, R, Cao, R, Yan, SD,Brett , J, Ramakrishnan, R, Tsang, TC, Simionescu, M, Stern, D 1994, The endothelial cell binding site for advanced glycation end products consists of a complex: an integral membrane protein and a lactoferrin-like polypeptide. *J Biol Chem* vol.269, pp.9882–9888.

UNIVERSITAS GADJAH MADA
EXPRESSION OF LUTEINIZING HORMONE RECEPTOR (LHR) IN THE LEYDIG CELL CULTURE OF SPRAGUE DAWLEY RAT THAT INDUCED BY ADVANCED GLYCATION END-PRODUCT 200MICRO GRAM PER MILI LITER AND GAMMA MANGOSTEEN 5MIKRO MOL
MUHAMMAD HERDHANA ASH SHIDIQI, Dr dr. Dicky Moch. Rizal,Sp. And., AIFM, M.Kes ; dr. Rustamaji,M.Kes
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Schmidt, AM, Vianna, M, Gerlach, M, Brett, J, Ryan, J, Kao, J, Esposito, C, Hegarty, H, Hurley, W, Clauss, M, Clauss, FW, Yu-Ching, EP, Tsang, TC, Stern, D 1992, Isolation and characterization of two binding proteins for advanced glycosylation end products from bovine lung which are present on the endothelial cell surface. *J Biol Chem*, vol.267, pp.14987-14997.

Schmidt, AM, Yan, SD, Wautier, JL, Stern, D 1999, Activation of receptor for advanced glycation end products: a mechanism for chronic vascular dysfunction in diabetic vasculopathy and atherosclerosis. *Circ Resp*, vol.84, pp.489-497.

Sherwood, L. 2011. *Textbook of Fundamentals Human physiology*, 4th edn, USA, West virginia.

Shibata, MA, Matoba, Y, Tosa, H, Iinuma, M, 2013, Effects of Mangosteen Pericarp Extracts Against Mammary Cancer. Laboratory of Anatomy and Histopathology. *Altern*

UNIVERSITAS GADJAH MADA
EXPRESSION OF LUTEINIZING HORMONE RECEPTOR (LHR) IN THE LEYDIG CELL CULTURE OF SPRAGUE DAWLEY RAT THAT INDUCED BY ADVANCED GLYCATION END-PRODUCT 200MICRO GRAM PER MILI LITER AND GAMMA MANGOSTEEN 5MIKRO MOL
MUHAMMAD HERDHANA ASH SHIDIQI, Dr dr. Dicky Moch. Rizal,Sp. And., AIFM, M.Kes ; dr. Rustamaji,M.Kes
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Integrative Med, vol.2, No.8, pp.2327-5162. Vailable on http://dx.doi.org/10.4172/2327-5162.1000139.

Singh, R, Barden, A, Mori, T, Beilin, L 2001, Advanced glycation end-product. Rev *Diabetologia*, vol.44, pp.129-146.

Sivakumar, R, Sivaraman, BP, Mohan-Babu, N, Jainul-Abideen, IM, Kalliyappan, P, Balasubramanian, K, 2006, Radiation Exposure Impairs Luteinizing Hormone Signal Transduction and Steroidogenesis in Cultured Human Leydig Cells. *Dep Endoc*, vol.91, no.2, pp.550–556.

Sugaya, K, Fukagawa, T, Matsumoto, K, Mita, K, Takahashi, E, Ando, A, Inoko, H, Ikemura, T 1994, Three genes in the human MHC class III region near the junction with the class II gene for receptor of advanced glycosylation end products, PBX2 homeobox gene and a notch homolog, human counterpart of mouse mammary tumor gene int-3. J *Genom*, vol.23, pp.408–419.

UNIVERSITAS GADJAH MADA
EXPRESSION OF LUTEINIZING HORMONE RECEPTOR (LHR) IN THE LEYDIG CELL CULTURE OF SPRAGUE DAWLEY RAT THAT INDUCED BY ADVANCED GLYCATION END-PRODUCT 200MICRO GRAM PER MILI LITER AND GAMMA MANGOSTEEN 5MIKRO MOL
MUHAMMAD HERDHANA ASH SHIDIQI, Dr dr. Dicky Moch. Rizal,Sp. And., AIFM, M.Kes ; dr. Rustamaji,M.Kes
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Tena-sempere, M, Rannikko, A, Kero, J, Zhang, FP, Huhtainiemi, IT 1997, Molecular Mechanisms of Reappearance of Luteinizing Hormone Receptor Expression and Function in Rat Testis after Selective Leydig Cell Destruction by Ethylene Dimethane Sulfonate. *Dep Physiol*, vol.138, no.1, pp.3340-3348.

Valko, M, Leibfritz, D, Moncol, J, Cronin, MT, Mazur, M, Telser, J 2007, Free radicals and antioxidants in normal physiological functions and human disease. *Inter J Biol Chem Cell*, vol.39, no.1 pp.44-84.

Wickens, AP 2001, Ageing and the free radical theory. *Resp Physiol*, vol.128, no.3, pp.379-391.

Yan, SF, Ramasamy, R, Naka, Y, Schmidt, AM 2003, Glycation, inflammation, and RAGE: a scaffold for the macrovascular complications of diabetes and beyond. *Circ Resp*, vol.93, pp.1159-1169.