



## DAFTAR PUSTAKA

- AAP, 2012. Levels of neonatal care. *Pediatrics* 130: 587–97.
- ACOG, AAP, 2016. Committee Opinion Number 644: The Apgar Score. *Am. Acad. Pediatr. Fetus Newborn* 1–4.
- ACOG committee opinion, 1992. Utility of umbilical cord blood acid-base assessment. *Int. J. Gynecol. Obstet.* 38: 61–62.
- Adcock, L., Stark, A., 2013. Systemic effects of perinatal asphyxia. *Uptodate*. available at <http://www.uptodate.com/contents/systemic-effects-of-perinatal-asphyxia> [cited 3.27.16].
- Adsett, D.B., Fitz, C.R., Hill, A., 1985. Hypoxic-ischaemic cerebral injury in the term newborn: correlation of CT findings with neurological outcome. *Dev. Med. Child Neurol.* 27: 155–160.
- Afjeh, S.-A., Sabzehei, M.-K., Esmaili, F., 2013. Neonatal resuscitation in the delivery room from a tertiary level hospital: risk factors and outcome. *Iran. J. Pediatr.* 23: 675–680.
- Agrawal, 2004. Role of lactate in critically ill children. *Indian J. Crit. Care Med.* 8: 173.
- Aiken, C.G., Bardgett, R.J.M., 2013. Neonatal Blood Gases and Outcome Following Perinatal Asphyxia. *Indian J. Neonatal Med. Res.* 1: 1–4.
- Akinbi, H., Abbasi, S., Hilpert, P.L., et al., 1994. Gastrointestinal and renal blood flow velocity profile in neonates with birth asphyxia. *J. Pediatr.* 125: 625–627.
- Allen, L.H., 2000. Anemia and iron deficiency: effects on pregnancy outcome. *Am. J. Clin. Nutr.* 71: 1280S–1284S.
- Andrews, D.A., Sawin, R.S., Ledbetter, D.J., et al., 1990. Necrotizing enterocolitis in term neonates. *Am. J. Surg.* 159: 507–509.
- Arnold, R.C., Shapiro, N.I., Jones, A.E., et al., 2009. Multicenter study of early lactate clearance as a determinant of survival in patients with presumed sepsis. *Shock* 32: 35–39.
- Aslam, H.M., Saleem, S., Afzal, R., et al., 2014. "Risk factors of birth asphyxia"; *Ital. J. Pediatr.* 40: 94.
- Backes, C.H., Markham, K., Moorehead, P., et al., 2011. Maternal preeclampsia and neonatal outcomes. *J. Pregnancy* 2011: 214365.
- Bakker, J., Gris, P., Coffernils, M., et al., 1996. Serial blood lactate levels can predict the development of multiple organ failure following septic shock. *Am. J. Surg.* 171: 221–226.
- Bakker, J., Nijsten, M.W., Jansen, T.C., 2013. Clinical use of lactate monitoring in critically ill patients. *Ann. Intensive Care* 3: 12.
- Baldari, C., Bonavolontà, V., Emerenziani, G., et al., 2009. Accuracy, reliability, linearity of Accutrend and Lactate Pro versus EBIO plus analyzer. *Eur. J. Appl. Physiol.* 107: 105–111.
- Barber, C.A., Wyckoff, M.H., 2006. Use and efficacy of endotracheal versus intravenous epinephrine during neonatal cardiopulmonary resuscitation in the delivery room. *Pediatrics* 118: 1028–1034.



- Behar, K.L., Rothman, D.L., Shulman, R.G., *et al.*, 1984. Detection of cerebral lactate in vivo during hypoxemia by  $^1\text{H}$  NMR at relatively low field strengths (1.9 T). *Proc. Natl. Acad. Sci. U. S. A.* 81: 2517–2519.
- Berseth, C.L., McCoy, H.H., 1992. Birth asphyxia alters neonatal intestinal motility in term neonates. *Pediatrics* 90: 669–673.
- Blair, E., Stanley, F.J., 1988. Intrapartum asphyxia: a rare cause of cerebral palsy. *J. Pediatr.* 112: 515–519.
- Blomkalns, A.L., 2007. Lactate - A Marker for Sepsis and Trauma. *EMCREG International* 43–49.
- Bryce, J., Boschi-Pinto, C., Shibuya, K., *et al.*, 2005. WHO estimates of the causes of death in children. *Lancet (London, England)* 365: 1147–1152.
- Cady, L.D., Weil, M.H., Afifi, A.A., *et al.*, 1973. Quantitation of severity of critical illness with special reference to blood lactate. *Crit. Care Med.* 1: 75–80.
- Casey, B.M., McIntire, D.D., Leveno, K.J., 2001. The Continuing Value of the Apgar Score for the Assessment of Newborn Infants. *N. Engl. J. Med.* 344: 467–471.
- Castle, V., Andrew, M., Kelton, J., *et al.*, 1986. Frequency and mechanism of neonatal thrombocytopenia. *J. Pediatr.* 108: 749–755.
- Cheung, P.Y., Robertson, C.M., 2000. Predicting the outcome of term neonates with intrapartum asphyxia. *Acta Paediatr.* 89: 262–264.
- Cowan, F., Rutherford, M., Groenendaal, F., Eken, P., Mercuri, E., Bydder, G. M., Meiners, L. C., Dubowitz, L. M. S. & de Vries, L.S., 2003. Origin and timing of brain lesions in term infants with neonatal encephalopathy. *Lancet* 361: 736–742.
- de Oliveira, T.G., Freire, P.V., Moreira, F.T., *et al.*, 2012. Apgar score and neonatal mortality in a hospital located in the southern area of Sao Paulo City, Brazil. *Einstein* 10: 22–28.
- Deshpande, S.A., Platt, M.P., 1997. Association between blood lactate and acid-base status and mortality in ventilated babies. *Arch. Dis. Child. Fetal Neonatal Ed.* 76: F15–F20.
- Dharmasetiawani, N., 2012. Resusitasi Neonatus Konsensus 2010. *Perinasia*. available at [http://www.perinasia.com/post/116?title=Resusitasi+Neonatus+\(Konsensus+2010\)](http://www.perinasia.com/post/116?title=Resusitasi+Neonatus+(Konsensus+2010)) [cited 5.2.16].
- Durkan, A.M., Alexander, R.T., 2011. Acute kidney injury post neonatal asphyxia. *J. Pediatr.* 158: e29–e33.
- Eerden, P., Bernstein, P., 2003. Summary of the Publication, “Neonatal Encephalopathy and Cerebral Palsy: Defining the Pathogenesis and Pathophysiology,” by the ACOG Task Force on Neonatal Encephalopathy and Cerebral Palsy. *Medscape*. available at <http://www.medscape.com/viewarticle/457882> [cited 4.25.16].
- Faix, R.G., Viscardi, R.M., DiPietro, M.A., *et al.*, 1989. Adult respiratory distress syndrome in full-term newborns. *Pediatrics* 83: 971–976.
- Fauchère, J.C., Bauschatz, A.S., Arlettaz, R., *et al.*, 2002. Agreement between capillary and arterial lactate in the newborn. *Acta Paediatr.* 91: 78–81.
- Fernandez, H.G.C., Vieira, A.A., Barbosa, A.D.M., 2012. The correlation between



- plasma lactate concentrations and early neonatal mortality. *Rev. Bras. Ter. Intensiva* 24: 184–187.
- Franklin, R., Peloso, J., 2006. Review of the clinical use of lactate. *AAEP Proc.* 52: 305–309.
- Gaieski, D.F., Drumheller, B.C., Goyal, M., et al., 2013. Accuracy of Handheld Point-of-Care Fingertip Lactate Measurement in the Emergency Department. *West. J. Emerg. Med.* 14: 58–62.
- Goldberg, R.N., Moscoso, P., Bauer, C.R., et al., 1986. Use of barbiturate therapy in severe perinatal asphyxia: a randomized controlled trial. *J. Pediatr.* 109: 851–6.
- Gunn, A.J., Bennet, L., Gunning, M.I., et al., 1999. Cerebral hypothermia is not neuroprotective when started after postischemic seizures in fetal sheep. *Pediatr. Res.* 46: 274–280.
- Gupta, B.D., Sharma, P., Bagla, J., et al., 2005. Renal failure in asphyxiated neonates. *Indian Pediatr.* 42: 928–934.
- Hamerman, C., Zadka, P., 1994. Asphyxia-related infant mortality rates. *Am. J. Perinatol.* 11: 290–294.
- Helmy, M., 2014. Mechanisms of Birth Asphyxia and a Novel Resuscitation Strategy.
- Huckabee, W., 1961. Abnormal resting blood lactate. I. The significance of hyperlactatemia in hospitalized patients. *Am. J. Med.* 30: 840–848.
- Jezova, H., Hotarkova, S., Muckova, C., et al., 2013. Birth asphyxia: Atlas of neonatal pathology. available at [https://atlases.muni.cz/atlas/novo/atl\\_en/main+novorozene+novorasycas.html](https://atlases.muni.cz/atlas/novo/atl_en/main+novorozene+novorasycas.html) [cited 5.3.16].
- Johnston, M. V, Fatemi, A., Wilson, M.A., et al., 2011. Treatment advances in neonatal neuroprotection and neurointensive care. *Lancet. Neurol.* 10: 372–82.
- Juliantari, I.G.A., Dasuki, D., 2014. Analisis Faktor Risiko Kematian Neonatal Dini Karena Asfiksia Di RSUP Nusa Tenggara Barat. available at [http://etd.repository.ugm.ac.id/index.php?mod=penelitian\\_detail&sub=PenelitianDetail&act=view&typ=html&buku\\_id=69339](http://etd.repository.ugm.ac.id/index.php?mod=penelitian_detail&sub=PenelitianDetail&act=view&typ=html&buku_id=69339) [cited 6.8.16].
- Karlsson, M., Blennow, M., Nemeth, A., et al., 2006. Dynamics of hepatic enzyme activity following birth asphyxia. *Acta Paediatr.* 95: 1405–1411.
- Kattwinkel, J., Perlman, J.M., Aziz, K., et al., 2010. Part 15: Neonatal resuscitation: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 122.
- Kemenkes, 2015. Profil Kesehatan Indonesia 2014. available at <http://www.depkes.go.id/resources/download/pusdatin/profil-kesehatan-indonesia/profil-kesehatan-indonesia-2014.pdf> [cited 4.24.15].
- Kiondo, P., Tumwesigye, N.M., Wandabwa, J., et al., 2014. Adverse neonatal outcomes in women with pre-eclampsia in Mulago Hospital, Kampala, Uganda: a cross-sectional study. *Pan Afr. Med. J.* 17 Suppl 1: 7.
- Kost, G.J., Nguyen, T.H., Tang, Z., 2000. Whole-blood glucose and lactate. Trilayer biosensors, drug interference, metabolism, and practice guidelines. *Arch. Pathol. Lab. Med.* 124: 1128–1134.
- Kruse, J., 2002. Blood Lactate Concentrations in Sepsis. Kluwer Academic Publishers, Boston.



- Kruse, J.A., Zaidi, S.A., Carlson, R.W., 1987. Significance of blood lactate levels in critically ill patients with liver disease. *Am. J. Med.* 83: 77–82.
- Lam, B.C.C., Yeung, C.-Y., 1992. Perinatal Features of Birth Asphyxia and Neurologic Outcome. *Pediatr. Int.* 34: 17–22.
- Lane, R., 2002. Blood lactate measurement could lead to earlier identification of patients for liver transplant. *Lancet.* available at [http://www.eurekalert.org/pub\\_releases/2002-02/l-blm021302.php](http://www.eurekalert.org/pub_releases/2002-02/l-blm021302.php) [cited 5.18.16].
- Lawn, J.E., Bahl, R., Bergstrom, S., et al., 2011. Setting research priorities to reduce almost one million deaths from birth asphyxia by 2015. *PLoS Med.* 8: e1000389.
- Leal, Y.A., Álvarez-Nemegyei, J., Velázquez, J.R., et al., 2012. Risk factors and prognosis for neonatal sepsis in southeastern Mexico: analysis of a four-year historic cohort follow-up. *BMC Pregnancy Childbirth* 12: 48.
- Lee, A.C.C., Mullany, L.C., Tielsch, J.M., et al., 2008. Risk factors for neonatal mortality due to birth asphyxia in southern Nepal: a prospective, community-based cohort study. *Pediatrics* 121: e1381–e1390.
- Lemeshow, S., David, W., 1997. Besar sampel dalam penelitian kesehatan (terjemahan). Gadjahmada University Press, Yogyakarta.
- Li, F., Wu, T., Lei, X., et al., 2013. The Apgar Score and Infant Mortality. *PLoS One* 8: 1–8.
- Li-juan, Q., YAang, L., Qiao, L., et al., 2011. Clinical Value of Early Lactate Clearance Rate on Evaluation of Prognosis in Severe Asphyxia Neonate. *J Appl Clin Pediatr* 26: 427–429.
- Lincetto, O., 2007. Birth Asphyxia - Summary of the Previous Meeting and Protocol Overview. available at <http://curoservice.com/Accessedon12/2/2015> [cited 2.5.16].
- Low, J.A., Galbraith, R.S., Muir, D.W., et al., 1988. Motor and cognitive deficits after intrapartum asphyxia in the mature fetus. *Am. J. Obstet. Gynecol.* 158: 356–361.
- Luft, F.C., 2001. Lactic acidosis update for critical care clinicians. *J. Am. Soc. Nephrol.* 12 Suppl 1: S15–S19.
- Lupton, B.A., Hill, A., Roland, E.H., et al., 1988. Brain swelling in the asphyxiated term newborn: pathogenesis and outcome. *Pediatrics* 82: 139–46.
- Maruniak-Chudek, I., Ćewietliński, J., 1998. Factors influencing serum lactate concentration in ICU newborns. *Crit. Care* 2: P143.
- Matter, M., Abdel-Hady, H., Attia, G., et al., 2010. Myocardial performance in asphyxiated full-term infants assessed by Doppler tissue imaging. *Pediatr. Cardiol.* 31: 634–642.
- Mir, N.A., Faquih, A.M., Legnain, M., 2010. Perinatal Risk Factors in Birth Asphyxia: Relationship of Obstetric and Neonatal Complications to Neonatal Mortality in 16,365 Consecutive Live Births. *Asia-Oceania J. Obstet. Gynaecol.* 15: 351–357.
- Mongale, P., Andrew, M., 2003. Developmental hemostasis: relevance to newborns and infants. In: Nathan and Oski's Hematology of Infancy and Childhood, Developmental hemostasis: relevance to newborns and infants. In:



Nathan and Oski's Hematology of Infancy and ChildhoodDevelopmental hemo, 6th ed. WB Saunders, Philadelphia.

Morales, P., Bustamante, D., Espina-marchant, P., *et al.*, 2011. Pathophysiology of perinatal asphyxia: can we predict and improve individual outcomes ? *EPMA J.* 2: 211–230.

Munde, A., Kumar, N., Beri, R.S., *et al.*, 2014. Lactate clearance as a marker of mortality in pediatric intensive care unit. *Indian Pediatr.* 51: 565–567.

Nathan, F., 2015. Bersihan laktat sebagai prediktor luaran pada sepsis neonatorum. Universitas Gadjah Mada.

Nguyen, H.B., Loomba, M., Yang, J.J., *et al.*, 2010. Early lactate clearance is associated with biomarkers of inflammation, coagulation, apoptosis, organ dysfunction and mortality in severe sepsis and septic shock. *J. Inflamm. (Lond).* 7: 6.

Padayachee, N., Ballot, D.E., 2013. Outcomes of neonates with perinatal asphyxia at a tertiary academic hospital in Johannesburg, South Africa. *South African J. Child Heal.*

Papile, L.-A., 2001. The Apgar Score in the 21st Century. *N. Engl. J. Med.* 344: 519–520.

Pérez, E.H., Dawood, H., Chetty, U., *et al.*, 2008. Validation of the Accutrend lactate meter for hyperlactatemia screening during antiretroviral therapy in a resource-poor setting. *Int. J. Infect. Dis.* 12: 553–556.

Perlman, J.M., Risser, R., 1995. Cardiopulmonary resuscitation in the delivery room. Associated clinical events. *Arch. Pediatr. Adolesc. Med.* 149: 20–25.

Perlman, J.M., Tack, E.D., 1988. Renal injury in the asphyxiated newborn infant: relationship to neurologic outcome. *J. Pediatr.* 113: 875–879.

Pfenninger, J., Tschaeppler, H., Wagner, B.P., *et al.*, 1991. The paradox of adult respiratory distress syndrome in neonates. *Pediatr. Pulmonol.* 10: 18–24.

Phibbs, C.S., Baker, L.C., Caughey, A.B., *et al.*, 2007. Level and volume of neonatal intensive care and mortality in very-low-birth-weight infants. *N. Engl. J. Med.* 356: 2165–2175.

Phypers, B., Pierce, J.M.T., 2006. Lactate physiology in health and disease. *Contin. Educ. Anaesth. Crit. Care Pain* 6: 128–132.

Pittard, A.J., 1999. Does blood lactate measurement have a role in the management of the critically ill patient? *Ann. Clin. Biochem.* 36 ( Pt 4): 401–407.

Rashkin, M.C., Bosken, C., Baughman, R.P., 1985. Oxygen delivery in critically ill patients. Relationship to blood lactate and survival. *Chest* 87: 580–584.

Rawlins, B., Kim, Y.-M., Haver, J., *et al.*, 2015. Case Study: Experience Applying and Tracking a Quality Improvement Approach for Maternal and Newborn Health Services in Sub-Saharan Africa. *World Health Popul.* 16.

Riikinen, R.S., Kero, P.O., Simell, O.G., 1991. Excitatory amino acids in cerebrospinal fluid in neonatal asphyxia. *Pediatr. Neurol.* 8: 37–40.

Robertson, C., Finer, N., 1985. Term infants with hypoxic-ischemic encephalopathy: outcome at 3.5 years. *Dev. Med. Child Neurol.* 27: 473–484.

Roche, 2012. Accutrend ® Plus User's Manual. Roche Diagnostic.

Roland, E.H., Jan, J.E., Hill, A., *et al.*, 1986. Cortical visual impairment following



- birth asphyxia. *Pediatr. Neurol.* 2: 133–137.
- Royal Prince Alfred Hospital, 2007. RPA Newborn Care Guidelines. available at <http://www.slhd.nsw.gov.au/rpa/neonatal/html/docs/asphyxia.pdf> [cited 6.8.16].
- Seri, I., Evans, J., Tulassay, T., 1998. Renal insufficiency and acute renal failure. In: Avery's Diseases of the Newborn, 7th ed. WB Saunders, Philadelphia.
- Shah, S., Tracy, M., Smyth, J., 2004. Postnatal lactate as an early predictor of short-term outcome after intrapartum asphyxia. *J. Perinatol.* 24: 16–20.
- Shankaran, S., Laptook, A.R., Ehrenkranz, R.A., et al., 2005. Whole-body hypothermia for neonates with hypoxic-ischemic encephalopathy. *N. Engl. J. Med.* 353: 1574–1584.
- Shankaran, S., Woldt, E., Koepke, T., et al., 1991. Acute neonatal morbidity and long-term central nervous system sequelae of perinatal asphyxia in term infants. *Early Hum. Dev.* 25: 135–148.
- Shireen, N., Nahar, N., Mollah, A., 2010. Risk Factors and Short-Term Outcome of Birth Asphyxiated Babies in Dhaka Medical College Hospital. *Bangladesh J. Child Heal.* 33: 83–89.
- Silva, S., Hennebert, N., Denis, R., et al., 2000. Clinical value of a single postnatal lactate measurement after intrapartum asphyxia. *Acta Paediatr.* 89: 320–323.
- Steward, J.E., McCormick, M.C., Leviton, A., 1997. Assessment of Outcomes Following Neonatal Asphyxia, in: Wright, L.L., Merenstein, G.B., Hirtz, D. (Eds.), Acute Perinatal Asphyxia in Term Infants: Report of the Workshop. DIANE Publishing, Maryland, pp. 175–186.
- Surjono, A., 1992. Pencegahan dan pengobatan terhadap infeksi pada neonatus dalam kumpulan makalah reuni dokter spesialis anak ke II FK UGM/RSUP Dr. Sardjito. Yogyakarta.
- Suzuki, S., Morishita, S., 1998. Hypercoagulability and DIC in high-risk infants. *Semin. Thromb. Hemost.* 24: 463–466.
- The Open University, 2014. Labour and delivery care HEAT module: Assessing the degree of asphyxia. available at <http://www.open.edu/openlearnworks/mod/oucontent/view.php?id=275&section=1.5> [cited 6.8.16].
- Valenza, F., Aletti, G., Fossali, T., et al., 2005. Lactate as a marker of energy failure in critically ill patients: hypothesis. *Crit. Care* 9: 588–593.
- Weil, M.H., Afifi, A.A., 1970. Experimental and Clinical Studies on Lactate and Pyruvate as Indicators of the Severity of Acute Circulatory Failure (Shock). *Circulation* 41: 989–1001.
- WHO, 2016. Neonatal mortality. available at [http://www.who.int/gho/child\\_health/mortality/neonatal\\_text/en/](http://www.who.int/gho/child_health/mortality/neonatal_text/en/) [cited 6.3.16].
- WHO, 2012. Maternal and Perinatal Health Profile. *MCA/WHO*. available at [http://www.who.int/maternal\\_child\\_adolescent/epidemiology/profiles/maternal/indn.pdf](http://www.who.int/maternal_child_adolescent/epidemiology/profiles/maternal/indn.pdf) [cited 3.15.16].
- WHO, UNFPA, UNICEF, 2010. AMDD: Monitoring emergency obstetric care: a handbook. Geneva WHO.
- WHO, UNICEF, 2013. Every newborn an action plan to end preventable deaths. available at



[http://www.who.int/maternal\\_child\\_adolescent/topics/newborn/every-newborn-action-plan-draft.pdf](http://www.who.int/maternal_child_adolescent/topics/newborn/every-newborn-action-plan-draft.pdf) [cited 6.1.16].

Wiberg, N., Källén, K., 2016. Fetal scalp blood lactate during second stage of labor: determination of reference values and impact of obstetrical interventions. *J. Matern. Fetal. Neonatal Med.* 1–6.

Wiberg, N., Källén, K., Herbst, A., et al., 2010. Relation between umbilical cord blood pH, base deficit, lactate, 5-minute Apgar score and development of hypoxic ischemic encephalopathy. *Acta Obstet. Gynecol. Scand.* 89: 1263–1269.

Zhang, Z., Xu, X., 2014. Lactate Clearance Is a Useful Biomarker for the Prediction of All-Cause Mortality in Critically Ill Patients: A Systematic Review and Meta-Analysis. *Crit. Care Med.* 42: 1–8.