

REFERENCE

- Achouri, A., Zhang, W. & Shiyang, X., 1998. Enzymatic hydrolysis of soy protein isolate and effect of succinylation on the functional properties of resulting protein hydrolysates. *Food Research International*, 31(9), pp.617–623.
- Adebowale, Y.A. et al., 2007. Isolation, fractionation and characterisation of proteins from Mucuna bean. *Food Chemistry*, 104(1), pp.287–299.
- Adler-Nissen, J., 1979. Determination of the Degree of Hydrolysis of Food protein Hydrolysates by Titrobenzene Sulfonic Acid. *Am. Chem. Soc.*, pp. 1256-1525
- Allen, J.C. & Wrieden, W.L., 1982. Influence of milk proteins on lipid oxidation in aqueous emulsion I. Casein, whey protein and α -lactalbumin. *J. Dairy Res.*, 49, pp.239–248.
- AOAC. 1995. *Official methods of analysis of the association of official analytical chemistry*. 16th edition. AOAC International, Virginia.
- Amza, T., Amadou, I, Zhu. K.X, Zhou, H.M, Amza, T. et al., 2011. Effect of extraction and isolation on physicochemical and functional properties of an underutilized seed protein: Gingerbread plum (*Neocarya macrophylla*). *Food Research International*, 44(9), pp.2843–2850.
- Aurand, L. M., Woods, A.E. & Wells, M.R 1987. *Food Composition and Analysis*. Van Nostrand Reinhold Co. New York
- Sugiyama, K., Egawa, M., Onzuka, H., and Kenkichi Oba, K.Characteristics of sardine muscle hydrolysates prepared by various enzymic treatments. *Nippon Suisan Gakkaishi* 57(3), pp 475-479.
- Berkelman, T & Garcia-Carreno, F., 2005. *Electrophoresis analysis*. In Wrolstat et al.*Handbook of food analytical chemistry*. Pigments, colorants, flavor, texture, and bioactive food components. Willey Interscience, pp 157- 217
- Bamdad, F., Wu, J. & Chen, L., 2011. Effects of enzymatic hydrolysis on molecular structure and antioxidant activity of barley hordein. *Journal of Cereal Science*, 54(1), pp.20–28.
- Barth, A., 2007. Infrared spectroscopy of proteins. *Biochimica et Biophysica Acta - Bioenergetics*, 1767(9), pp.1073–1101.

- Benjakul, S. & Morrissey, M.T., 1997. Protein Hydrolysates from Pacific Whiting Solid Wastes. *Journal of Agricultural and Food Chemistry*, 45(9), pp.3423–3430.
- Broering, J.M. & Bommarius, A.S., 2005. Evaluation of hofmeister effects on the kinetic stability of proteins. *Journal of Physical Chemistry B*, 109(43), pp.20612–20619.
- Bustomi, S., Roswati, T., Sudrajat, R., Leksono, B., Kosasih, A.S., Anggraeni, I., Syamsuwida, Dm., Lisnawati, Y., Mile, Y., Djaenudin, D., Mahfudz & Rachman, E., 2008. Nyamplung (*Calophyllum inophyllum* L.) Sumber biofuel yang potensial. Litbang Kehutanan. Jakarta.
- Chan, W.K.M., Decker, E.A., Lee, J.B., Butterfield, D.A., 1994. EPR Spin-Trapping Studies of the Hydroxyl Radical Scavenging Activity of Carnosine and Related Dipeptides. *Journal of Agricultural and Food Chemistry*, 42, pp.1407–1410.
- Chang, S.K.C. 2003. Protein Analysis. In *Food Analysis* edited by Nielsen, S.S. Plenum Publisher. New York.
- Cheison, S.C., Wang, Z. & Xu, S.Y., 2007. Multivariate strategy in screening of enzymes to be used for whey protein hydrolysis in an enzymatic membrane reactor. *International Dairy Journal*, 17(4), pp.393–402.
- Chen, C., Chi, Y.J., Zhao, M.Y. & Xu, W., 2012. Influence of degree of hydrolysis on functional properties, antioxidant and ACE inhibitory activities of egg white protein hydrolysate. *Food Science and Biotechnology*, 21(1), pp.27–34.
- Chen, C., Chi, Y.J., Zhao, M.Y. & Lv, L., 2012. Purification and identification of antioxidant peptides from egg white protein hydrolysate. *Amino Acids*, 43(1), pp.457–466.
- Chen, H.M. et al., 1996. Antioxidant Activity of Designed Peptides Based on the Antioxidative Peptide Isolated from Digests of a Soybean Protein. *Journal of Agricultural and Food Chemistry*, 44(9), pp.2619–2623.
- Chew, P.G., Casey, A.J. & Johnson, S.K., 2003. Protein quality and physico-functionality of Australian sweet lupin (*Lupinus angustifolius* cv. Gungurru) protein concentrates prepared by isoelectric precipitation or ultrafiltration. *Food Chemistry*, 83(4), pp.575–583.

- Clemente, A., 2001. Enzymatic protein hydrolysates in human nutrition. *Trends in Food Science and Technology*, 11(7), pp.254–262.
- Creighton, T. E. 1993. *Protein: structure and molecular properties*. Freeman.New York
- Damodaran, S., 1996. *Functional Properties Ch 4 Food Protein Properties and Characterization*. S. Nakai and H. W. Modler (Eds). VCH Publisher. Inc.New York.
- Darmawan, R., Bringe, N.A. & de Mejia, E.G., 2010. Antioxidant Capacity of Alcalase Hydrolysates and Protein Profiles of Two Conventional and Seven Low Glycinin Soybean Cultivars. *Plant Foods for Human Nutrition*, 65(3), pp.233–240.
- Decker, E.A. & Crum, A.D., 1991. Inhibition of oxidative rancidity in salted ground pork by carnosine. *Journal of Food Science*, 56(5), pp.1179–1181.
- Demirhan, E., Apar, D.K. & Özbek, B., 2011. Sesame cake protein hydrolysis by alcalase: Effects of process parameters on hydrolysis, solubilisation, and enzyme inactivation. *Korean Journal of Chemical Engineering*, 28(1), pp.195–202.
- Díaz, M. & Decker, E.A., 2004. Antioxidant Mechanisms of Caseinophosphopeptides and Casein Hydrolysates and Their Application in Ground Beef. *Journal of Agricultural and Food Chemistry*, 52(26), pp.8208–8213.
- Elias, R.J., McClements, D.J. & Decker, E.A., 2005. Antioxidant activity of cysteine, tryptophan, and methionine residues in continuous phase ??-lactoglobulin in oil-in-water emulsions. *Journal of Agricultural and Food Chemistry*, 53(26), pp.10248–10253.
- Fabian, C.B., Huynh, L.H. & Ju, Y.H., 2010. Precipitation of rice bran protein using carrageenan and alginate. *LWT - Food Science and Technology*, 43(2), pp.375–379.
- Faraji, H., Decker, E.A. & Aaron, D.K., 1991. Suppression of Lipid Oxidation in Phosphatidylcholine Liposomes and Ground Pork by Spray-Dried Porcine Plasma. *Journal of Agricultural and Food Chemistry*, 39(7), pp.1288–1290.
- Friday, J.B and D.Okano. 2006. *Calophyllum inophyllum* (kamani) Clusiaceae (syn. Guttiferae) (mangosteen family). Species Profiles for Pacific Island Agroforestry www.traditionaltree.org

- González-Tello, P. et al., 1994. Enzymatic hydrolysis of whey proteins: I. Kinetic models. *Biotechnology and bioengineering*, 44(4), pp.523–528.
- Grzonka Z, F. Kasprzykowski & Wiczak, 2007. Cysteine Protease. In *industrial enzymes structure, function and applications*. Springer.
- Haard, N.F., 2001. Conditions of enzymic catalyzed reaction. . In Sikorski, Z.E. *chemical and functional properties of food proteins*. Tehnomic. Publishing Co. inc, Lancaster, p 155-181
- Hernandez, M., Camanas, R. & Alvarezcoque, M., 1990. Determination of total free amino acids with -phthalaldehyde and -acetyl--cysteine. *Microchemical Journal*, 42(3), pp.288–293.
- Hu, F.Y., Chi, C.F., Wang, B., Deng, S.G., 2015. Two novel antioxidant nonapeptides from protein hydrolysate of skate (*Raja porosa*) muscle. *Marine Drugs*, 13(4), pp.1993–2009.
- Huang, G., Ren, L. & Jiang, J., 2011. Purification of a histidine-containing peptide with calcium binding activity from shrimp processing byproducts hydrolysate. *European Food Research and Technology*, 232(2), pp.281–287.
- Illanes, A., 2008. *Enzyme biocatalysis. Principles and applications*. Springer
- Ivashov, V.I, Neklyudov, A.D., Fedorova, N.V. & Khromova, R.A., 1992. Some macrokinetic characteristics of protein proteolysis. *Dokl Akad Nauk* ;322:989–93.
- Jai ganesh, R., Nazeer, R.A. & Sampath Kumar, N.S., 2011. Purification and identification of antioxidant peptide from black pomfret, *Parastromateus niger* (Bloch, 1975) viscera protein hydrolysate. *Food Science and Biotechnology*, 20(4), pp.1087–1094.
- Ju, Z.Y. & Kilara, A., 1998. Textural {Properties} of {Cold}-set {Gels} {Induced} from {Heat}-denatured {Whey} {Protein} {Isolates}. *Journal of Food Science*, 63(2), pp.288–292.
- Karamac, M., Amarowicz, R. & Kostyra, H., 2002. Effect of Temperature and Enzyme / Substrate Ratio on the Hydrolysis of Pea Protein Isolates by Trypsin. *Czech J. Food Sci.*, 20(1), pp.1–6.

- Khorshid, N., Hossain, M.M. & Farid, M.M., 2007. Precipitation of food protein using high pressure carbon dioxide. *Journal of Food Engineering*, 79(4), pp.1214–1220.
- Kim, J.M., Whang, J.H. & Suh, H.J., 2004. Enhancement of angiotensin I converting enzyme inhibitory activity and improvement of the emulsifying and foaming properties of corn gluten hydrolysate using ultrafiltration membranes. *European Food Research and Technology*, 218(2), pp.133–138.
- Kong, B. & Xiong, Y.L., 2006. Antioxidant activity of zein hydrolysates in a liposome system and the possible mode of action. *Journal of Agricultural and Food Chemistry*, 54(16), pp.6059–6068.
- Koolman, J. & Roehm, K.H., 2005. Color Atlas of Biochemistry. Second edition Thieme Stuttgart · New York
- Laidler, K.J., Branko, F., & Peterman, 2009. Temperature Effects in Enzyme Kinetics. In Purich, D.L. *Contemporary Enzyme Kinetics and Mechanism*. Elsevier. USA.
- Leskovacs, V. 2004. *Comprehensive enzyme kinetics*. Kluwer Academic Publishers, New York, pp 192
- Li, B., Chen, F., Ji C., Wu, Y., 2007. Isolation and identification of antioxidative peptides from porcine collagen hydrolysate by consecutive chromatography and electrospray ionization-mass spectrometry. *Food Chemistry*, 102(4), pp.1135–1143.
- Li, H. & Aluko, R.E., 2010. Identification and inhibitory properties of multifunctional peptides from pea protein hydrolysate. *Journal of Agricultural and Food Chemistry*, 58(21), pp.11471–11476.
- Lieske, B. & Konrad, G., 1996. Physico-chemical and functional properties of whey protein as affected by limited papain proteolysis and selective ultrafiltration. *International Dairy Journal*, 6(1), pp.13–31.
- Lowe, G., 1970. The Structure and Mechanism of Action of Papain. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 257(813), pp.237–248.
- Mahmoud, M.I. 1994. Physicochemical and functional properties of protein hydrolysates in nutritional products. *Food technology* 48 (10), pp 89-95

- Mangaroni, A.G. 2003. *Enzyme kinetics. A modern approach*. John Wiley & Son, Inc Publication.
- Ma´rquez , M.C. & Va ´zquez, M.A., 1999. Modeling of enzymatic protein hydrolysis. *Process Biochemistry*, 35(1-2), pp.111–117.
- Marambe, P.W.M.L.H.K., Shand, P.J. & Wanasundara, J.P.D., 2008. An in-vitro investigation of selected biological activities of hydrolysed flaxseed (*Linum usitatissimum* L.) Proteins. *JAOCS, Journal of the American Oil Chemists’ Society*, 85(12), pp.1155–1164.
- Marrufo-Estada, D.M., Segura-Campos, M.R., Chel-Guerrero, L.A., and Betancur-Ancona, D.A., 2013. Defatted *Jatropha curcas* flour and protein isolate as materials for protein hydrolysates with biological activity. *Food Chemistry*, 136, p.77-83.
- Millan, F., Alaiz, M., Hernandez-Pinzon,I., Sanchez , R & Bautista, J., 1994. Study of Lipid Components of *Lupinus mutabilis* Meal and Isolates. *Bioscience, Biotechnology and Biochemistry*, 58(12), pp.2258–2260.
- Mondor,M., Ippersiel, D., Lamarche, F. And Boyee J.I., 2004. Production of Soy Protein Concentrates Using a Combination of Electroacidification and Ultrafiltration. *J.Agric.FoodChem* , 52, pp. 6991–6996
- Naqash, S.Y. & Nazeer, R.A., 2012. Optimization of enzymatic hydrolysis conditions for the production of antioxidant peptides from muscles of *Nemipterus japonicus* and *Exocoetus volitans* using response surface methodology. *Amino Acids*, 43(1), pp.337–345.
- Nazeer, R.A. & Srividhya, T.S., 2011. Antioxidant peptides from the protein hydrolysates of *conus betulinus*. *International Journal of Peptide Research and Therapeutics*, 17(3), pp.231–237.
- Oomah, b.d. & bushuk, w., 1983. Characterization of lupine proteins. *Journal of Food Science*, 48(1), pp.38–41.
- Paredes-Lopez, O., Ordorica-Falomir, C., & Olivares-Vazquez, M.R.,1991. Chickpea protein isolates: Physicochemical, functional and nutritional characterization. *Journal of Food Science*, 56(3), pp.726–729
- Park, J.H., Ye, M. & Park, K., 2005. Biodegradable polymers for microencapsulation of Drugs.pdf. , pp.146–161.

- Powers, J.R., 2003. Application of Enzymes in Food Analysis. In *Food Analysis* edited by Nielsen, S.S dalam Plenum Publisher. New York.
- Putra, M.G.S., 2010. *Isolasi dan karakterisasi isolat bungkil nyamplung (*Calophyllum inophyllum*)*, Tesis, M.Sc., Universitas Gadjah Mada, Yogyakarta
- Qi, M., Hettiarachchy, N.S. & Kalapathy, U., 1997. Solubility and Emulsifying Properties of Soy Protein Isolates Modified by Pancreatin. *Journal of Food Science*, 62(6), pp.1110–1115.
- Rajapakse, N., Mendis, E., Jung, WJ., Je, JY., Kim, SK., 2005. Purification of a radical scavenging peptide from fermented mussel sauce and its antioxidant properties. *Food Research International*, 38(2), pp.175–182.
- Rawlings, N.D, Fraser,R., Morton & Barret, A.L., 2007. An Introduction to Peptidases and The Merops Database. In *Industrial Enzymes Structure, Function and Applications* . Springer.
- Ruan, C.Q., Chi, Y.J. & Zhang, R.D., 2010. Kinetics of hydrolysis of egg white protein by pepsin. *Czech Journal of Food Sciences*, 28(5), pp.355–363.
- Safari, R. et al., 2012. Use of Hydrolysates from Yellowfin Tuna (*Thunnus albacares*) Heads as a Complex Nitrogen Source for Lactic Acid Bacteria. *Food and Bioprocess Technology*, 5(1), pp.73–79.
- Sakanaka, S. & Tachibana, Y., 2006. Active oxygen scavenging activity of egg-yolk protein hydrolysates and their effects on lipid oxidation in beef and tuna homogenates. *Food Chemistry*, 95(2), pp.243–249.
- dos Santos, S.D, Martins, V.M., Salas-Mellado, M. & Prentice, C., 2011. Evaluation of Functional Properties in Protein Hydrolysates from Bluewing Searobin (*Prionotus punctatus*) Obtained with Different Microbial Enzymes. *Food and Bioprocess Technology*, 4(8), pp.1399–1406.
- Lo Scalzo, R., 2008. Organic acids influence on DPPH scavenging by ascorbic acid. *Food Chemistry*, 107(1), pp.40–43.
- Schmidl, M. K., Taylor, S. L., & Nordlee, J. A., 1994. Use of hydrolysate-based products in special medical diets. *Food Technology*, 48, pp 77–85.

- Sgarbieri, V.C. & Galeazzi, M. a, 1978. Some physicochemical and nutritional properties of a sweet lupin (*Lupinus albus* var. multolupa) protein. *Journal of agricultural and food chemistry*, 26(6), pp.1438–42.
- Shahidi, F., Synowiecki, J. & Balejko, J., 1995. Proteolytic hydrolysis of muscle proteins of harp seal (*groenlandica*). *Journal of Agricultural and Food Chemistry*, 42(11), pp.2634–2638.
- Shantha, N.C. & Decker, E.A., 1995. Conjugated linoleic acid concentrations in cooked beef containing antioxidants and hydrogen donors. *Journal of Food Lipids*, 2(1), pp.57–64.
- Shih, F.F. & N.F. Campbell, 1993. *Enzymic modification of soy proteins to improve their functional properties for food use. Food Flavor and Safety*. Washington, D.C American Chemical Society 528, pp 181-191
- Spellman, D., McEvoya, E., O’Cuinnb, G., FitzGerald, R.J., 2003. Proteinase and exopeptidase hydrolysis of whey protein: Comparison of the TNBS, OPA and pH stat methods for quantification of degree of hydrolysis. *International Dairy Journal*, 13(6), pp.447–453.
- Suetsuna, K., Ukeda, H. & Ochi, H., 2000. Isolation and characterization of free radical scavenging activities peptides derived from casein. *The Journal of Nutritional Biochemistry*, 11(3), pp.128–131.
- Sugiyama, K., Egawa, M., Onzuka, H. & Oba, K., 1991. Characteristics of sardine muscle hydrolysates prepared by various enzymic treatments. *Nippon Suisan Gakkaisi* 57, pp. 475-479
- Sun, Q., Shen, H. & Luo, Y., 2011. Antioxidant activity of hydrolysates and peptide fractions derived from porcine hemoglobin. *Journal of Food Science and Technology*, 48(1), pp.53–60.
- Taylor, M.J. & Richardson, T., 1980. Antioxidant activity of skim milk: Effect of heat and resultant sulfhydryl groups. *Journal of Dairy Science*, 63(11), pp.1783–1795.
- Theodorou, L.G., Bieth, J.G. & Papamichael, E.M., 2007. The catalytic mode of cysteine proteinases of papain (C1) family. *Bioresource Technology*, 98(10), pp.1931–1939.

- Tong, L.M., Sasaki, S., McClements, D.J. and Decker, E.A., 2000. Mechanisms of the antioxidant activity of a high molecular weight fraction of whey. *Journal of Agricultural and Food Chemistry*, 48(5), pp.1473–1478.
- Vasileva-Tonkova, E., Nustorova, M. & Gushterova, A., 2007. New protein hydrolysates from collagen wastes used as peptone for bacterial growth. *Current Microbiology*, 54(1), pp.54–57.
- Venkatesan, N. & Rege, D. V., 1973. Nutritional evaluation of the seed proteins of *Calophyllum inophyllum*, Linn. and *Bassia latifolia*. *Sci. Fd. Agric* (24), pp 1317-1323
- Veselá, M. & Friedrich, J., 2009. Amino acid and soluble protein cocktail from waste keratin hydrolysed by a fungal keratinase of *Paecilomyces marquandii*. *Biotechnology and Bioprocess Engineering*, 14(1), pp.84–90.
- Vioquea, J., Sánchez-Vioquea, R., Clementea, A., Pedrochea, J., Bautistab, J. and Francisco Millana, F., 1999. Production and characterization of an extensive rapeseed protein hydrolysate. *Journal of the American Oil Chemists' Society*, 76(7), pp.819–823.
- Wang, JS., Zhao, WM, Zhao QZ. & Jiang, YM., 2007. Antioxidant properties of papain hydrolysates of wheat gluten in different oxidation systems. *Food Chemistry*, 101(4), pp.1658–1663.
- Whitaker, J.R. 2000. Enzyme catalyzed reactions experimental factors that affect rates. In Whitaker, J.r., Vorage, A.G.J., Wong D.W.S. *Handbook of Food Enzymology*. Marcel Dekker, Inc, New York, pp 43- 48
- Wu, H., Chen, H. & Shiau, C., 2003. Free amino acids and peptides as related to antioxidant properties in protein hydrolysates of mackerel (*Scomber austriasicus*). *Food Research International*, 36(9-10), pp.949–957.
- Wu, W.U., Hettiarachchy, N.S. & Qi, M., 1998. Hydrophobicity, solubility, and emulsifying properties of soy protein peptides prepared by papain modification and ultrafiltration. *Journal of the American Oil Chemists' Society*, 75, pp.845–850.
- Zayas, J. F., 1997. *Functionality of Protein in Food*. Springer Pub. Jerman.
- Zhang, J., Zhang, H., Wang, L., Guo X., Wang X. & Yao, H., 2009. Antioxidant activities of the rice endosperm protein hydrolysate: Identification of the active peptide. *European Food Research and Technology*, 229(4), pp.709–719.

- Zhu, K., Zhou, H. & Qian, H., 2006. Antioxidant and free radical-scavenging activities of wheat germ protein hydrolysates (WGPH) prepared with alcalase. *Process Biochemistry*, 41(6), pp.1296–1302.
- Zieliński, H. & Kozłowska, H., 2000. Antioxidant activity and total phenolics in selected cereal grains and their different morphological fractions. *Journal of Agricultural and Food Chemistry*, 48(6), pp.2008–2016