

## DAFTAR PUSTAKA

- Akaike, H. 1974. A new look at the statistical model identification. *IEEE Trans. Autom. Control* 19 (6), 716-723.
- Ali, M.A., G. Gyulai., N.Hidvégi., B. Kerti., F.M.A. Al Hemaïd., A.K. Pandey and J. Lee. 2014. The changing epitome of species identification – DNA barcoding. *Saudi Journal of Biological Sciences* 21:204-231.
- Altschul, S.F., W. Gish., W. Miller., E.W. Myers and D.J. Lipman. 1990. Basic local alignment search tool. *Journal of Molecular Biology* 215(3): 403-410.
- Anonim. 2012. *National Strategy and Action Plan: Indonesia 2012 – 2015*. BAPPENAS by Mangroves for the Future. Jakarta. pp: 12-13.
- Anonim. 2004. *Seagrass in the South China Sea: Reversing Environmental Degradation Trends in the South China Sea and Gulf of Thailand*. UNEP/GEF Regional Working Group on Seagrass. Bangkok, Thailand.
- CBOL Plant Working Group. 2009. A DNA barcode for land plants. *Proceeding of National Academy of Sciences* 106(31): 12794-12797.
- Cesar, H. 1996. *Economic Analysis of Indonesia Coral Reefs*. Report Progress: toward environmentally and socially sustainable development. Environment Department.
- Chase, M.W., R.S. Cowan., P.M. Hollingsworth., C. van der Berg., S. Madrinan., G. Petersen., O. Seberg., T. Jørgensen., K.M. Cameron., M. Carine., N. Pedersen., T.A.J. Hedderson., F. Conrad., G.A. Salazar., J.E. Richardson., M.L. Hollingsworth., T.G. Barraclough., L. Kelly and M. Wilkinson. 2007. A proposal for standarised protocol to barcode all land plants. *Taxon* 56:295-299.
- Chen, L.Y., J.M. Chen., R.W. Gituru and Q.F. Wang. 2012. Generic phylogeny, historical biogeography and character evolution of the cosmopolitan aquatic plant family Hydrocharitaceae. *BMC Evolutionary Biology* 12 article 30.
- Christanty, L., M. K. Moosa, Soekarno and M. Abrar. 2008. *Ekosistem Pesisir dan Laut: Potensi dan Pemanfaatan Ramah Lingkungan*. COREMAP-LIPI. Jakarta.
- Çiftci, Y., and I. Okamus. 2003. Fish Population Genetics and Molecular Markers : II- Molecular Markers and Their Applications in Fisheries and Aquaculture. *Turkish Journal of Fisheries and Aquatic Sciences*. 3:51–79.
- Coyer, J. A., G. Hoarau., J. Kuo., A. Tronholm., J. Veldsink and J.L. Olsen. 2013. Phylogeny and temporal divergence of the seagrass family Zosteraceae using one nuclear and three chloroplast loci. *Systematics and Biodiversity* 1-14 pp.
- De Vere, N., T.C.G. Rich. C.R. Ford., S.A. Trinder., C. Long., C.M. Moore., D. Satterthwaite., H. Davies., J. Allainguillaume., S. Ronca., T. Tatarinova., H. Garbett., K. Walker and M.J. Wilkinson. 2012. DNA barcoding the native flowering plants and conifers of Wales. *PLoS ONE* 7: e37945.
- Drummond, A.J. & A. Rambaut. 2007. BEAST: Bayesian evolutionary analysis by sampling tree. *BMC Evolutionary Biology*. 7:214.

- Fadholi, H.P., A.P. Nugroho, M. Sarto. 2014. Distribusi dan akumulasi merkuri (Hg) pada kerang bulu (*Anadara* sp. di Sekotong, Lombok Barat-NTB. *Proceeding on 4<sup>th</sup> International Conference on Basic Science in conjunction with 5<sup>th</sup> International Conference on Global Resource Conservation Science*. Universitas Negeri Malang.
- Fazekas, A.J., M.L. Kuzmina, S.G. Newmaster and P.M. Hollingsworth. 2012. DNA Barcoding Methods for Land Plants. *In: DNA Barcodes: Methods and Protocols*. Kress, W.J. and D.L Erickson (Eds.) *In: Methods in Molecular Biology* 858: 223-252.
- Fazekas, A.J., K.S. Burgess., P.R. Kesanakurti., S.W. Graham., S.G. Newmaster., B.C. Husband., D.M. Percy., M. Hajibabaei and S.C.H. Barrett. 2008. Multiple multilocus DNA barcodes from the plastid genome discriminate plant species equally well. *PLoS ONE* 3(7): e2802.
- Forest, F., R. Grenyer., M. Rouget., T.J. Davies., R.M. Cowling., D.P. Faith., A. Balmford., J.C. Manning., S.Proches., M. Van der Bank., G. Reeves., T.A. Hedderson and V. Savolainen. 2007. Preserving the evolutionary potential of floras in biodiversity hotspots. *Nature* 445(7129):757-760.
- Göltenboth, F., K.H. Timotius, P.P. Milan, J. Margraf. 2012. *Ekologi Asia Tenggara: Kepulauan Indonesia*. Penerbit Salemba Teknika. Jakarta. pp: 31-46.
- Groombridge, B. and M.D. Jenkins. 2002. *World Atlas of Biodiversity: Earth's Living Resources in 21<sup>st</sup> Century*. UNEP World Conservation Monitoring Center by University of California Press. London. pp: 134-137.
- Guindon, S. and O. Gascuel. 2003. A Simple, fast and accurate method to estimate large phylogenies by maximum-likelihood. *Systematic Biology* 52: 696-704.
- Hebert, P.D.N., A. Cywinska., S.L. Ball and J.R. deWaard. 2003. Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London. Series B-Biological Sciences* 270: 313-321.
- Hollingsworth, P.M. 2011. Refining the DNA barcode for land plants. *Proceeding of National Academy of Sciences* 108(49): 19451-19452.
- Hollingsworth, M.L., A.A. Clark., L.L. Forest., J. Richardson., R.T. Pennington., D.G. Long., R. Cowan., M. W. Chase., M. Gaudeul and P.M. Hollingsworth. 2009. Selecting barcoding loci for plants: evaluation of seven candidate loci with species-level sampling in three divergent groups of land plants. *Molecular Ecology Resources* 9:439-457.
- Hurst, G.D.D. and F.M. Jiggins. 2005. Problems with mitochondrial DNA as a marker in population, phylogeographic and phylogenetic studies: the effects of inherited symbionts. *Proceedings. Biological Sciences / The Royal Society*. 272, 1525–1534.
- Iles, W.J.D., S.Y. Smith and S.W. Graham. Refining our understanding of the phylogenetic backbone of Alismatales. *In: Early Events in Monocot Evolution*. Wilkin, P. and S.J. Mayo (Eds.). Systematics Association special volume. Cambridge University Press. Sequence data submitted to GenBank on 12 Jan 2011. Unpublished paper.

- Ito, Y. and Tanaka, N. 2011. Hybridisation in a tropical seagrass genus *Halodule* (Cymodoceaceae), inferred from plastid and nuclear DNA phylogenies. *Telopea* 13(1-2): 219-231.
- Ito, Y., T.O. Toma., J. Murata and N. Tanaka. 2010. Hybridization and polyploidy of an aquatic plant, *Ruppia* (Ruppiaceae), inferred from plastid and nuclear DNA phylogenies. *American Journal of Botany* 97(7):1156-1167.
- Kato, Y., K. Aioi., Y. Omori., N. Takahata and Y. Satta. 2003. Phylogenetic analyses of *Zostera* species based in *rbcL* and *matK* nucleotide sequences: Implications for the origin and diversification of seagrasses in Japanese waters. *Genes and Genetic Systems* 78(5): 329-342.
- Kearse, M., R.Moir., A. Wilson., S. Stones-Havas., M. Cheung., S. Sturrock., S. Buxton., A. Copper., S. Markowitz., C. Duran., T. Thierer., B. Ashton., P. Meintjes and A. Drummond. 2012. Geneious basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28 (12): 1647-1649.
- Kim, J.S. and J.H. Kim. 2013. Comparative genome analysis and phylogenetic relationship of order Liliales insight from the complete plastid genome sequences of two lilies (*Lilium longiflorum* and *Alstroemeria aurea*). *PLoS ONE* 8(6): e68180.
- Kress, J.W. and D.L. Erickson. 2007. A two-locus global DNA barcode for land plants: the coding *rbcL* gene complements the non-coding *trnH-psbA* spacer region. *PLoS ONE* 2(6): e508.
- Kress, J.W., K.J. Wurdack., E.A. Zimmer., L.A. Weigt and D.H. Janzen. 2005. Use of DNA barcodes to identify flowering plants. *Proceeding of National Academy of Sciences* 102(23): 8369-8374.
- Kuriandewa, T.E., W. Kiswara., H. Malikusworo and S. Soemodihardo. 2003. In: *World Atlas of Seagrass*. Green, E.P. and F.T. Short (Eds.). UNEP World Conservation Monitoring Center by University of California Press. London. pp: 171-183.
- Krebs, C.J. 2009. *Ecology: The Experimental Analysis of Distribution and Abundance*. 6<sup>th</sup> edition. Benjamin Cummings. USA.
- Kuo, J. 2007. New monocious seagrass of *Halophila sulawesii* (Hydrocharitaceae) from Indonesia. *Aquatic botany* 87: 171-175.
- Kuo, J. and C. den Hartog. 2001. Seagrass Taxonomy and Identification Key. in: Short, F.T., R.G. Coles & C.A. Short (Eds). *Global Seagrass Research Method*. Elsevier Science B.V. Amsterdam. Netherlands. pp: 31-58.
- Lahaye, R., M. van der Bank., D. Bogarin., J. Warner and F. Pupulin. 2008a. DNA barcoding the floras of biodiversity hotspots. *Proceedings of the National Academy of Sciences* 105: 2923-2928.
- Lahaye, R., V. Savolainen., S. Duthoit., O. Maurin., M. van der Bank. 2008b. A test of *psbK-psbI* and *atpF-atpH* as potential plant DNA barcodes using the flora of Kruger National Park (South Africa) as a model system. *Nature Proceedings*. ([hdl.handle.net/10101/npre.2008.1896.1](http://hdl.handle.net/10101/npre.2008.1896.1))

- Les, D.H., M.L. Moody., W.L Jacobs and R.J. Bayer. 2002. Systematic of seagrasses (Zosteraceae) in Australia and New Zealand. *Systematic Botany* 27(3): 468-484.
- Les, D.H., M.A. Cleland and M. Waycott. 1997. Phylogenetic studies in Alismatidae II: evolution of marine angiosperms (seagrasses) and hydrophyly. *Systematic Botany* 22(3): 443-463.
- Les, D.H., D.K. Garvin and C.F. Wimpee. 1993. Phylogenetic studies in the monocot subclass Alismatidae: evidence for a reappraisal of the aquatic order Najales. *Molecular Phylogenetic Evolution* 2(4): 304-314.
- Li, F.W., L.Y. Kuo., C.J. Rothfels., A. Ebihara., W.L. Chiou., M.D. Windham and K.M. Pryer. 2011. *rbcl* and *matK* earn two thumbs up as the core DNA barcode for ferns. *PLoS ONE* 6(10): e26597.
- Li, X., and Z. Zhou. 2009. Phylogenetic studies of the core Alismatales inferred from morphology and *rbcl* sequences. *Progress in Natural Science* 19(8): 931-945.
- Li, X.X. and Z.K. Zhou. Monocotyledons phylogeny based on three genes (*matK*, *rbcl* and 18S rDNA) sequences. Taxonomical and Ethnobotanical Department, Kunming Institute of Botany, The Chinese Academy of Sciences. Heilongtan, Kunming, Yunnan, China. Data submitted to GenBank on 3 Mar 2005. Unpublished paper.
- Lucas, C., T. Thangaradjou & J. Papenbrock. 2012. Development of a DNA barcoding system for seagrasses: Successful but not simple. *PLoS ONE* 7(1): e29987.
- Maabuat, P.V., J. Sampekalo & H.E.I. Simbala. 2012. Keanekaragaman lamun di pesisir pantai Molas, kecamatan Bunaken kota Manado. *Jurnal Bioslogos* 2(1): 20-27.
- McKenzie, L.J., S.J. Campbell & C.A. Roder. 2003. *Seagrass Watch: Manual for Mapping & Monitoring Seagrass Resources by Community (Citizen) Volunteers*. Northern Fisheries Center, Department of Primary Industries. Cairns. Townsville. pp: 11-14.
- Mount, D.W. 2008. Choosing a Method for Phylogenetic Prediction. In *Cold Spring Harbor Protocols*. Cold Spring Harbor Laboratory Press, pp .1-3.
- Newmaster, S.G., A.J. Fazekas., R.A.D. Steeves and J. Janovec. 2008. Testing candidate plant barcode regions in the Myristicaceae. *Molecular Ecology Notes* 8: 480-490.
- Olsen, J.L., W.T. Stam., J.A. Coyer., T.B.H. Reusch., M. Billingham., C. Bostrom., E. Calvert., H. Christie., S. Granger., R. la Lumiere., N. Milchakova., M.P.O le Secq., G. Procaccini., B. Sanjabi., E. Serrao., J. Vedsink., S. Widdicombe & S.W. Echeverria. 2004. North Atlantic phylogeography and large-scale population differentiation of the seagrass *Zostera marina* L. *Molecular Ecology* 13(7): 1923-1941.
- Osathanukul, M., S. Chomdej., A. Jampeetong., K. Buddhachat., S. Singtonat and S. Ounjai. Traditional approach combining with DNA and anatomy data for the identification of two variants of *Halophila ovalis* seagrass found in

- Thailand. Biology Chiang Mai University. Thailand. Data submitted to GenBank on 12 Jul 2012. Unpublished paper.
- Papenbrock, J. 2012. Highlights in seagrasses phylogeny, physiology and metabolism: What makes them special?. *International Scholarly Research Network Botany* 1-15 pp.
- Petersen, G., O. Seberg., J.I. Davis and Stevenson. 2006. RNA editing and phylogenetic reconstruction in two monocot mitochondrial genes. *Taxon* 55(4):871-886.
- Pennisi, E. 2007. Taxonomy wanted: a barcode for plants. *Science* 318:190-191.
- Posada, D. 2008. In Press, jModelTest: Phylogenetic Model Averaging. *Molecular Biology and Evolution*.
- Procaccini, G., L. Mazzella., R.S. Alberte and D.H. Les. 1999. Chloroplast tRNA (Leu) (UAA) intron sequences provide phylogenetic resolution of seagrass relationships. *Aquatic Botany* 62(4): 269-283.
- QIAGEN. 2001. *DNeasy® Plant Mini Kit and DNeasy Plant Maxi Kit Handbook for DNA Isolation from Plant Tissue*. QIAGEN.
- Rambaut, A. 2009. FigTree. Tree Figure Drawing Tool Version 1.3.1. Institute of Evolutionary Biology, University of Edinburgh.
- Reid, C., J. Marshall., D. Logan & D. Kleine. 2011. *Terumbu Karang dan Perubahan Iklim*. (terjemahan dari: *Coral Reefs and Climate Change: The Guide for Education and Awareness*). Coral Watch, The University of Queensland. Brisbane-Australia.
- Ronquist, F., M. Teslenko., P.V.D. Mark., D.L. Ayres., A. Darling., A. Höhna., B. Larget., L. Liu., M.A. Suchard and J.P. Huelsenbeck. MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst. Biol* 61(3): 539-542.
- Saarela, J.M., P.C. Sokoloff., L.J. Gillespie., L.L. Consaul and R.D. Bull. 2013. DNA barcoding the Canadian arctic flora: core plastid barcodes (*rbcL* + *matK*) for 490 vascular plant species. *PLoS ONE* 8(10): e77982.
- Seagrass Watch. Tanpa tahun. *Indonesia Seagrass*. [www.seagrasswatch.org/indonesia.html](http://www.seagrasswatch.org/indonesia.html). [Web]. Akses tanggal 6 Maret 2014.
- Selvaraj, D., J.I. Park., M.Y. Chung., Y.G. Cho., S. Ramalingam & I.S. Nou. 2013. Utility of DNA barcoding for plant biodiversity conservation. *Plant Breeding Biotechnology* 1(4): 320-332.
- Shalihah, Q., M. Nuhamunada., S. Rahmawati., I. Riyanto., W. Pratama, R.E. Ranis., A.K. Pawiro., M. Ulfa and Y.H. Pratiwi. 2012. Biodiversity and percent coverage of seagrass in Gili Genting beach, West Lombok, Indonesia. *Poster presentation on LIPI-JSPS Joint International Seminar*.
- Short, F.T., G.E. Moore and K.A. Peyton. 2010. *Halophila ovalis* in the tropical Atlantic ocean. *Aquatic Botany* 93(3): 141-146.
- Short, F., T. Carruthers., W. Denisson and M. Waycott. 2007. Global seagrass distribution and diversity: A bioregional model. *Journal of Experimental Marine Biology and Ecology* 350: 3-20.

- Supriyadi, I.H. 2010. Pemetaan padang lamun di perairan teluk Toli-toli dan pulau sekitarnya, Sulawesi Barat. *Oseanologi dan Limnologi di Indonesia* 36(2): 147-164.
- Tamura, K., D. Peterson., N. Peterson., G. Stecher., M. Nei and S. Kumar. 2011. MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance and maximum parsimony methods. *Mol. Biol. Evol* 9 pp. doi: 10.1093/molbev/msr121.
- Tanaka, N., H. Setoguchi and J. Murata. 1997. Phylogeny of the family Hydrocharitaceae inferred from *rbcl* and *matK* gene sequence data. *Journal of Plant Research* 110(1099): 329-337.
- Tomlinson, P.B. and U. Posluzny. 2001. Generic limits in the seagrass family Zosteraceae. *Taxon* 50: 429-437.
- Uchimura, M., E.J. Faye., S. Shimada., T. Inoue and Y. Nakamura. 2008. A reassessment of *Halophila* species (Hydrocharitaceae) diversity with special reference to Japanese representatives. *Botanica Marina* 51(4): 258-268.
- Vijayan, K. and C.H. Tsou. 2010. DNA barcoding in plants: taxonomy in a new perspective. *Current Science* 99(11): 1530-1541.
- Waycott, M., D.W. Freshwater., R.A. York., A. Calladine and W.J. Kenworthy. 2002. Evolutionary trends in the seagrass genus *Halophila* (Thouars): insight from molecular phylogeny. *Bulletin of Marine Science* 71(3): 1299-1308.
- Waycott, M., K. McMahon, J. Mellors., A. Calladine and D.Kleine. 2004. *A Guide to Tropical Seagrasses of the Indo-West Pacific*. James Cook University. Townsville, Australia. 72 pp.
- Yaakub, S.M., R.L.F. Lim., W.L. Lim & P.A. Todd. 2013. The diversity and distribution of seagrass in Singapore. *Nature in Singapore* 6:105-111.