

## DAFTAR PUSTAKA

- Annison, G., & Topping, D. L. (1994). Nutritional role of resistant starch: chemical structure vs physiological function. *Annual Review of Nutrition*, 14: 297–320.
- Anonim. (1999). *Definition, Diagnosis and Classification of Diabetes Mellitus and its Complications, Report of a WHO Consultation, Part 1: Diagnosis and Classification of Diabetes Mellitus*. World Health Organization Department of Noncommunicable Disease Surveillance, Geneva.
- Anonim. (2011). *Feed Conversion Ratio (FCR)*. Technical Bulletin 07, USAID.
- Anonim. (2012). Diagnosis and Classification of Diabetes Mellitus. *Journal of Diabetes Care*, 35, 1: S64-S71.
- Anonim<sup>a</sup>. (2013). *Riset Kesehatan Dasar (Riskesdas) 2013*. Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan Republik Indonesia, Jakarta.
- Anonim<sup>b</sup>. (2013). Konsumsi Rata-rata per Kapita Setahun Beberapa Bahan Makanan di Indonesia, 2009-2013. <http://www.deptan.go.id/Indikator/tabe-15b-konsumsi-rata.pdf>. [20 Maret 2014].
- Anonim<sup>a</sup>. (2014). *Situasi dan Analisis Diabetes*. Infodatin, Pusat Data dan Informasi Kementerian Kesehatan Republik Indonesia, Jakarta.
- Anonim<sup>b</sup>. (2014). Risk factors Diabetes, International Diabetes Federation. <http://www.idf.org/about-diabetes/risk-factor/.html>. [21 Mei 2015].
- Anwer, T. (2014). Melatonin Ameliorates Hyperinsulinemia, Glucose Intolerance And Insulin Resistance In STZ-Nicotinamide Induced Type 2 Diabetic Rats. *International Journal of Pharmacy and Pharmaceutical Sciences*, 6, 2: 133-136.
- AOAC. (1995). *Official of Analysis of The Association of Official Analytical Chemistry*. AOAC Inc, Airlington.
- Arcila, J.A., & Rose, D.J. (2015). Repeated cooking and freezing of whole wheat flour increases resistant starch with beneficial impacts on in vitro fecal fermentation properties. *Journal of functional foods*, 12: 230–236.
- Argasasmita, T.U. (2008). Karakterisasi Sifat Fisikokimia dan Indeks Glikemik Varietas Beras Beramilosa Rendah dan Tinggi. *Skripsi*. Fakultas Teknologi Pertanian IPB, Bogor.
- Asp, N.G., Johansson, C.G., Hallmer, H., & Siljestrom, M. (1983). Rapid Enzymatic Assay of Insoluble and Soluble Dietary Fiber. *Journal. Agric. Food Chem.*, 31: 476-482.
- Banerjee, M., & Vats, P. (2014). Reactive metabolites and antioxidant gene polymorphisms in Type 2 diabetes mellitus. *Journal Redox Biology*, 2: 170–177.

- Benzie, I., & Strain, J. (1996). The Ferric Reducing Ability of Plasma (FRAP) as a Measure of “Antioxidant Power”: The FRAP Assay. *Analytical Biochemistry*, 239: 70–76.
- Bird, A. R., Brown, I. L., and & Topping, D. L. (2000). Starches, Resistant Starches, the Gut Microflora and Human Health. *Journal Curr. Issues Intest. Microbiol*, 1(1): 25-37.
- Brouns, F., Kettlitz, B., & Arrigoni, E. (2002). Resistant starch and “the butyrate revolution”. *Journal Trends in Food Science & Technology*, 13: 251–261.
- Denardin, C. C., Walter, M., Silva, L. P. D., Souto, G. D., & Fagundes, C. A.A. (2007). Effect of amylose content of rice varieties on glycemic metabolism and biological responses in rats. *Journal Food Chemistry*, 105: 1474–1479.
- Eprintsev, A. T., Shevchenko, Y.M., & Popov, V.N. (2008). Carbohydrate Metabolism in the Liver of Rats in Food Deprivation and Experimental Diabetes. *Biology Bulletin*, 35 (1): 99–101.
- Everard, A., & Cani, P.D. (2013). Diabetes, obesity and gut microbiota. *Best Practice & Research Clinical Gastroenterology*, 27: 73–83.
- Frei, M., Siddhuraju, P., & Becker, K. (2003). Studies on in vitro starch digestibility and the glycemic index of six different indigenous rice cultivars from the Philippines. *Journal Food Chemistry*, 83: 395–402.
- Galuh, A., Aprilia P, F., Kohartono, G., Novia, J., Widyawati, P.S., Suteja, A.M., & Suseno, T.I.P. (2013). Perbedaan Kandungan Senyawa Bioaktif dan Aktivitas Antioksidan Tepung Beras Organik Varietas Lokal (Putih Varietas Cianjur, Merah Varietas Saodah, dan Beras Hitam Varietas Jawa). *Prosiding Seminar Nasional: Menggagas Kebangkitan Komoditas Unggulan Lokal Pertanian dan Kelautan, Fakultas Pertanian Universtas Trunojoyo Madura*: 801-809.
- Gandhi, G.R., Ignacimuthu, S., & Paulraj, M.G. (2011). Solanum torvum Swartz. fruit containing phenolic compounds shows antidiabetic and antioxidant effects in streptozotocin induced diabetic rats. *Journal Food and Chemical Toxicology*, 49: 2725–2733.
- Gellar, L., & Nansel, T.R. (2009). High and Low Glycemic Index Mixed Meals and Blood Glucose in Youth with Type 2 Diabetes or Impaired Glucose Tolerance. *Journal of Pediatrics*, 154:455-458.
- Gibson, G.R., & Roberfroid, M.B. (1995). Dietary Modulation of the Human Colonie Microbiota: Introducing the Concept of Prebiotics. *Journal American Institute of Nutrition*: 1401-1412.
- Gibney, M., Ian, A., Helen, M. (2003). *Nutrition and Metabolism*. Blackwell Publishing, United Kingdom.
- Goni, I., Garcia-Diz, L., Manas, E., & Saura-Calixto, F. (1996). Analysis of resistant starch: a method for foods and food products. *Journal Food Chemistry*, 56: 445449.

- Haenen D., Zhang, J., Souza, D.S.C., Bosch, G., Van, D.M.I.M., Van, A.J., Van, D.B.J.J., Perez, G.O., Smidt, H., Kemp, B.M.M., Hooiveld, G.J. (2013). A diet high in resistant starch modulates microbiota composition, SCFA concentrations, and gene expression in pig intestine. *Journal Nutrition*, 143 (3): 274-283.
- Harini, S., R., Roekistiningsih, & Rahmi, Y. (2010). Perbedaan Nilai Indeks Glikemik Beras Hitam (*Oryza sativa L. Indica*), Beras Merah (*Oryza nivara*), dan Beras Putih (*Oryza sativa*). Program Studi Ilmu Gizi FKUB, Malang.
- Harish, K., Varghese, T. (2006). Probiotics in humans—evidence based review. *Calicut Medical Journal*, 4 (4):1-11.
- Herawati, E.R.N., Santosa, U., Astuti, M., Suharwaadji, & Ariani, D. (2013). Pengaruh Konsumsi Ekstrak Antosianin Ubi Jalar Ungu (*Ipomoea batatas L.*) terhadap Glukosa Darah, dan Produk Oksidasi Lipid (Malonaldehyde) Tikus Hiperglikemia Induksi Aloksan. Prosiding Seminar Nasional Konsumsi Pangan Sehat dengan Gizi Seimbang Menuju Tubuh Sehat Bebas Penyakit: 191-197. Gadjah Mada University Press, Yogyakarta.
- Hijova, E., Chmelarova A. (2007). Short Chain Fatty Acids and Colonic Health. *Journal Bratisl Lek Listy*, 108, 8: 354-358.
- Hu, P., Zhao, H., Duan, Z., Linlin, Z., & Wu, D. (2004). Starch digestibility and the estimated glycemic score of different types of rice differing in amylose contents. *Journal of Cereal Science*, 40: 231–237.
- Hu, E.A., Pan, An., Malik, V., & Sun, Q. (2012). White rice consumption and risk of type 2 diabetes: meta-analysis and systematic review. *BMJ*, 344:e1454.
- Irawan, M.A. (2007). Glukosa & Metabolisme Energi. *Jurnal Polton Sports Science & Performance Lab*, 1 (6): 1-5.
- Johnston, K., Sharp, P., Clifford, M., & Morgan, L. (2005). Dietary polyphenols decrease glucose uptake by human intestinal Caco-2 cells. *FEBS Letters*, 579: 1653–1657.
- Juliano, B.O. (1972). *The rice caryopsis and its composition*. Di dalam D.F. Houston (ed). *Rice Chemistry and Technology*. American Association of Chemists, Inc..St. Paul. Minnesona, pp.
- Jung, E.H., Kim, S.R., Hwang, I.K., & Ha, T.Y. (2007). Hypoglycemic Effects of a Phenolic Acid Fraction of Rice Bran and Ferulic Acid in C57BL/KsJ-db/db Mice. *Journal Agric. Food Chem*, 55: 9800–9804.
- Karam, J.H., Patricia, P.R., Salber, and Forsham, P.H. (1996). *Pancreatic Hormones and Diabetes Mellitus*, In Greenspan, F.S., *Basic and Clinical Endocrinology*, 3rd Ed, 593-649, Prentice-Hall International Inc., London.

- Kim, W. K., Chung, M. K., Kang, N. E., Kim, M. H., & Park O. J. (2003). Effect of resistant starch from corn or rice on glucose control, colonic events, and blood lipid concentrations in streptozotocin-induced diabetic rats. *Journal of Nutritional Biochemistry*, 14: 166–172.
- Kleessen, B., Stoof G., Proll, J., Schmiedl D., Noack J., & Blaut, M. (1997). Feeding resistant starch affects fecal and cecal microflora and short-chain fatty acids in rats. *Journal Animal Science*, 75:2453-2462.
- Kwak, J. H., Paik, J. K., Kim, H. I., Kim, O. Y., Shin, D. Y., Kim, H., Lee, J H., & Lee, J. H. (2012). Dietary treatment with rice containing resistant starch improves markers of endothelial function with reduction of postprandial blood glucose and oxidative stress in patients with prediabetes or newly diagnosed type 2 diabetes. *Journal Atherosclerosis*, 224: 457-464.
- Lai, M.W.K., Lee, Y., Chan, C.S., Sung, IK., Wong, W.K., Wong, K.W., Yip, T.P., & Siu, S.C. (2014). Effect Of Low Glycaemic Index Diet On Blood Glucose In Chinese Type 2 Diabetic Patients: Randomized Controlled Trial. *Posters Diabetes Research and Clinical Practice*, 106S1: S69.
- Larsen N., Vogensen F.K., van den Berg F.W.J., Nielsen D.S., Andreasen A.S., Pedersen B. K., Al-Soud, W.A., . Sørensen, S.J., Hansen, L.H., & Jakobsen, M. (2010). Gut Microbiota in Human Adults with Type 2 Diabetes Differs from Non-Diabetic Adults. *Journal PLoS ONE*, 5(2): e9085.
- Lehninger, A.L. (1982). *Dasar-dasar Biokimia Jilid 2*. (Diterjemahkan oleh Maggy Thenawidjaja). Erlangga, Jakarta.
- Lubijarsih, M.A. (2001). Pengaruh Berbagai Proses Pengolahan terhadap Kadar *Resistent Starch* (RS) dan Indeks Glikemik Uwi (*Dioscorea alata* LINN). *Tesis*. Sekolah Pascasarjana UGM, Yogyakarta.
- Mansour, A., Hosseini, S., Larijani, B., Pajouhi, M., & Mohajeri-Tehrani, M. R. (2013). Nutrients related to GLP1 secretory responses. *Journal Nutrition*, 29: 813–820.
- Marsono, Y., & Topping, D.L. (1993). Complex carbohydrates in Australian rice products-influence of microwave cooking and food processing. *Journal Food Science and Technology*, LWT 26: 364-370.
- Marsono, Y. (2002). Pengaruh Pengolahan terhadap Pati Resisten Pisang Kepok (*Musa paradisiacal fa. typical*) dan Pisang Tanduk (*Musa paradisiacal fa. corniculata*). *Jurnal Agritech*, 22: 56-59.
- Marsono, Y. (2004). Serat Pangan dalam Perspektif Ilmu Gizi. Pidato Pengukuhan Jabatan Guru Besar pada Fakultas Teknologi Pertanian UGM, Yogyakarta.
- Marsono, Y. (2008). Prospek Pengembangan Makanan Fungsional. *Jurnal Teknologi Pangan dan Gizi*, 7 (1): 19-27.
- Marsono, Y., P. Wiyono dan Zaki Utama, (2013). Indeks Glikemik dan Sifat Hipoglikemik Pangan Fungsional untuk Penderita Diabetes, Berbasis Tepung Garut. Dalam Marliyati SA dan Hardiensyah (eds) Prosiding C

Simposium Penelitian Terkini Pangan dan Gizi Bidang Inovasi Produk, Mutu dan Keamanan Pangan. Hlm. 1-11. 27 Juni 2013. PERGIZI PANGAN Indonesia, Jakarta.

- Matsui<sup>a</sup>, T., Ueda, T., Oki, T., Sugita, K., Terahara, N., & Matsumoto, K. (2001).  $\alpha$ -Glucosidase Inhibitory Action of Natural Acylated Anthocyanins. 1. Survey of Natural Pigments with Potent Inhibitory Activity. *Journal Agric. Food Chem.*, 49: 1948-1951.
- Matsui<sup>b</sup>, T., Ueda, T., Oki, T., Sugita, K., Terahara, N., & Matsumoto, K. (2001).  $\alpha$ -Glucosidase Inhibitory Action of Natural Acylated Anthocyanins. 2.  $\alpha$ -Glucosidase Inhibition by Isolated Acylated Anthocyanins. *Journal Agric. Food Chem.*, 49: 1952-1956.
- Matsui, T., Ebuchi, S., Kobayashi, M., Fukui, K., Sugita, K., Terahara, N., & Matsumoto, K. (2002). Anti-hyperglycemic Effect of Diacylated Anthocyanin Derived from *Ipomoea batatas* Cultivar Ayamurasaki Can Be Achieved through the  $\alpha$ -Glucosidase Inhibitory Action. *Journal Agric. Food Chem.*, 50: 7244-7248.
- Melati, S. (2009). Potensi prebiotik Resistant Starch dari Pati Kentang. *Tesis*. Sekolah Pascasarjana UGM, Yogyakarta.
- Nurdyansyah, F. (2014). Karakteristik Mikrobiologis, Fisik dan kimia Digesta Tikus Diabetes Induksi STZ-NA yang diberi Diet Kefir Kombinasi Susu Kambing dan Susu Kedelai. *Tesis*. Sekolah Pascasarjana UGM, Yogyakarta.
- Nurhidajah<sup>a</sup>, Astuti, M., Sardjono, & Murdiati, A. (2004). Evaluasi Sifat fisik, sensori, dan kimia beras merah yang diperkaya kappa-karagenan dan ekstrak antosianin. *Prosiding Seminar Nasional Konsumsi Pangan Sehat dengan Gizi Seimbang Menuju Tubuh Sehat Bebas Penyakit*: 191-197. Gadjah Mada University Press, Yogyakarta.
- Nurhidajah<sup>b</sup> & Nurrahman. (2014). Pemanfaatan Kecambah Beras Merah untuk Perbaikan Kondisi Glikemik Tikus yang Diinduksi STZ-Na. *Laporan akhir*. Penelitian Hibah Bersaing DIPA (Daftar Isian Pelaksanaan Anggaran) Koordinasi Perguruan Tinggi Swasta Wilayah VI Semarang.
- Phillips, J. (1995). Effect resistant starch on fecal bulk and fermentation dependent events in human. *The American Journal of Clinical Nutrition*, 62: 121-130.
- Pramitasari, R. (2014). Potensi Antioksidatif Minuman Fungsional Berbasis Beras Hitam (*Oryza Sativa L. Indica*) dan Kedelai Hitam (*Glycine Max L. Merr.*) untuk Lansia Penyandang Diabetes Mellitus Tipe 2. *Tesis*. Sekolah Pascasarjana UGM, Yogyakarta.
- Puddu, A., Sanguineti, R., Montecucco, F., & Viviani, G.L. (2014). Evidence for the Gut Microbiota Short-Chain Fatty Acids as Key Pathophysiological Molecules Improving Diabetes. *Journal Mediators of Inflammation*: 1-9.

- Purwani, E.Y., Yuliani, S., Indrasari, S.D., Nugraha, S., & Thahir, R. (2007). Sifat Fisiko-Kimia Beras dan Indeks Glikemiknya. *Jurnal Teknologi dan Industri Pangan*, 18 (1): 59-66.
- Purwaningsih, H., & Kristantini. (2009). Menyelamatkan Sumber Daya Genetik Padi Beras Merah. *Warta Plasma Nutraf Indonesia*, 21:4-6.
- Rajilic-Stojanovic, M., & Fellow, R. (2013). Function of the microbiota. *Journal Best Practice & Research Clinical Gastroenterology*, 27: 5–16.
- Ramely, M., Dworzak, S., Hippe, B., Zwieler, J., Aumuller, E., Brath, H., & Haslberger, A. (2013). Abundance and Diversity of Microbiota in Type Diabetes and Obesity. *Journal Diabetes Metabolism*, 4 (3): 1-8.
- Reeves, P.G., F.H. Nielsen, and G.C. Fahey, Jr. (1993). AIN-93 purified diets for laboratory rodents: Final report of the American Institute of Nutrition and ad hoc writing committee on the reformulation of the AIN-76A diet. *Journal Nutr.* 123:1939-1951.
- Rodríguez-Cabezas, M. E., Camuesco, D., Arribas, B., Garrido-Mesa, N., Comalada, M., Bailón, E., Cueto-Sola, M., Utrilla, P., Guerra-Hernández, E., Pérez-Roca, C., Gálvez, J., & Zarzuelo, A. (2010). The combination of fructooligosaccharides and resistant starch shows prebiotic additive effects in rats. *Journal Clinical Nutrition*, 29: 832-839.
- Sakuraia, Masaru., Nakamura, K., Miurab, K., Takamura, T., Yoshitad, K., Morikawaa, Y., Ishizakie, M., Kidof, T., Naruseg, Y., Suwazonoh, Y., Kanekoc, S., Sasaki, S., & Nakagawaa, H. (2012). Dietary glycemic index and risk of type 2 diabetes mellitus in middle-aged Japanese men. *Journal Metabolism*, 61 (1): 47–55.
- Sancho, R. A. S., & Pastore, G. M. (2012). Evaluation of the effects of anthocyanins in type 2 diabetes. *Journal Food Research International*, 46: 378–386 .
- Sharma, A.K., Mohan, P., & Nayak, B.B. (2005). Probiotics: Making a comeback. *Indian Journal Pharmacol*, 37:358-365.
- Shih, C., Chen, S., Hou W., & Cheng H. (2007). A high-resistance-starch rice diet reduces glycosylated haemoglobin levels and improves the antioxidant status in diabetic rats. *Journal Food Research International*, 40: 842–847.
- Shou, Y., Xu, F., Sun, X., Bao, J. & Beta, T. (2014). Identification and quantification of phenolic acids and anthocyanins as antioxidants in bran, embryo, and endosperm of white, red, and black rice kernels (*Oryza sativa* L.). *Journal of Cereal Science*, 56 (2): 211-218.
- Siregar, S.B. (1994). *Ransum Ternak Ruminansia*. Penebar Swadaya, Jakarta.
- Sompong, R., Siebenhandl-Ehn, S., Linsberger-Martin, G., & Berghofer, E. (2011). Physicochemical and antioxidative properties of red and black rice varieties from Thailand, China and Sri Lanka. *Journal Food Chemistry*, 124: 132–140.

- Suardi, D. (2005). Potensi Beras Merah untuk Peningkatan Mutu Pangan. *Jurnal Litbang Pertanian*, 24, 3: 93-100.
- Sudarmadji, S., Bambang Haryono dan Suhardi. (1996). *Analisa Bahan Makanan dan Pertanian*. Liberty, Yogyakarta.
- Suhartatik, N, Karyantina, M., Mustofa, A., Cahyanto, M.N., Raharjo, S., & Rahayu, E.S. (2013). Stabilitas Ekstrak Antosianin Beras Ketan (*Oryza sativa var. glutinosa*) Hitam Selama Proses Pemanasan dan Penyimpanan. *Journal Agritech*, 33 (4): 384-390.
- Szkudelski, T. (2012). Streptozotocin-nicotinamide-induced diabetes in the rat. Characteristics of the experimental model. *Journal Experimental Biology and Medicine*, 237: 481-490.
- Widowati, S., M. Astawan, D. Muchtadi, dan T. Wresdiyati. (2007). Pemanfaatan ekstrak teh hijau (*Camellia sinensis* O. Kuntze) dalam pengembangan beras pratanak fungsional. Prosiding Seminar Nasional PATPI 2007, Bandung 17 18 Juli 2007.
- Widowati, Sri, B.A. Santosa, S., Astawan, M., & Akhyar. (2009). Penurunan Indeks Glikemik Berbagai Varietas Beras melalui Proses Pratanak. *Jurnal Pascapanen*, 6, 1: 1-9.
- Winarno, F.G. (2004). *Kimia Pangan dan Gizi*. PT. Gramedia Pustaka Utama, Jakarta.
- Wu, Y., Chen, Z., Li, X., & Wang, Z. (2010). Retrogradation properties of high amylose rice flour and rice starch by physical modification. *Journal Food Science and Technology*, 43: 492-497.
- Yadav, H., Lee, J.H., Lloyd, J., Walter, P., & Rane, S.G. (2013). Beneficial metabolic effects of a probiotic via butyrate induced GLP-1 secretion. *Manuscript*. The American Society for Biochemistry and Molecular Biology, Inc.
- Yao, Y., Sang, W., Zhou, M., & Ren, G. (2010). Antioxidant and r-Glucosidase Inhibitory Activity of Colored Grains in China. *Journal Agric. Food Chem.*, 58: 770-774.
- Zaupa, M., Calani, L., Rio, D.D., Brighenti, F., & Pellegrini, N. (2015). Characterization of total antioxidant capacity and (poly)phenolic compounds of differently pigmented rice varieties and their changes during domestic cooking. *Journal Food Chemistry*, 187: 338-347.
- Zhou, Z.K., Wanga, F., Rena, X.C., Wanga, Y., & Blanchard, C. (2015). Resistant starch manipulated hyperglycemia/hyperlipidemia and related genes expression in diabetic rats. *Journal of Biological Macromolecules*, 75:316-321.