

DAFTAR PUSTAKA

- Adham M, Kurniawan AN, Muhtadi A, Roezin A, Hermani B, Gondhowiardjo S, et al., (2012). Nasopharyngeal Carcinoma in Indonesia : epidemiology, incidence, signs and symptoms at presentation. *Chin J Cancer* **31**(4).
- Agathangelou, A., G. Niedobitek, R. Chen, J. Nicholls, W. Yin & L. S. Young., (1995). Expression of immune regulatory molecules in Epstein-Barr virus-associated nasopharyngeal carcinomas with prominent lymphoid stroma: evidence for a functional interaction between epithelial tumour cells and infiltrating lymphoid cells. *Am J Pathol* **147**: 1152-1160.
- Akshal S. Patel, Joshua E. Allen, David T. Dicker, Kristi L. Peters, Jonas M. Sheehan, Glantz MJ, et al., (2011). Identification and enumeration of circulating tumor cells in the cerebrospinal fluid of breast cancer patients with central nervous system metastases. *Oncotarget* **2** (10).
- Alatas H, Karyomanggolo WT, Musa DA, Boediarso A, Oesman IN in Sastroasmoro S, Ismail S., (2008). Dasar-dasar Metodologi Penelitian Klinis. CV Sagung Seto. 3rd **Edition**: 92-110.
- Allen, M. D., L. S. Young and C W Dawson, (2005). The Epstein-Barr virus-encoded LMP2A and LMP2B proteins promote epithelial cell spreading and motility. *Journal of Virology* **79**(3): 1789-1802.
- Alix-Panabie C, Pantel K., (2013) Circulating Tumor Cells: Liquid Biopsy of Cancer. Mini-Reviews. *Clinical Chemistry* **59**:1 110–118.
- Altman DG, McShane LM Sauerbrei W, Taube SE. Reporting Recommendations for Tumor Marker Prognostic Studies (REMARK); Explanation and Elaboration. Guidelines and Guidance. Plos Medicine., May 2012, Vol 9, Issue 5
- Altiok, E., J. Minarovits, L. F. Hu, B. Contreras- Brodin, G. Klein & I. Ernberg., (1992). Host-cell-phenotype- dependent control of the BCR2/BWR1 promoter complex regulates the expression of Epstein-Barr virus nuclear antigens 2-6. *Proc Natl Acad Sci USA* **89**: 905- 909.
- Arrand JR., (2000). Expressed but enigmatic RNAs. *EBV Rep* **7**:145 -9.
- Babcock GJ, Hochberg D, Thorley-Lawson AD., (2000). The expression pattern of Epstein-Barr virus latent genes in vivo is dependent upon the differentiation stage of the infected B-cell. *Immunity* **13**:497-506.
- Babcock GJ, Thorley-Lawson DA., (2000). Tonsillar memory B-cells, latently infected

- with Epstein-Barr virus, express the restricted pattern of latent genes previously found only in Epstein-Barr virus-associated tumors. *Proc Natl Acad Sci USA* **97**:12250-5.
- Bates RC, Bellovin DI, Brown C et al. (2005). Transcriptional activation of integrin $\beta 6$ during the epithelial-mesenchymal transition defines a novel prognostic indicator of aggressive colon carcinoma. *Journal of Clinical Investigation* **115** (2): 339–347.
- Bauman, R. W., (2007). *Microbiology with disease by taxonomy*. San Fransisco, Pearson Education Inc. **2nd Edition**: 682-683, 689-691.
- Bayliss GJ, Wolf H., (1980). Epstein-Barr virus-induced cell fusion. *Nature* **287**: 164-5.
- Beck A, Pätzolt D, Grabenbauer GG, Nicholls JM, Herbst H, Young LS, et al., (2001). Expression of cytokine and chemokine genes in Epstein-Barr virus-associated nasopharyngeal carcinoma: comparison with Hodgkin's disease. *J Pathol* **194**:145–51.
- Ben-Sasson, S. A., Klein G., (1981). Activation of the Epstein-Barr virus genome by 5-azacytidine in latently infected human lymphoid cell lines. *Int J Cancer* **28**: 131- 135.
- Bernard, R., (1996). *Herpesviridae : General description, taxonomy and classification*. London, Plenum Press: 23.
- Bestor, T. H., (2000). The DNA methyltransferases of mammals. *Hum Mol Genet* **9**: 2395-2402.
- Brooks LA, Lear AL, Young LA, Rickinson AB., (1993). Transcripts from the Epstein-Barr virus BamHI-A fragment are detectable in all three forms of virus latency. *J Virol* **67**:3182-90.
- Brooks, L., Q. Y. Yao, A. B. Rickinson, et al., (1992). Epstein-Barr virus latent gene transcription in nasopharyngeal carcinoma cells: coexpression of EBNA1, LMP1, and LMP2 transcripts. *Journal of Virology* **66**: 2689-2697.
- Busson P, Braham K, Ganem G, Thomas F, Grausz D, Lipinski M, et al., (1987). Epstein-Barr virus-containing epithelial cells from nasopharyngeal carcinoma produce interleukin. *Proc Natl Acad Sci U S A* **84**:6262–6.
- Brooks PC, Stromblad S, Sanders LC et al., (1996). Localization of matrix metalloproteinase MMP-2 to the surface of invasive cells by interaction with integrin $\alpha v \beta 3$. *Cell* **85**(5): 683–693.
- Busson, P., R. McCoy, R. Sadler, et al., (1992). Consistent transcription of the Epstein-Barr virus LMP2 gene in nasopharyngeal carcinoma. *Journal of Virology* **66**(5): 3257-3262.

- Caldwell RG, Wilson JB, Anderson SJ, Longnecker R., (1998). Epstein -Barr virus LMP2a drives B-cell development and survival in the absence of normal B-cell receptor signals. *Immunity* **9**: 405-11.
- Chan, A. P., P. M. Teo and P. J. Johnson, (2002). Review of nasopharyngeal carcinoma. *Annals of Oncology* **13**: 1007-1015.
- Chan, A. T., Y. M. Lo, B. Zee, L. Y. Chan, B. B. Ma, S.F.Leung, et al., (2002). Plasma Epstein-Barr virus DNA and residual disease after radiotherapy for undifferentiated nasopharyngeal carcinoma. *J Natl Cancer Inst* **94**: 1614- 1619.
- Chan KH, Gu YL, Ng F, Ng PS, Seto WH, Sham JS, et al., (2003). EBV specific antibody-based and DNA-based assays in serologic diagnosis of nasopharyngeal carcinoma. *Int. J. Cancer* **105**: 706–709.
- Chan, S. L., E. Hui, S. F. Leung, et al., (2006). Radiological, pathological and DNA remission in recurrent metastatic nasopharyngeal carcinoma. *BMC Cancer* **6**: 259.
- Chen H, Smith P, Ambinder RF, Hayward SD., (1999). Expression of Epstein-Barr virus BamHI-A rightward transcripts in latently infected B-cells from peripheral blood. *Blood* **93**:3026-32.
- Chen HL, Lung MM, Sham JT, Choy DTK, Griffin BE, Ng MH., (1992). Transcription of the BamHI A region of the EBV genome in NPC tissues and B-cells. *Virology* **191**:193-201.
- Chen, J., (2012). Roles of the PI3K/Akt pathway in Epstein-Barr virus induced cancers and therapeutic implications. *World Journal of Virology* **1**(6): 154-161.
- Cheung, S. T., D. P. Huang, A. B. Hui, K. W. Lo, C. W. Ko, Y. S. Tsang, et al., (1999). Nasopharyngeal carcinoma cell line (C666-1) consistently harbouring Epstein-Barr virus. *Int J Cancer* **83**: 121-126.
- Chiang, A. K., Q. Tao, G. Srivastava & F. C. Ho., (1996). Nasal NK- and T-cell lymphomas share the same type of Epstein-Barr virus latency as nasopharyngeal carcinoma and Hodgkin's disease. *Int J Cancer* **68**: 285-290.
- Chi Man Tsang, Wen Deng, Yim Ling Yip, Mu-Sheng Zeng, Kwok Wai Lo and Sai Wah Tsao., (2014). Epstein-Barr virus infection and persistence in nasopharyngeal epithelial cells. *Clin J Cancer* **33** (11).
- Chindavijak S, Tiwawech D, Sangariyanich E, (2008). “Quantitative Detection of Plasma Cell-free Epstein-Barr Virus DNA in Nasopharyngeal Carcinoma Patients by Real-time PCR.” *Thai Cancer Journal* **28** (1).
- Cho, W., (2007). Nasopharyngeal carcinoma : Molecular biomarker discovery and progress. *Molecular Cancer* **6**(1).

- Christopher W. Dawson, Rebecca J. Port, Lawrence S. Young., (2012). The role of the EBV-encoded latent membrane proteins LMP1 and LMP2 in the pathogenesis of nasopharyngeal carcinoma (NPC). *Seminars in Cancer Biology* **22**: 144–153.
- Chung, J., R. E. Bachelder, E. A. Lipscomb, et al., (2002). Integrin ($\alpha 6\beta 4$) regulation of eIF-4E activity and VEGF translation : a survival mechanism for carcinoma cells. *The Journal Cell of Biology* **158**(1): 165-174.
- Claire, S.-L. and R. Martin, (2011). Epstein-Barr Virus Infection of Polarized Epithelial Cells via the Basolateral Surface by Memory B Cell-Mediated Transfer Infection. *PLoS Pathogens* **7**(5).
- Clemens ML., (1994). Functional significance of the Epstein-Barr virus-encoded small RNAs. *EBV Rep* **1**:107-11.
- Dawson B, Trapp RG., (2004). Basic and Clinical Biostatistics.. Mc Graw Hill. **4th Edition**: 314-316.
- Dawson, C. W., R. J. Port and L. S. Young, (2012). The role of the EBV encoded latent membrane proteins LMP1 and LMP2 in the pathogenesis of nasopharyngeal carcinoma (NPC). *Seminar in Cancer Biology* **22**: 144-153.
- Dawson, C. W., G. Tramountanis, A. G. Eliopoulos, et al., (2003). Epstein-Barr virus latent membrane protein 1 (LMP1) activates the phosphatidylinositol 3-kinase/Akt pathway to promote cell survival and induce actin filament remodeling. *The Journal of Biological Chemistry* **278**(6): 3694-3704.
- Deacon, E. M., G. Pallesen, G. Niedobitek, J. Crocker, L. Brooks, A. B. Rickinson & L. S. Young., (1993). Epstein-Barr virus and Hodgkin's disease: transcriptional analysis of virus latency in the malignant cells. *J Exp Med* **177**: 339-349.
- Decaussin, G., F. Sbih-Lammali, M. de Turenne- Tessier, A. Bouguermouh & T. Ooka., (2000). Expression of BARF1 gene encoded by Epstein-Barr virus in nasopharyngeal carcinoma biopsies. *Cancer Res* **60**: 5584-5588.
- Denzer K, Kleijmeer, Hiljnen HFG, Stoorvogel W, Geuze HJ, (2000). “Exosome : from internal vesicle of the multivesicular body to intercellular signaling device.” *Journal of cell Science* **113**: 3365-3374.
- Desgrosellier JS, Cheresch DA., (2010). Integrins in cancer: biological implications and therapeutic opportunities. *Nature Reviews Cancer* **10**: 890.
- Dutta A Li J, Lu H, Akech J, Pratap J, Wang T et al., (2014). The integrin $\alpha v\beta 6$ promotes metastasis of prostate cancer cells to bone. *IBMS BoneKEy* **11**: 581.
- Dutta A, Jing Li J, Lu H, Akech J, Pratap J, Wang T,et al., (2014). Integrin $\alpha v\beta 6$ Promotes an Osteolytic Program in Cancer Cells by Upregulating MMP2. *American Association for Cancer Research*.

- Ellen T. Chang, Hans-Olov Adami., (2006). The Enigmatic Epidemiology of Nasopharyngeal Carcinoma. *Cancer Epidemiol Biomarkers Prev* **15**(10).
- Ernberg, I., K. Falk, J. Minarovits, P. Busson, T. Tursz, M. G. Masucci & G. Klein., (1989). The role of methylation in the phenotype-dependent modulation of Epstein-Barr nuclear antigen 2 and latent membrane protein genes in cells latently infected with Epstein-Barr virus. *J Gen Virol* **70**: 2989-3002.
- Evi S. Lianidou, Athina Markou. Circulating Tumor Cells in Breast Cancer: Detection Systems, Molecular Characterization, and Future Challenges.
- Fachiroh, J., D. Paramita, B. Hariwiyanto, et al., (2006). Single assay combination of Epstein-Barr Virus EBNA1 and VCA-p18 derived synthetic peptides for measuring EBV-IgG and IgA antibodies in nasopharyngeal carcinoma patients options for field screening. *Journal of Clinical Microbiology* **44**(4): 1459-1467.
- Falasca, M., (2010). PI3K/Akt signalling pathway specific inhibitors: a novel strategy to sensitize cancer cells to anti-cancer drugs. *Current pharmaceutical design* **16**(12): 1410-1416.
- Felding-Habermann B, O'Toole TE, Smith JW et al., (2001). Integrin activation controls metastasis in human breast cancer. *Proceedings of the National Academy of Sciences of the United States of America* **98**(4): 1853–1858.
- Feng, B. J., W. Huang, Y. Y. Shugart, M. K. Lee, F. Zhang, J. C. Xia, H. Y et al., (2002). Genome-wide scan for familial nasopharyngeal carcinoma reveals evidence of linkage to chromosome 4. *Nat Genet* **31**: 395-399.
- Fielding CA, Sandvej K, Mehl A, et al., (2001). Epstein-Barr virus LMP-1 natural sequence variants differ in their potential to activate cellular signalling pathways. *J Virol* **75**:9129-41.
- Fingerroth JD, Diamond ME, Sage DR et al., (1999). CD21-dependent infection of epithelial cell line, 293, by Epstein-Barr virus. *J Virology* **73**: 2115-25.
- Franco R, Cantile M, Marino FZ, Pirozzi G., (2012). Circulating tumor cells as emerging tumor biomarkers in lung cancer. *J Thorac Dis* **4**(5): 438-439.
- Fruehling, S., R. Longnecker., (1997). The immunoreceptor tyrosine-based activation motif of Epstein-Barr virus LMP2A is essential for blocking BCR-mediated signal transduction. *Virology* **235**: 241-251.
- Fruehling, S., R. Swart, K. M. Dolwick, E. Kremmer & R. Longnecker., (1998). Tyrosine 112 of latent membrane protein 2A is essential for protein tyrosine kinase loading and regulation of Epstein-Barr virus latency. *J Virol* **72**: 7796-7806 .
- Fotheringham, J. A., N. E. Coalson and N. Raab-Traub, (2012). Epstein-Barr virus latent membrane protein-2A induces ITAM/Syk- and Akt-dependent epithelial

- migration through α V-integrin membrane translocation. *Journal of Virology* **86**(19): 10308-10320.
- Gao, Y., Y. J. Lu, S. A. Xue, H. Chen, N. Wedderburn & B. E. Griffin., (2002). Hypothesis: a novel route for immortalization of epithelial cells by Epstein-Barr virus. *Oncogene* **21**: 825-835.
- Geminard C, Nault F, Johnstone RM, Vidal M, (2001). "Characteristic of the Interaction between Hsc70 and the Transferrin Receptor in Exosome Released during Reticulocyte Maturation." *The Journal of Biological Chemistry* **276** (12) : 9910-9916.
- Gilman AG., (1987). G proteins: transducers of receptor-generated signals. *Annu. Rev. Biochem.* **56** (1): 615-49.
- Glickman JN, Howe JG, Steitz JA., (1988). Structural analyses of EBER1 and EBER2 ribonucleoprotein particles present in Epstein-Barr virus-infected cells. *J Virol* **62**:902-11.
- Gruhne, B., R. Sompallae and M. G. Masucci, (2009). Three Epstein-Barr virus latency proteins independently promote genomic instability by inducing DNA damage, inhibiting DNA repair and inactivating cell cycle checkpoints. *Oncogene* **28**(45): 3997-4008.
- Gulley, M. L., S. L. Glasser, F. E. Craig, et al., (2002). Guidelines for interpreting EBER in situ hybridization and LMP1 immunohistochemical tests for detecting Epstein-Barr virus in Hodgkin lymphoma. *American Journal of Clinical Pathology* **117**(2): 259-267.
- Habermann BF, O'Toole TE, Smith JW, Franssvea E, Ruggeri ZM, Ginsberg MH, et al., (2001). "Integrin activation controls in human breast cancer." *PNAS* **98**(4).
- Herbst, H., F. Dallenbach, M. Hummel, G. Niedobitek, S. Pileri, N. Muller-Lantzsch & H. Stein., (1991). Epstein-Barr virus latent membrane protein expression in Hodgkin and Reed-Sternberg cells. *Proc Natl Acad Sci U S A* **88**: 4766-4770.
- Heussinger, N., M. Buttner, G. Ott, et al., (2004). Expression of the Epstein-Barr virus (EBV)-encoded latent membrane protein 2A (LMP2A) in EBV-associated nasopharyngeal carcinoma. *The Journal of Pathology* **203**(2): 696-699.
- Higuchi M, Izumi KM, Kieff E., (2001). Epstein-Barr virus latent- infection membrane proteins are palmitoylated and raft-associated: protein 1 binds to the cytoskeleton through TNF receptor cytoplasmic factors. *Proc Natl Acad Sci USA* **98**: 4675-80.
- Hildesheim, A., R. J. Apple, C. J. Chen, S. S. Wang, Y. J. Cheng, W. Klitz, et al., (2002). Association of HLA class I and II alleles and extended haplotypes with nasopharyngeal carcinoma in Taiwan. *J Natl Cancer Inst* **94**: 1780-1789.

- Hino R, Uozaki H, Inoue IY, Shintani Y, Ushiku T, Sakatani T, et al., (2008). Survival Advantage of EBV-Associated Gastric Carcinoma: Survivin Up-regulation by Viral Latent Membrane Protein 2A. *Cancer Res* 2008 **68**(5).
- Hino R, Uozaki H, Murakami N, Ushiku T, Shinozaki A, Ishikawa S, et al., (2009). Activation of DNA Methyltransferase 1 by EBV Latent Membrane Protein 2A Leads to Promoter Hypermethylation of PTEN Gene in Gastric Carcinoma. *Cancer Res* **69**(7).
- Hitt MM, Allday MJ, Hara T, et al., (1989). EBV gene expression in an NPC related tumor. *EMBO J* **8**: 2639-51.
- Ho JHC., (1972). Current knowledge of the epidemiology of nasopharyngeal carcinoma. In: Biggs P, de The G, and Payne L (eds), *Oncogenesis and herpesviruses* : 357- 366.
- Hongchuan, J., M. Yanning, S. Qi, et al., (2012). Circulating Methylated DNA as Biomarkers for Cancer Detection. InTech.
- Hong RL, Lin CY, Ting LL, Ko JY, Hsu MM., (2004). Comparison of clinical and molecular surveillance in patients with advanced naso- pharyngeal carcinoma after primary therapy: the potential role of quantitative analysis of circulating Epstein-Barr virus DNA. *Cancer* **100**:1429–1437.
- Howe JG, Steitz JA., (1986). Localization of Epstein-Barr virus- encoded small RNAs by in situ hybridization. *Proc Natl Acad Sci USA* **83**:9006-10.
- Humphries, M., (2000). Integrin structure. *Biochemical Society Transactions* **28**(4): 311-339.
- Hsiao JR., Jin YT, Tsai ST., (2000). Detection of cell free Epstein- Barr virus DNA in sera from patients with nasopharyngeal carcinoma. *Cancer* **94**:723–729.
- Huang YT, Liu MY, Tsai CH, Yeh TH., (2010). Upregulation of interleukin-1 by Epstein- Barr virus latent membrane protein 1 and its possible role in nasopharyngeal carcinoma cell growth. *Head Neck* **32**:869–76.
- Huang YT, Sheen TS, Chen CL, Lu J, Chang Y, Chen JY, et al., (1999). Profile of cytokine expression in nasopharyngeal carcinomas: a distinct expression of interleukin 1 in tumor and CD4+ T cells. *Cancer Res* **59**:1599–605.
- Hui, AB, Or YY, Takano H, Tsang RK, To KF, Guan XY, et al., (2005). Array-based comparative genomic hybridization analysis identified cyclin D1 as a target oncogene at 11q13.3 in nasopharyngeal carcinoma. *Cancer Res* **65**: 8125-8133.
- Hu, L. F., J. Minarovits, S. L. Cao, B. Contreras- Salazar, L. Rymo, K. Falk, et al., (1991). Variable expression of latent membrane protein in nasopharyngeal carcinoma can be related to methylation status of the Epstein-Barr virus BNLF-1 5'-flanking region. *J Virol* **65**: 1558-1567.

- International Agency for Research on Cancer., (1997). Epstein-Barr virus and Kaposi's sarcoma herpesvirus 8. Monographs on evaluation of carcinogenic risks to humans, **70**, Lyon, France: WHO.
- Iwakiri, D., T. Minamitani and M. Samantha, (2013). Epstein-Barr virus latent membrane protein 2A contributes to anoikis resistance through ERK activation. *Journal of Virology* **87**(14): 8227-8234.
- Jay S. Desgrosellier and David A. Cheresh., (2010). Integrins in cancer : biological implications and therapeutic opportunities. Nature review. *Cancer* **10**: 9 - 20.
- Jones, P. A., S. B. Baylin., (2002). The fundamental role of epigenetic events in cancer. *Nat Rev Genet* **3**: 415-428.
- Jun Ma, Su Mei Cao., (2010). Epidemiology of Nasopharyngeal Carcinoma, Nasopharyngeal Cancer. *Medical Radiological* : 1-7.
- Kennedy, G., J. Komano & B. Sugden., (2003). Epstein-Barr virus provides a survival factor to Burkitt's lymphomas. *Proc Natl Acad Sci U S A* **100**: 14269-14274.
- Khanna, R., S. R. Burrows and D. J. Moss, (1995). Immune regulation in Epstein-Barr virus associated disease. *Microbiological Reviews* **59**(3): 387-405.
- Khanim F, Yao QY, Niedobitek G, et al., (1996). Analysis of Epstein-Barr virus gene polymorphisms in normal donors and in virus- associated tumors from different geographic locations. *Blood* **88**:3491-501.
- Kieff E., (1996). Epstein-Barr virus and its replication. In: Fields BN, Knipe DM, Howley P, et al, editors. *Fields virology*. Philadelphia: Lippincott-Raven 2343-96.
- Kim, T. H., H. I. Kim, Y. Soung, et al., (2009). Integrin (α6β4) signals through Src to increase expression of S100A4, a metastasis promoting factor: Implications for cancer cell invasion. *Molecular Cancer Research* **7**: 1605-1612.
- Kintner, C., B. Sugden., (1981). Conservation and progressive methylation of Epstein-Barr viral DNA sequences in transformed cells. *J Virol* **38**, 305-316.
- Kirschner AN, Omerovic J, Popov B, Longnecker R, Jardetzky TS., (2006). Soluble Epstein-Barr Virus Glycoprotein gH, gL, and gp42 Form a 1:1:1 Stable Complex That Acts Like Soluble gp42 in B-Cell Fusion but Not in Epithelial Cell Fusion. *Journal of Virology*: 9444-54.
- Kis LL., (2009). The role of the microenvironment on regulation of Epstein-Barr virus latent gene expression. Karoloska Institute, Stocholm.
- Klaus, P. and A.-P. Catherine, (2012). Detection methods of circulating tumor cells. *Journal of Thoracic Disease* **4**(5): 446-447.
- Klein G., (1979). The relationship of the virus to nasopharyngeal carcinoma. In:

- Epstein MA and Achong BG (eds), *The Epstein-Barr Virus*: 339-350. Springer-Verlag, Berlin.
- Kligys, K. R., Y. Wu, S. B. Hopkinsin, et al., (2012). $\alpha 6\beta 4$ integrin, a master regulator of expression of integrins in human keratinocytes. *Journal of Biological Chemistry* **287**(22): 17975-17984.
- Knox PG, Li Q-X, Rickinson AB et al., (1996). In vitro production of stable Epstein-Barr virus- positive epithelial cell clones which resemble the virus : cell interaction observed in nasopharyngeal carcinoma. *Virology* **215**: 40 – 50.
- Knutson J., (1990). The level of c-fgr RNA is increased by EBNA-2, an Epstein-Barr virus gene required for B-cell immortalization. *J Virol* **64**:2530 - 4.
- Kong QL, Hu LJ, Cao JY, Huang YJ, Xu LH, Liang Y, et al., (2010). Epstein-Barr Virus-Encoded LMP2A Induces an Epithelial– Mesenchymal Transition and Increases the Number of Side Population Stem-like Cancer Cells in Nasopharyngeal Carcinoma. *PLoS Pathogens* **6** (6).
- Korcum, A. F., E. Ozyar and A. Ayhan, (2006). Epstein-Barr virus genes and nasopharyngeal cancer. *Turkish Journal of Cancer* **36**(3): 97-107.
- Kuhn-Hallek I, Sage DR, Stein L, Groelle H, Fingerroth JD., (1995). Expression of recombination activating genes (RAG-1 and -2) in Epstein-Barr virus-bearing B-cells. *Blood* **85**:1289-99.
- Kusano, S., N. Raab-Traub., (2001). An Epstein-Barr virus protein interacts with Notch. *J Virol* **75**, 384-395.
- Laing KG, Matys V, Clemens MJ., (1995). Effects of expression of the Epstein-Barr virus small RNA EBER-1 in heterologous cells on protein synthesis and cell growth. *Biochem Soc Trans* **23**:311S.
- Lan YY, Hsiao JR, Chang KC, Chang JSM, Chen CW, Lai HC, et al., (2012). Epstein-Barr Virus Latent Membrane Protein 2A Promotes Invasion of Nasopharyngeal Carcinoma Cells through ERK/Fra-1-Mediated Induction of Matrix Metalloproteinase 9. *Journal of Virology* **86**(12): 6656–6667.
- Laux G, Perricaudet M, Farrell PJ., (1988). A spliced Epstein-Barr virus gene expressed in immortalized lymphocytes is created by circularization of the linear viral genome. *EMBO J* **7**:769 -74.
- Lawrence S. Young., (2014). Epstein-Barr virus at 50—future perspectives. *Chinese Journal* .
- Leight ER, Sugden B., (2000). EBNA1: a protein pivotal to latent infection by Epstein-Barr virus. *Rev Med Virol* **10**:83 -100.

- Li, H. P. and Y. S. Chang, (2003). Epstein-Barr virus latent membrane protein 1: structure and functions. *Journal of Biomedical Science* **10**(5): 490-504.
- Li, Z., Y. Lifang and S. L. Quan, (2012). Pathologic significance of EBV encoded RNA in NPC. InTech.
- Liao LJ and Lai MS., (2012). Epstein-Barr Virus Serology in the Detection and Screening of Nasopharyngeal Carcinoma in Carcinogenesis, Diagnosis, and Molecular Targeted Treatment for Nasopharyngeal Carcinoma, Dr. Shih-Shun Chen (Ed.). InTech.
- Liudmila, C. S. and M. H.-F. Lindsey, (2011). Fusion of Epstein-Barr Virus with Epithelial Cells Can Be Triggered by $\alpha v \beta 5$ in Addition to $\alpha v \beta 6$ and $\alpha v \beta 8$, and Integrin Binding Triggers a Conformational Change in Glycoproteins gHgL. *Journal of Virology* **85**(24): 13214-13223.
- Liudmila, C. S., S. L. Nishimura and M. H.-F. Lindsey, (2009). Fusion of epithelial cells by Epstein-Barr virus proteins is triggered by binding of viral glycoproteins gHgL to integrins $\alpha v \beta 6$ or $\alpha v \beta 8$. *Proceedings of the National Academy of Sciences of the United States of America* **106**(48): 20464-20469.
- Lo K, Huang D., (2002). Genetic and epigenetic changes in nasopharyngeal carcinoma. *Sem Cancer Biol* **12**:451–62.
- Lo KW, Huang DP, Lau KM., (1995). Gene alterations in nasopharyngeal carcinoma. *Cancer Res* **55**: 2039-2043.
- Lo, KW, To KF, Huang DP., (2004). Focus on nasopharyngeal carcinoma. *Cancer Cell* **5**: 423-428.
- Lo KW, Teo PM, Hui AB, To KF, Tsang YS, Chan SY, et al., (2000). High resolution allelotype of microdissected primary nasopharyngeal carcinoma. *Cancer Res* **60**: 3348-3353.
- Lo, Y. M., L. Chan, K. Lo, et al., (1999). Quantitative analysis of cell free Epstein-Barr virus DNA in plasma of patients with nasopharyngeal carcinoma. *Cancer Research* **59**: 1188-1191.
- Lo, Y. M., L. Y. Chan, A. T. Chan, S. F. Leung, K. W. Lo, J. Zhang, et al., (1999). Quantitative and temporal correlation between circulating cell-free Epstein-Barr virus DNA and tumor recurrence in nasopharyngeal carcinoma. *Cancer Res* **59**: 5452-5455.
- Lo, Y. M., A. T. Chan, L. Y. Chan, S. F. Leung, C. W. Lam, D. P. Huang & P. J. Johnson., (2000). Molecular prognostication of nasopharyngeal carcinoma by quantitative analysis of circulating Epstein-Barr virus DNA. *Cancer Res* **60**: 6878-6881.

- Longan, L. and R. Longnecker, (2000). Epstein-Barr virus latent membrane protein 2A has no growth-altering effects when expressed in differentiating epithelia. *The Journal of General Virology* **81**(9): 2245-2252.
- Longnecker, R., (2000). Epstein-Barr virus latency: LMP2, a regulator or means for Epstein-Barr virus persistence. *Adv Cancer Res* **79**: 175-200.
- Longnecker, R. and E. Kieff, (1990). "A second Epstein-Barr virus membrane protein (LMP2) is expressed in latent infection and colocalizes with LMP1." *Journal of Virology* **64**(5): 2319-2326.
- Lopez-Verrilli MA, Court FA, (2013). "Exosomes: mediators of communication in eukaryotes." *Biol Res* **46**: 5-11.
- Lung HL, Cheng Y, Kumaran MK, Liu ET, Murakami Y, Chan CY, et al., (2004). Fine mapping of the 11q22-23 tumor suppressive region and involvement of TSLC1 in nasopharyngeal carcinoma. *Int J Cancer* **112**: 628-635.
- Luo, B., Y. Wang, X. F. Wang, H. Liang, L. P. Yan, B. H. Huang & P. Zhao., (2005). Expression of Epstein- Barr virus genes in EBV-associated gastric carcinomas. *World J Gastroenterol* **11**: 629-633.
- Lynch, D. T., J. S. Zimmerman and D. T. Rowe, (2002). Epstein-Barr virus latent membrane protein 2B (LMP2B) co-localizes with LMP2A in perinuclear regions in transiently transfected cells. *The Journal of General Virology* **83**(5): 1025-1035.
- Makrilia N, Kollias A, Manolopoulos L, Syrigos K., (2009). Cell adhesion molecules: role and clinical significance in cancer. *Cancer Investigation* **27**(10): 1023–1037.
- Mariani Costantini RM, Falcioni R, Battista P, Kennel SJ, Colasante A, Venturo I, et al., (1990). "Integrin ($\alpha 6/\beta 4$) Expression in Human Lung Cancer as Monitored by Specific Monoclonal Antibodies." *American Association for Cancer Research*.
- Marzaini, D. S., D. L. Tobing, S. B. Kresno, et al., (2009). The accuracy of plasma EBV-DNA quantification using LMP2A as primer to detect distant metastasis after radiation of nasopharyngeal cancer in Dharmas National Cancer Center. *Indonesia Journal of Cancer* **3**(2): 47-51.
- Mazeron, M. C., (1996). Value of anti-Epstein-Barr antibody detection in the diagnosis and management of undifferentiated carcinoma of the nasopharynx. *Bull Cancer Radiother* **83**: 3-7.
- Mee JB, Johnson CM, Morar N, Burslem F, Groves RW., (2007). The psoriatic transcriptome closely resembles that induced by interleukin-1 in cultured

- keratinocytes: dominance of innate immune responses in psoriasis. *Am J Pathol* **171**:32–42.
- Mercurio AM, Rabinovitz I, Shaw LM., (2001). The $\alpha 6\beta 4$ integrin and epithelial cell migration. *Current Opinion in Cell Biology* **13** (5): 541–545.
- Michael D. Allen, Lawrence S. Young, and Christopher W. Dawson., (2005). The Epstein-Barr Virus-Encoded LMP2A and LMP2B Proteins Promote Epithelial Cell Spreading and Motility. *Journal of Virology* : 1789–1802.
- Middledrop, J., A. Brink, A. V. D. Brule, et al., (2003). Pathogenic roles for Epstein-Barr virus (EBV) gene products in EBV associated proliferative disorders. *Critical Reviews in Oncology/Hematology* **45**(1): 1-36.
- Miller, C. L., A. L. Burkhardt, J. H. Lee, et al., (1995). Integral membrane protein 2 of Epstein-Barr virus regulates reactivation from latency through dominant negative effects on protein-tyrosine kinases. *Immunity* **2**(2): 155-166.
- Miller WE, Earp HS, Raab-Traub N., (1995). The Epstein-Barr virus latent membrane protein 1 induces expression of the epidermal growth factor receptor. *J Virol* **69**:4390–8.
- Min Yu, Shannon Stott, Mehmet Toner, Shyamala Maheswaran, Daniel A. Haber. Circulating tumor cells: approaches to isolation and characterization. *J. Cell Biol* **192** (3): 373–382.
- Miyamoto, S., H. Teramoto, O. A. Coso, et al., (1995). Integrin function : Molecular hierarchies of cytoskeletal and signaling molecules. *Journal of Cell Biology* **131**.
- Mizejewski GJ., (1999). Role of Integrins in Cancer: Survey of Expression Patterns. Issue . Proceedings of the Society for Experimental Biology and Medicine Proceedings of the Society for Experimental Biology and Medicine **222**(2): 124–138.
- Moody CA, Scott RS, Su T, Sixbey JW., (2003). Length of Epstein-Barr virus termini as a determinant of epithelial cell clonal emergence. *J Virol* **77**:8555–61.
- Morris MA, Dawson CW, Young LS., (2009). Role of the Epstein-Barr virus-encoded latent membrane protein 1, LMP1, in the pathogenesis of nasopharyngeal carcinoma. *Future Oncol* **5**:811–25.
- Morrison, J. A., A. J. Klingehutz and N. Raab-Traub, (2003). Epstein-Barr virus latent membrane protein 2A activates β -catenin signaling in epithelial cells. *Journal of Virology* **77**(22): 12276-12284.
- Morrison, J. A. and N. Raab-Traub, (2005). "Roles of the ITAM and PY motifs of Epstein-Barr virus latent membrane protein 2A in the inhibition of epithelial

- cell differentiation and activation of {beta}-catenin signaling." *Journal of Virology* **79**(4): 2375-2382.
- Murakami, M., K. Lan, C. Subramanian & E. S. Robertson., (2005). Epstein-Barr virus nuclear antigen 1 interacts with Nm23-H1 in lymphoblastoid cell lines and inhibits its ability to suppress cell migration. *J Virol* **79**: 1559-1568.
- Murray, P. G., C. M. Constandinou, J. Crocker, L. S. Young & R. F. Ambinder., (1998). Analysis of major histocompatibility complex class I, TAP expression, and LMP2 epitope sequence in Epstein-Barr virus-positive Hodgkin's disease. *Blood* **92**: 2477-2483.
- Murray, P. G. and L. S. Young, (2001). Epstein-Barr virus infection : basis of malignancy and potential for therapy. *Cambridge Journals Online* **3**: 1-20.
- Mutirangura A., Pornthanakasem W, Theamboonlers A, Sriuranpong V, Lertsanguansinchi P, Yenrudi S, et al., (1998). Epstein-Barr viral DNA in serum of patients with nasopharyngeal carcinoma. *Clin. Cancer Res* **4**:665–669.
- Nicholls, J. M., A. Agathangelou, K. Fung, X. Zeng & G. Niedobitek., (1997). The association of squamous cell carcinomas of the nasopharynx with Epstein-Barr virus shows geographical variation reminiscent of Burkitt's lymphoma. *J Pathol* **183**, 164-168.
- Niedobitek, G., (2000). Epstein-Barr virus infection in the pathogenesis of nasopharyngeal carcinoma. *Journal Clinical Pathology* **53**: 248-254.
- Niedobitek, G., M. I. Hansmann, H. Herbst, I. S. Young, D. Dienemann, C. A. Hartmann, et al., (1991). Epstein-Barr virus and carcinomas: Undifferentiated carcinomas but not squamous cell carcinomas of the nasopharynx are regularly associated with the virus. *J Pathol* **165**: 17-24.
- Oh,S.T.,J.S.Seo,U.Y.Moon,K.H.Kang,D.J.,Shin,S.K.Yoon,et al., (2004). A naturally derived gastric cancer cell line shows latency I Epstein-Barr virus infection closely resembling EBV- associated gastric cancer. *Virology* **320**: 330-336.
- Okano, M., D. W. Bell, D. A. Haber & E. Li., (1999). DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development. *Cell* **99**: 247-257.
- Ordonez, B. P., (2007). An update on Epstein-Barr virus and nasopharyngeal carcinogenesis. *Head and Neck Pathology* **1**: 141-145.
- Pang, M. F., K. W. Lin and S. C. Peh, (2009). The signaling pathways of Epstein-Barr virus-encoded latent membrane protein 2A (LMP2A) in latency and cancer. *Cellular and Molecular Biology Letters* **14**(2): 222-247.
- Pathmanathan, R., U. Prasad, R. Sadler, K. Flynn & N. Raab-Traub., (1995). Clonal proliferations of cells infected with Epstein-Barr virus in preinvasive lesions

- related to nasopharyngeal carcinoma. *N Engl J Med* **333**: 693-698.
- Pathmanathan, R., U. Prasad, G. Chandrika, R. Sadler, K. Flynn & N. Raab-Traub., (1995). Undifferentiated, nonkeratinizing, and squamous cell carcinoma of the nasopharynx. Variants of Epstein-Barr virus-infected neoplasia. *Am J Pathol* **146**: 1355- 1367.
- Pegtel, D. M., A. Subramanian, T. S. Sheen, et al., (2005). Epstein-Barr virus encoded LMP2A induces primary epithelial cell migration and invasion : Possible role in nasopharyngeal carcinoma metastasis. *Journal of Virology* **79**(24): 15430-15442.
- Portis, T., L. Cooper, P. Dennis, et al., (2002). The LMP2A signalosome--a therapeutic target for Epstein-Barr virus latency and associated disease. *Frontiers in Bioscience : A Journal and Virtual Library* **1**(7): 414-426.
- Portis, T. and R. Longnecker, (2004). Epstein-Barr virus (EBV) LMP2A mediates B-lymphocyte survival through constitutive activation of the Ras/PI3K/Akt pathway. *Oncogene* **23**: 8619-8628.
- Pratesi C, Maria, Bortolin MT, D'Andrea M, Vaccher E, Barzan L, Bidoli E, et al., (2003). "Quantitative plasma/serum EBV DNA load by LMP2A determination in an Italian cohort of NPC patients." *Journal of Clinical Virology* **28**:155-164.
- Pricipe S, Hui AB-Y, Bruce J, Sinha A, Liu F-F, Klisinger T, (2013). "Tumor-derived exosomes and microvesicles in head and neck cancer : Implications for tumor biology and biomarker discovery." *Proteomic* **13**: 1608-1623.
- Raab-Traub, N., (2002). Epstein-Barr virus in the pathogenesis of NPC. *Seminar in Cancer Biology* **12**(6): 431-441.
- Raab-Traub, N., K. Flynn., (1986). The structure of the termini of the Epstein-Barr virus as a marker of clonal cellular proliferation. *Cell* **47**: 883-889.
- Raab-Traub, N., K. Flynn, G. Pearson, A. Huang, P. Levine, A. Lanier & J. Pagano., (1987). The differentiated form of nasopharyngeal carcinoma contains Epstein-Barr virus DNA. *Int J Cancer* **39**: 25-29.
- Rabinovitz, I. and A. Mercurio, (1997). The integrin alpha6beta4 functions in carcinoma cell migration on laminin-1 by mediating the formation and stabilization of actin containing motility structures. *The Journal Cell of Biology* **139**(7): 1873-1874.
- Renato, F., C. Monica, Z. M. Federica, et al., (2012). Circulating tumor cells as emerging tumor biomarkers in lung cancer. *Journal of Thoracic Disease* **4**(5): 438-439.
- Richardson K., (2015). Enhanced Detection of Low-Level DNA Mutations Using

Multiplexed ICE COLD PCR Coupled to NGS or ddPCR. 22nd International Molecular Medicine Tri-Conference.

- Robertson, K. D., A. P. Wolffe., (2000). DNA methylation in health and disease. *Nat Rev Genet* **1**: 11-19.
- Rovedo, M. and R. Longnecker, (2007). Epstein-Barr virus latent membrane protein 2B (LMP2B) modulates LMP2A activity. *Journal of Virology* **81**: 89-94.
- Rovedo, M. A.,(2008). Analysis of latent membrane protein (LMP) 2A signaling through Src Family Kinases and the role of LMP2B in modulating LMP2A signaling. Evanston, Illinois.Northwestern University
- Sall, A., S. Caserta, P. Jolicoeur, L. Franqueville, M. de Turenne-Tessier & T. Ooka., (2004). Mitogenic activity of Epstein-Barr virus-encoded BARP1 protein. *Oncogene* **23**: 4938-4944.
- Sample J, Young LS, Martin B, et al., (1990). Epstein-Barr virus types 1 and 2 differ in their EBNA3A, EBNA3B and EBNA3C genes. *J Virol* **64**:4084-92.
- Scholle, F., R. Longnecker and N. Raab-Traub, (1999). Epithelial cell adhesion to extracellular matrix proteins induces tyrosine phosphorylation of the Epstein-Barr virus latent membrane protein 2: a role for C-terminal Src kinase. *Journal of Virology* **73**(6): 4767-4775.
- Scholle, F., K. M. Bendt & N. Raab-Traub., (2000). Epstein- Barr virus LMP2A transforms epithelial cells, inhibits cell differentiation, and activates Akt. *J Virol* **74**: 10681-10689.
- Schorey JF, Cheng Y, Singh PP, Smith VL, (2015). "Exosomes and other extracellular vesicles in host-pathogen interactions." *EMBO Report* **16**: 24-43.
- Sejeong Shin, Laura Wolgamott, and Sang-Oh Yoon., (2012). Integrin Trafficking and Tumor Progression. Hindawi Publishing Corporation. *International Journal of Cell Biology* (7).
- Seto, E., L. Yang, J. Middeldorp, T. S. Sheen, J. Y. Chen, M. Fukayama., (2005). Epstein-Barr virus (EBV)-encoded BARP1 gene is expressed in nasopharyngeal carcinoma and EBV- associated gastric carcinoma tissues in the absence of lytic gene expression. *J Med Virol* **76**, 82-88.
- Shair, K. H., C. I. Schnegg and N. Raab-Traub, (2008). EBV latent membrane protein 1 effects on plakoglobin, cell growth, and migration. *Cancer Research* **68**(17): 6997-7005.
- Shao, J. Y., I. Ernberg, P. Biberfeld, T. Heiden, Y. X. Zeng & L. F. Hu., (2004). Epstein-Barr virus LMP1 status in relation to apoptosis, p53 expression and leucocyte infiltration in nasopharyngeal carcinoma. *Anticancer Res* **24**: 2309-2318.

- Shaw, L. M., I. Rabinovitz, H. H. Wang, et al., (1997). Activation of Phosphoinositide 3-OH kinase by the $\alpha 6 \beta 4$ integrin promotes carcinoma invasion. *Journal of Cell Biology* **91**: 949-960.
- Sheen TS, Huang YT, Chang YL, Ko JY, Wu CS, Yu YC, et al., (1999). Epstein-Barr virus-encoded latent membrane protein 1 co-expresses with epidermal growth factor receptor in nasopharyngeal carcinoma. *Jpn J Cancer Res* **90**:1285–92.
- Sheng, W., G. Decaussin, A. Ligout, K. Takada & T. Ooka., (2003). Malignant transformation of Epstein-Barr virus- negative Akata cells by introduction of the BART1 gene carried by Epstein-Barr virus. *J Virol* **77**: 3859-3865.
- Shin S, Wolgamott L, Yoon S., (2012). Integrin Trafficking and Tumor Progression. *International Journal of Cell Biology*.
- Sinclair AJ, Palmero I, Peters G, Farrell PJ., (1994). EBNA-2 and EBNA-LP cooperate to cause G0 to G1 transition during immortalization of resting human B lymphocytes by Epstein- Barr virus. *EMBO J* **13**:3321 - /8.
- Smith PR, Gao Y, Karran L, Jones MD, Snudden D, Griffin BE., (1993). Complex nature of the major viral polyadenylated transcripts in Epstein/Barr virus-associated tumors. *J Virol* **67**:3217 -25.
- Smith, P. R., J. O. de, D. Turner, M. Hollyoake, C. E. Karstegl, B. E. Griffin., (2000). Structure and coding content of CST (BART) family RNAs of Epstein-Barr virus. *J Virol* **74**: 3082-3092.
- Srinivas S, Sixbey JW., (1995). Epstein-Barr virus induction of recombinase-activating genes RAG1 and 2. *J Virol* **69**:8155-/8.
- Stevens, S. J., S. A. Verkuijlen, B. Hariwiyanto, et al., (2005). Diagnostic value of measuring Epstein-Barr virus (EBV) DNA load and carcinoma specific viral mRNA in relation to anti-EBV immunoglobulin A (IgA) and IgG antibody levels in blood of nasopharyngeal carcinoma patients from Indonesia. *Clinical Microbiology* **43**(7): 3066-3073.
- Stewart, S., C. W. Dawson, K. Takada, et al., (2004). Epstein-Barr virus encoded LMP2A regulates viral and cellular gene expression by modulation of the MF-KB transcription factor pathway. *PNAS* **101**(44): 15730-15735.
- Sugiura, M., S. Imai, M. Tokunaga, S. Koizumi, M. Uchizawa, K. Okamoto & T. Osato., (1996). Transcriptional analysis of Epstein-Barr virus gene expression in EBV- positive gastric carcinoma: unique viral latency in the tumour cells. *Br J Cancer* **74**: 625-631.
- Su-Mei Cao, Malcolm J Simons, Chao-Nan Qian., (2011). The preventive and prevention of nasopharyngeal carcinoma in China. *Clin J Cancer* **30**(2).
- Tabuchi K., (2011). Early Detection of Nasopharyngeal Carcinoma. Hindawi

Publishing Corporation. *International Journal of Otolaryngology*.

- Takada K, Nanbo A., (2001). The role of EBERs in oncogenesis. *Semin Cancer Biol* **11**:461-7.
- Tannock, I., R. Hill, R. Bristow, et al., (2005). Viruses and cancer in the basic of oncology. I. Tannock, R. Hill, R. Bristow and L. Harrington. Boston, Mc Graw Hill Co, Inc.
- Tao Jin, Hao Peng, Hongyan Wu., (2013). Clinical value of circulating liver cancer cells for the diagnosis of hepatocellular carcinoma: A meta-analysis. *Biomedical Report* **1**: 731-736.
- Tao Q, Chan AT., (2007). Nasopharyngeal carcinoma: molecular pathogenesis and therapeutic developments. *Expert Rev Mol Med* **9**:1-24.
- Tao, Q., F. C. Ho, S. L. Loke & G. Srivastava., (1995). Epstein- Barr virus is localized in the tumour cells of nasal lymphomas of NK, T or B cell type. *Int J Cancer* **60**: 315- 320.
- Tao, Q., K. D. Robertson., (2003). Stealth technology: how Epstein-Barr virus utilizes DNA methylation to cloak itself from immune detection. *Clin Immunol* **109**: 53-63.
- Thorley-Lawson DA., (2001). Epstein-Barr virus: exploiting the immune system. *Nat Rev Immunol* **1**:75/82.
- Thorley-Lawson DA, Miyashita EM., (1996). Epstein-Barr virus and the B-cell: that's all it takes. *Trends Microbiol* **4**:204-8.
- Thornburg, N. J., S. Kusano & N. Raab-Traub., (2004). Identification of Epstein-Barr virus RK-BARF0- interacting proteins and characterization of expression pattern. *J Virol* **78**: 12848-12856.
- Tierney, R. J., H. E. Kirby, J. K. Nagra, J. Desmond, A. I. Bell & A. B. Rickinson., (2000). Methylation of transcription factor binding sites in the Epstein-Barr virus latent cycle promoter Wp coincides with promoter down-regulation during virus-induced B-cell transformation. *J Virol* **74**: 10468-10479.
- TierneyR,StevenN,YoungL,RickinsonA., (1994). Epstein-Barrvirus latency in blood mononuclear cells: analysis of viral gene transcription during primary infection and in the carrier state. *J Virol* **68**(11):7374-85.
- Toczyski DP, Steitz JA., (1991). EAP, a highly conserved cellular protein associated with Epstein-Barr virus small RNAs (EBERs). *EMBO J* **10**:459-66.
- Tsang, C. M., W. Deng, Y. L. Yip, et al., (2014). Epstein-Barr virus infection and persistence in nasopharyngeal epithelial cells. *Chinese Journal of Cancer* **33**(11): 549-555.

- Tsao, S. W., G. Tramoutanis, C. W. Dawson, et al., (2002). The significance of LMP1 expression in nasopharyngeal carcinoma. *Seminar in Cancer Biology* **12**(6): 473-487.
- Tsao, S. W., C. M. Tsang, P. S. Pang, et al., (2012). The biology of EBV infection in human epithelial cells. *Seminar in Cancer Biology* **22**(2): 137-143.
- Wang F, Tsang SF, Kurilla MG, Cohen JI, Kieff E., (1991). Epstein-Barr virus nuclear antigen 2 transactivates latent membrane protein LMP1. *J Virol* **64**:3407 -16.
- Wang, Q., S. W. Tsao, T. Ooka, J. M. Nicholls, H. W. Cheung, S. Fu., (2005). Anti-apoptotic role of BARP1 in gastric cancer cells. *Cancer Lett* Jul 26, Epub ahead of print.
- Wasil LR, Tomaszewski MJ, Hoji A, Rowe DT., (2013). The Effect of Epstein-Barr Virus Latent Membrane Protein 2 Expression on the Kinetics of Early B Cell Infection. *Plos One* **8**(1).
- Weiss, L. M., L. A. Movahed, A. E. Butler, S. A. Swanson, H. F. Frierson, P. H. Cooper, et al., (1989). Analysis of lymphoepithelioma and lymphoepithelioma-like carcinomas for Epstein-Barr viral genomes by in situ hybridization. *Am J Surg Pathol* **13**, 625-631.
- Wendel, M., L. Bazhenova, R. Boshuzien, et al., (2012). Fluid biopsy for circulating tumor cell identification in patients with early-and late-stage non-small cell lung cancer: a glimpse into lung cancer biology. *Physical Biology* **9**(1): 016005.
- Wu H, Kapoor P, Frappier L., (2002). Separation of the DNA replication, segregation, and transcriptional activation functions of Epstein-Barr nuclear antigen 1. *J Virol* **76**:2480-90.
- Xiong, W., Z. Y. Zeng, J. H. Xia, K. Xia, S. R. Shen, X. L. Li, D. et al., (2004). Susceptibility locus at chromosome 3p21 linked to familial nasopharyngeal carcinoma. *Cancer Res* **64**, 1972-1974.
- Xuan SH, Zhou YG, Pan JQ, Zhu W, Xu P., (2013). Overexpression of integrin α v in the human nasopharyngeal carcinoma associated with metastasis and progression. *Cancer Biomark* **13**(5):323-8.
- Xue, S. A., L. G. Labrecque, Q. L. Lu, S. K. Ong, I. A. Lampert, P. Kazembe, et al., (2002). Promiscuous expression of Epstein-Barr virus genes in Burkitt's lymphoma from the central African country Malawi. *Int J Cancer* **99**: 635-643.
- Yang Chenjie, Robbins PD, (2011). "The Roles of Tumor-Derived Exosomes in Cancer Pathogenesis." Hindawi Publishing Corporation. Clinical and Developmental Immunology.

- Yang, L., K. Aoz, K. Oshimi, et al., (2004). Epstein-Barr virus encoded RNA promote growth of EBV infected T cells through interleukin-9 induction. *Cancer Research* **64**(1): 5332-5337.
- Yamamoto N, Takizawa T, Iwanaga Y, Shimizu N, Yamamoto N., (2000). Malignant transformation of B lymphoma cell line BJAB by Epstein-Barr virus-encoded RNAs. *FEBS Lett* **484**:153-8.
- Ye S-B, Li Z-L, Luo D-H, Huang Bi-Jun, Chen YU-Suan, Zhang X-S,, Cui J, Zeng Y-I, et al., (2014). "Tumor-derived exosome promote tumor progression and T-cell dysfunction through the regulation of enriched exosomal microRNAs in human nasopharyngeal carcinoma." *Oncotarget* **5**(14): 5439-5452.
- Ying, J., Srivastava G, Hsieh WS, Gao Z, Murray P, Liao SK, et al., (2005). The stress-responsive gene GADD45G is a functional tumor suppressor, with its response to environmental stresses frequently disrupted epigenetically in multiple tumors. *Clin Cancer Res* **11**: 6442-6449.
- Yong-Lin, C., J. Li, A. Y. Lu, et al., (2014). Diagnostic Significance of combined detection of Epstein-Barr virus antibodies, VCA/IgA, Rta/IgG and EBNA1/IgA for nasopharyngeal carcinoma. *Asian Pac J Cancer Prev* **15**(5): 2001-2006.
- Young, L. S., A. B. Rickinson., (2004). Epstein-Barr virus: 40 years on. *Nat Rev Cancer* **4**: 757-768.
- Young LS, Murray PG., (2003). Epstein-Barr virus and oncogenesis: from latent genes to tumours. *Oncogene* **22**: 5108-5121.
- Yu, M. C., J. M. Yuan., (2002). Epidemiology of nasopharyngeal carcinoma. *Semin Cancer Biol* **12**: 421-429.
- Yu, M., S. Stott, M. Toner, et al., (2011). Circulating tumor cells: approaches to isolation and characterization. *Journal of Cell Biology* **192**(3): 372-382.
- Zheng, H., L. L. Li, D. S. Hu, et al., (2007). Role of Epstein-Barr virus encoded latent membrane protein 1 in the carcinogenesis of nasopharyngeal carcinoma. *Cellular and Molecular Immunology* **4**(3): 185-196.
- Zheng H, Li LL, Hu DS, Deng XY, Cao Y., (2007). Role of Epstein-Barr virus encoded latent membrane protein 1 in the carcinogenesis of nasopharyngeal carcinoma. *Cell Mol Immunol* **4**:185-96.
- Zhou, L., W. Jiang, C. Ren, Z. Yin, X. Feng, W. Liu, et al., (2005). Frequent Hypermethylation of RASSF1A and TSLC1, and High Viral Load of Epstein-Barr Virus DNA in Nasopharyngeal Carcinoma and Matched Tumor-Adjacent Tissues. *Neoplasia* **7**: 809-815.
- Zimber-Strobl U, Strobl LJ., (2001). EBNA2 and Notch signalling in Epstein -Barr virus mediated immortalization of B lymphocytes. *Semin Cancer Biol* **11**:423 -

34.

Zur Hausen, A., A. A. Brink, M. E. Craanen, J. M. Middeldorp, C. J. Meijer & A. J. van den Brule., (2000). Unique transcription pattern of Epstein-Barr virus (EBV) in EBV-carrying gastric adenocarcinomas: expression of the transforming BARF1 gene. *Cancer Res* **60**: 2745- 2748.

Zur Hausen, H., H. Schulte-Holthausen, G. Klein, W. Henle, G. Henle, P. Clifford & L. Santesson., (1970). EBV DNA in biopsies of Burkitt tumors and anaplastic carcinomas of the nasopharynx. *Nature* **228**: 1056-1058.