



DAFTAR PUSTAKA

Agnello D, Bigini P, Villa P, Mennini T, Cerami A, Brines ML, Ghezzi P (2002). Erythropoietin exerts an anti-inflammatory effect on the CNS in a model of experimental autoimmune encephalomyelitis. *Brain Res*;952:128-34.

Baciu I, Oprisiu C, Derevenco P, Vasile V, Muresan A, Hriscu M, Chis I (2000). The brain and other sites of erythropoietin production. *Rom J Physiol*;37: 3-14.

Bolanos JP, Almada A (1999). Roles of nitric oxide in brain hypoxia ischemia. *Biochim Biophys Acta*;1411: 415-36.

Brines ML, Ghezzi P, Keenan S, Agnello D, de Lanerolle NC, Cerami C, Cerami A (2000). Erythropoietin crosses the blood-brain barrier to protect against experimental brain injury. *Proc Natl Acad Sci USA*; 97:10526-31.

Bruick RK, McKnight SL (2001). A conserved family of prolyl-4-hydroxylases that modify HIF. *Science*;294:1337- 40.

Buemi M, Allegra A, Corica F, Floccari F, D'Avella D, Aloisi C, Calapai G, Iacopino G, Frisina N (2000). Intravenous recombinant erythropoietin does not lead to an increase in cerebrospinal fluid erythropoietin concentration. *Nephrol Dial Transplant*;15:422-3.

Buemi M, Cavallaro E, Floccari F, Sturiale A, Trimarchi M, Corica F, Frisina N (2003). The pleiotropic effects of erythropoietin in the central nervous system. *J Neuropathol Exp Neurol*;62:228-36.

Byts N, Siren AL (2009). Erythropoietin: a multimodal neuroprotective agent. *Exp Transl Stroke Med* ;1:4.

Dirnagl U, Iadecola C, Moskowitz MA (1999). Pathobiology of ischemic stroke: an integrated view. *Trends in Neuroscience* ; 22 (9): 391-7.

Elmahdy H, El Mashad AR, El Bahrawi H, ElGohary T, El Barbary A, Aly H (2010). Human Recombinant Erythropoietin in Asphyxia Neonatorum: Pilot Trial. *Pediatrics*:125; 1135-42.

Epstein AC, Gleadle JM, McNeill LA, Hewitson KS, O'Rourke J, Mole DR, Mukherji M, Wilson MI, Dhanda A (2001). C. elegans EGL-9 and mammalian homologs define a family of dioxygenases that regulate HIF by prolyl hydroxylation. *Cell*;107:43-54.

Ehrenreich H, Hasselbatt M, Dembowski C (2002). Erythropoietin therapy for acute stroke is both safe and beneficial. *Mol Med*; 8: 495-505.



Fanaroff AA, Martin RJ, Walsh MC (2006). Response to asphyxia. Dalam: Neonatal perinatal medicine, disease of the fetus and infant. Ed 8. Fanaroff AA. Mosby Elsevier-Philadelphia:483-90.

Ferriero DM (2004). Neonatal brain injury. *N Engl J Med* ;351: 1985-95

Fisher JW (2003). Erythropoietin: physiology and pharmacology update. *Exp Biol Med*; 228: 1-14.

Forstermann U, Schmidt HH, Pollock JS, Sheng H, Mitchell JA, Warner TD, Nakane M, Murad F (1991). Isoforms of nitric oxide synthase. Characterization and purification from different cell types. *Biochem Pharmacol* ;42:1849-57.

Frankenburg WK, Dodds J, Acher P, Shapiro H, Bresnick B (1992). The Denver II: A Major Revision and Restandardization of Denver Developmental Screening Test. *Pediatrics*;89: 91-7.

Gomella TL, Cunningham MD, Eyal FG (2004). Perinatal asphyxia. Dalam Neonatology. Ed 5. Gomella TL. Lange Medical book-Newyork:480-9.

Guix FX, Uribealago I, Coma M, Munoz FJ (2005). The physiology and pathophysiology of nitric oxide in the brain. *Prog Neurobiol*;76:126-52.

Gunn AJ (2000). Cerebral hypothermia for prevention of brain injury following perinatal asphyxia. *Curr Opin Pediatr*;12(2):111-15.

Hewitson KS, McNeill LA, Riordan MV, Tian YM, Bullock A, Welford RW, Elkins JM, Gleadle JM (2002). Hypoxia-inducible factor (HIF) asparagine hydroxylase is identical to factor inhibiting HIF (FIH) and is related to the cupin structural family. *J Biol Chem*;277:26351-5.

Ivan M, Kondo K, Yang H F, Kim W, Valiando J, Ohh M, Salic A, Lane WS, Kaelin WG (2001). HIF α targeted for VHL-mediated destruction by proline hydroxylation: Implications for O₂ sensing. *Science* ;292:464-8.

Jaakkola P, Mole DR, Tian YM, Wilson MI, Gielbert J, Gaskell SJ, Von Kriegsheim A, Mukherji M, Schofield CJ (2001). Targeting of HIF α to the von Hippel-Lindau ubiquitylation complex by O₂-regulated prolyl hydroxylation. *Science* ; 292:468-72.

Johnston MV, Nakajima W, Hagberg H (2002). Mechanism of Hypoxic neurodegeneration in developing brain. *Neuroscientist*;8(3):212-20.

Jiang BH, Semenza GL, Bauer C, Marti HH (1996). Hypoxia-inducible factor 1 levels vary exponentially over a physiologically relevant range of O₂ tension. *Am J Physiol*; 271:C1172 -80.



Juul SE, Anderson DK, Li Y, Christensen RD (1998). Erythropoietin and erythropoietin receptor in the developing human central nervous system. *Pediatr Res*;43:40-9.

Juul SE (2002). Erythropoietin in the central nervous system, and its use to prevent hypoxic-ischemic brain damage. *Acta Paediatr*;438:36-42.

Juul SE, Mc Pherson RJ, Farrel FX, Jolliffe L, Ness DJ, Gleason CA (2004). Erythropoietin concentrations in cerebrospinal fluid of non human primate and fetal sheep following high dose recombinant erythropoietin. *Biol Neonate*; 85: 138-49.

Juul SE (2010). Erythropoietin as neonatal neuroprotective agent. *Neo Reviews*;11(2):e78.

Kaufman JS, Reda DJ, Fye CL, Henderson WG, Kleinman JG, Vaamonde CA (1998). Subcutaneous versus intravenous administration of erythropoietin in hemodialysis patients. *N Engl J Med*; 339:578-83.

Kattwinkel J (2013). Dalam Textbook of Neonatal resuscitation. Ed 5. Kattwinkel J. American Academy of Pediatrics and American Heart Association-USA: 8-10.

Lando D, Gorman JJ, Whitelaw ML, Peet DJ (2003). Oxygen-dependent regulation of hypoxia-inducible factors by prolyl and asparaginyl hydroxylation. *Eur J Biochem*; 270:781-90.

Leviton A, Nelson KB (1992). Problems with definitions and classifications of newborn encephalopathy. *Pediatr Neurol*; 8(2): 85-90.

Levene MI (1995). The asphyxiated newborn infant. Dalam Fetal and neonatal neurology and neuro-surgery. Ed. Levene MI, Lilford RJ. Churchill Livingstone-Edinburg:405-26.

Liu C, Shen K, Liu Z, Noguchi CT (1997). Regulated human erythropoietin receptor expression in mouse brain. *J Biol Chem*;272:32395-400.

Low JA (2004). Determining the contribution of asphyxia to brain damage in the neonate. *J Obstet Gynaecol Res*; 30(4): 276-86.

Maiese K, Li F, Chong ZZ (2005). New Avenues of Exploration for Erythropoietin. *JAMA* ;293(1):90-5.

Marlow N, Rose AS, Rands CE, Draper ES (2005). Neuropsychological and educational problems at school age associated with neonatal encephalopathy. *Arch Dis Child Fetal Neonatal*;90:380-7.



Marti HH, Gassmann M, Wenger RH, Kuitikova F, Morganti-Kossmann MC, Kosmann T, Trentz O, Bauer C (1997). Detection of erythropoietin in human liquor, intrinsic erythropoietin production in the brain. *Kidney Int*; 51:416-8.

Marti HH, Bernaudin M, Petit E, Bauer C (2000). Neuroprotection and angiogenesis: a dual role of erythropoietin in brain ischemia. *New Physiol Sci*; 15:225-9

Marti HH (2004). Erythropoietin and the hypoxic brain. *J Exp Biol*; 207(18):3233-42.

Matsushita H, Johnston MV, Lange MS, Wilson MA (2003). Protective effect of erythropoietin in neonatal hypoxic ischemia in mice. *Dev Neuro Sci*; 14:1757-61.

Masson N, Ratcliffe PJ (2003). HIF prolyl and asparaginyl hydroxylases in the biological response to intracellular O₂ levels. *J Cell Sci*; 116:3041-9.

Miller SP, Ramaswamy V, Michelson D, Barkovich AJ (2005). Patterns of brain injury in term neonatal encephalopathy. *J Pediatr*; 146:453-60.

Miranda KM, Espey MG, Wink DA (2001). A rapid simple spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide*; 5(1):62-71.

Nunez JL, McCarthy MM (2008). Androgen predispose males to GABA-mediated excitotoxicity in the developing hippocampus. *Exp Neurol*; 210(2): 699-708.

Perlman JM, Risser R (1996). Can asphyxiated infants at risk for neonatal seizures be rapidly identified by current high-risk markers. *Pediatrics* ; 97: 456-62.

Pugh CW, Ratcliffe PJ (2003). Regulation of angiogenesis by hypoxia: role of the HIF system. *Nat Med*; 9:677-84.

Risau W (1997). Mechanism of angiogenesis. *Nature* ;386:671-4.

Robert CV, Jeffrey MP (1997). Interventions for Perinatal Hypoxic-Ischemic Encephalopathy. *Pediatrics*; 100(6):1004-114.

Ronen GM, Buckley D, Penney S, David RN (2007). Longterm prognosis in children with neonatal seizures: a population-based study. *Neurology*; 69(19):1816-22.

Sastroasamoro S, Madiyono B, Moeslichan S, Budiman I, Purwanto H (2008). Perkiraan besar sampel. In: Sastroasamoro S, Ismael S. Dasar-dasar metodologi penelitian klinis. 3th ed. Jakarta : CV Sagung Seto;302-29.



Savman K (2008). The brain and resuscitation. *Neo Reviews*;9:e513-9.

Semenza GL, Wang GL (1992). A nuclear factor induced by hypoxia via de novo protein synthesis binds to the human erythropoietin gene enhancer at a site required for transcriptional activation. *Mol Cell Biol*;12:5447-54.

Shahshahani S, Vameghi R, Azari N, Sajedi F, Kazemnejad A (2010). Validity and reliability determination of Denver Developmental Screening Test-II in 0-6 years old in Tehran. *Iran J Pediatr*;20: 313-22.

Sie LT, Van der Knaap MS, Oosting J, De Vries LS, Lafeber HN, Valk J (2000). MR patterns of hypoxic-ischemic brain damage after prenatal, perinatal or postnatal asphyxia. *Neuropediatrics*;31: 128-36.

Spandou E, Soubasi V, Papoutsopoulou S, Simeonidou C, Kaiki AA, Guiba TO (2005). Erythropoietin prevents long-term sensorimotor deficits and brain injury following neonatal hypoxia-ischemia in rats. *Brain Res* ;1045:22-30.

Sun Y, Zhou C, Polk P, Nanda A, Zhang JH (2004). Mechanisms of erythropoietin-induced brain protection in neonatal hypoxia-ischemia rat model. *J Cereb Blood Flow Metab*;24:259-70.

Tan CC, Eckardt KU, Firth JD, Ratcliffe P J (1992). Feedback modulation of renal and hepatic erythropoietin mRNA in response to graded anemia and hypoxia. *Am J Physiol*;263:F474-81.

Van de Berg, WD (2002). Perinatal asphyxia induced neuronal loss by apoptosis in the neonatal rat striatum: a combined Tunel and stereological study. *Exp Neurol*; 174(1): 29-36.

Volpe JJ (2008). Hypoxic Ischemic Encephalopathy: clinical aspect. Dalam *Neurology of the newborn*. Ed 5. Volpe JJ. Saunders Elsevier-Philadelphia:400-80.

Wagner CL, Eicher DJ, Barbosa E, Holden KR (1999). The use of hypothermia: a role in the treatment of neonatal asphyxia. *Pediatr Neurol*;21:429-43.

Wanner RM, Spielman P, Stroka DM, Camenisch G, Camenisch I, Scheid E, Bauer C, Gassmann M, Wenger RH (2000). EPOlones induces erythropoietin expression via hypoxia-inducible factor-1 α activation. *Blood*; 96:1558-65.

Wenger RH (2002). Cellular adaptation to hypoxia: O₂-sensing protein hydroxylases, hypoxia-inducible transcription factors, and O₂-regulated gene expression. *FASEB J*;16:1151-62.



World Health Organization (1999). Basic Newborn Resuscitation. A Practical Guide-Revision. Geneva:WHO. Diunduh dari: www.who.int/reproductive-health/publication/newborn_resuscitation/index.html.

World Health Organization (2007). Birth Asphyxia-Summary of the previous meeting and protocol overview. Milano 11 Juni 2007. Geneva:WHO.

World Health Organization (2014). Situation of newborn and child health in South-East Asia. Progress towards MDG 4. WHO-India:105-25.

Wu H, Lee SH, Gao J, Liu X, Iruela-Arispe ML (1999). Inactivation of erythropoietin leads to defects in cardiac morphogenesis. *Development*;126:3597-605.

Zhu C, Kang W, Xu F, Cheng X, Zhang Z, Jia L, Guo X, Xiong H, Simbruner G, Wang X (2009). Erythropoietin Improved Neurologic Outcomes in Newborns With Hypoxic Ischemic Encephalopathy. *Pediatrics*;124:e218-6.