

DAFTAR PUSTAKA

- Abdelhamid I, Lasram K, Meilud G, Ben Halim N, Refi R, Samb A, Abdelhak S, *et al*, 2013, E23K variant in KCNJ11 gene is associated with susceptibility to type 2 diabetes in the Mauritanian population, *Prim. Care Diab*, 345:1-5
- Abed AA, Ayesb BM dan Hamdona OM, 2013, Single Nucleotide Polymorphism E23K of KCNJ11 Gene and Other Risk Factors Associated With Type-2 Diabetes Mellitus In Gaza, *Int. J. Chem. and Life Sciences*, 02 (04) : 1146-52
- Alsmadi O, Al-Rubeaan K, Wakil SM, Imtiaz F, Mohamed G, Al-Saud H, Al-Saud NA, *et al*, 2008, Genetic Study of Saudi Diabetes (GSSD): significant association of the KCNJ11 E23K polymorphism with type 2 diabetes, *Diabetes Metab Res Rev* 24: 137-140.
- American Diabetes Association, 2005, Diagnosis and classification of diabetes mellitus, *Diabetes Care*, 28 (Suppl 1): S37-S42.
- American Diabetes Association, 2010, Standards of medical care in diabetes-2010, *Diabetes Care*, 33 (Suppl 1): S11-S61.
- American Diabetes Association, 2014, Diagnosis and classification of diabetes mellitus, *Diabetes Care*, 34 (Suppl 1): S81-S90.
- Amin S, Arlina dan Karman AS, 2008, *Identifikasi bentuk kearifan lokal masyarakat Maluku Utara dan strategi implementasinya dalam pembangunan*, Laporan penelitian hibah bersaing.
- Ashcroft FM, 2005, ATP-sensitive potassium channelopathies: focus on insulin secretion, *JCI*, 115 : 2047-58.
- Ashcroft FM, dan Rorsman P, 2004, Type 2 diabetes mellitus: not quite exciting enough?, *Hum. Mol. Genet*, 13: R21-R31.
- Baroso I, Luan J, Middelberg RPS, Harding AH, Franks PW, Clayton D, *et al*, 2003, Candidate gene association study in type 2 diabetes indicates a role for genes involved in β -cell function as well as insulin action, *PLoS Biol*, 1(3): 445
- Boodram LG, Miyake K, Hayes MG, Bel GI, dan Cockburn BN, 2011, Association of the KCNJ11 variant E23K with type 2 diabetes in Indo-Trinidadians, *West Indian Med J*, 60 (6) : 604
- Bratanova-Tochkova TK, Cheng H, Daniel S, Gunawardana S, Liu Y, Musa JM, Schermerhorn T, *et al*, 2002, Triggering and Augmentation Mechanisms, Granule Pools, and Biphasic Insulin Secretion, *Diabetes*, 51 (suppl. 1): S83-90).
- Bryan J, Muñoz A, Zhang X, Düfer M, Drews G, Krippeit-Drews P dan Aguilar-Bryan L, 2007, ABCC8 and ABCC9: ABC transporters that regulate K^+ channels, *Pflugers Arch - Eur J Physiol*, 453:703-18.
- Bryan J, Wanda H, Carriles V, Zhao G, Babenko AP, dan Bryan LA, 2004, Toward linking structure with function in ATP-sensitive K^+ channels, *Diabetes*, 53 (Suppl. 3): S104-S112.

- Cejkova P, Novota P, Cerna M, Kolostova K, Novakova D, Kucera P, Novak J, *et al*, 2007, KCNJ11 E23K Polymorphism and Diabetes Mellitus with Adult Onset in Czech Patients, *Folia Biol (Praha)*, 53(5): 173-75.
- Chistiakov DA, Potapov VA, Khodirev DS, Shamkhalova MS, Shestakova MV dan Nosikov VV, 2008, The KCNJ11 E23K and ABCC8 exon 31 variants contribute to susceptibility to type 2 diabetes, glucose intolerance and altered insulin secretion in a Russian population, *Diabetes Metab. Syndr. Clin. Res. Rev*, 2(3): 185-91.
- Dahlan MS, 2013, *Besar Sampel dan Cara Pengambilan Sampel dalam Penelitian Kedokteran dan Kesehatan*, Salemba Medika, Jakarta.
- Evliaoglu O, Sancaktar E, Sogut E, Basarali MK, Uzuncan N, dan Karaca B, 2011, Association of a single nucleotide polymorphism in the SUR1 gene with type 2 diabetes and obesity in Turkish patients, *J Clin Exp Invest*, 2 (2): 161-67
- Ezenwaka C, Kalloo R, Uhlig M, Schwenk R dan Eckel J, 2005, The E23K variant in the Kir6.2 subunit of the ATP-sensitive K⁺ channel does not augment impaired glucose tolerance in Caribbean subjects with a family history of type 2 diabetes, *J Endocrinol*, 185: 439–44.
- Fischer A, Mohlig M, Weickert MO, Schueler R, Martin O, Pfeiffer AFH, *et al*, 2008, KCNJ11 E23K Affects Diabetes Risk and Is Associated With the Disposition Index, *Diabetes Care*, 31: 87-89.
- Florez JC, Jablonski KA, Kahn SE, Franks PW, Dabelea D, Hamman RF, *et al*, 2007, Type 2 Diabetes–Associated Missense Polymorphisms KCNJ11 E23K and ABCC8 A1369S Influence Progression to Diabetes and Response to Interventions in the Diabetes Prevention Program, *Diabetes* 56: 531-536.
- Frizzel JP, 2001, *Handbook of Pathophysiology*, Springhouse Corporation, Philadelphia.
- Gloyn AL, Weedon MN, Owen KR, Turner MJ, Knight BA, Hitman G, *et al*, 2007, Large-Scale Association Studies of Variants in Genes Encoding the Pancreatic β -Cell K_{ATP} Channel Subunits Kir6.2 (KCNJ11) and SUR1 (ABCC8) Confirm That the KCNJ11 E23K Variant Is Associated With Type 2 Diabetes, *Diabetes* 52(2): 568-72.
- Goksel DL, Fischbach K, Duggirala R, Mitchell BD, Lydia AB, Blangero J, *et al*, 1998, Variant in sulfonylurea receptor-1 gene is associated with high insulin concentrations in non-diabetic Mexican Americans: SUR-1 gene variant and hyperinsulinemia, *Hum Genet* 103: 280–85.
- Haghvirdizadeh P, Mohamed Z, Abdullah NA, Haghvirdizadeh P, Haerian MS dan Haerian BS, 2014, Review Article; KCNJ11: Genetic Polymorphisms and Risk of Diabetes Mellitus, *J Diab Res*, 1-9.
- Hani EH, Clement K, Velho G, Vionnet N, Hager J, Philippi A, *et al*, 1997, Genetic Studies of the Sulfonylurea Receptor Gene Locus in NIDDM and in Morbid Obesity Among French Caucasians, *Diabetes* 46(4): 688-94.

- [http://: www.editor@malutpost.co.id/](http://www.editor@malutpost.co.id/) tanggal 11 Juni dan 24 Desember 2013
- <http://www.ncbi.nlm.nih.gov/>
- Jiang YD, Chuang LM, Pei D, Lee YJ, Wei JN, Sung FC dan Chang TJ, 2014, Genetic Variations in the Kir6.2 Subunit (KCNJ11) of Pancreatic ATP-Sensitive Potassium Channel Gene Are Associated with Insulin Response to Glucose Loading and Early Onset of Type 2 Diabetes in Childhood and Adolescence in Taiwan, *Int J Endocrinol.* 1-7.
- Koster JC, Kurata HT, Enkvetchakul D dan Nichols CG, 2008, DEND Mutation in Kir6.2 (KCNJ11) Reveals a Flexible N-Terminal Region Critical for ATP-Sensing of the K_{ATP} Channel, *Biophysical J*, 95: 4689-97.
- Koster JC, Permutt MA, dan Nichols CG, 2005, Perspectives in Diabetes: Diabetes and Insulin Secretion The ATP-Sensitive K^+ Channel (K_{ATP}) Connection, *Diabetes* 54: 3065–72.
- Lasram K, Ben Halim N, Hsouna S, Refi R, Arfa I, Ghazouani W, Jamoussi H, *et al*, 2014, Evidence for Association of the E23K Variant of KCNJ11 Gene with Type 2 Diabetes in Tunisian Population: Population-Based Study and Meta-analysis, *Biomed Res Int.* 265274-9
- Laukkanen O, Pihlajamaki J, Lindstrom J, Eriksson J, Valle TT, Hamalainen H, *et al*, 2004, Polymorphisms of the SUR1 (ABCC8) and Kir6.2 (KCNJ11) genes predict the conversion from impaired glucose tolerance to type 2 diabetes. The Finnish diabetes prevention study, *J Clin Endocrinol Metab*, 89: 6286-90.
- Lebar FM, 1972, *Ethnic groups of insular southeast Asia. Volume 1: Indonesia, Andaman islands, and Madagascar*, Human Relations Area Files Press, New Haven, USA.
- Li YY, 2013, The KCNJ11 E23K gene polymorphism and type 2 diabetes mellitus in the Chinese Han population: a meta-analysis of 6,109 subjects, *Mol Biol Rep*, 40: 141–146.
- Linawati Y, 2009, *Polimorfisme p.R1273R gena Sulfonylurea receptor-1 (SUR1) sebagai faktor risiko diabetes mellitus tipe 2 pada populasi Jawa (tesis)*, Pascasarjana Universitas Gadjah Mada.
- Lyssenko V, Almgren P, Anevski D, Orho-Melander M, Sjogren M, Saloranta C, Tuomi T, *et al*, 2005, Genetic Prediction of Future Type 2 Diabetes, *PLoS Med*, 2(12): e345.
- Lyssenko V, Jonsson A, Almgren P, Pulizzi N, Isomaa B, Tuomi T, *et al*, 2008, Clinical risk factors, DNA variants, and the development of type 2 diabetes, *N Engl J Med*, 359: 2220-32.
- Meloni AR, Deyoung MB dan Lowe C, 2012, GLP-1 receptor activated insulin secretion from pancreatic β -cells: mechanism and glucose dependence, *Diabetes Obes and Metab*, 15: 15-27.
- Melton T, Peterson R, Redd AJ, Saha N, Sofro ASM, Martinson J, dan Stoneking M, 1995, Polynesian genetic affinities with southeast asian populations as identified by mtDNA analysis, *Am J Hum Genet*, 57: 403-414.

- Merentek E, 2006, *Resistensi Insulin pada Diabetes Mellitus tipe 2*, Poliklinik Endokrin Metabolik, Bagian Penyakit Dalam Rumah Sakit Umum Gowa, Makassar.
- Miki T, Nagashima K dan Seino S, 1999, Review: The structure and function of the ATP-sensitive K⁺ channel in insulin-secreting pancreatic β -cells, *J Mol Endocrinol.* 22: 113–23.
- Moore DJ, Gregory JM, Crystal YAK, dan Simmons JH, 2009, Mitigating micro- and macro-vascular complications of diabetes beginning in adolescence, *Vasc Health Risk Manag*, 5: 1015-31.
- Murray RK, Granner DK & Rodwell VW, 2009, *Biokimia Harper*, ed. 27, EGC, Jakarta.
- Nielsen EMD, Hansen L, Carstensen B, Echwald SM, Driysholm T, Glumer C, *et al*, 2003, The E23K variant of kir6.2 associates with impaired post-OGTT serum insulin response and increased risk of type 2 diabetes, *Diabetes* 52: 573-577.
- Nussbaum RL, Roderick RM, dan Huntington FW, 2001, *Thompson & Thompson Genetic in Medicine*, 6th revised edition, Saunders Philadelphia, USA, 9: 79-94.
- Odgerel Z, Lee HS, Erdenebileg N, Gandbold S, Luvsanjamba M, Sambuughin N, Sonomtseren S, *et al*, 2013, Genetic variants in potassium channels are associated with type 2 diabetes in Mongolian population, *J Diabetes*, 4 (3): 238-242.
- Permutt MA, Wasson J, dan Cox N, 2005, Genetic epidemiology of diabetes, *J. Clin Invest*, 115: 1431–39.
- Prabandari AS, 2009, *Polimorfisme p.E23K gena Potassium inwardly rectifying channel sub family J member 11 (KCNJ11) sebagai faktor risiko diabetes mellitus tipe 2 di Yogyakarta (tesis)*, Pascasarjana Universitas Gadjah Mada.
- Proks P, Reimann F, Green N, Gribble F, dan Ashcroft F, 2002, Sulfonylurea Stimulation of Insulin Secretion, *Diabetes*, 51 (Suppl 3): S368-S376.
- Radha KS, Vimalaswaran R, Deepa dan Mohan V, 2003, The genetic of diabetes mellitus, *Indian J Med Res* 117, pp 225-38.
- Rastegari A, Rabbani M, Sadeghi HM, Imani EF, Hasanzadeh A, dan Moazen F, 2015, Association of KCNJ11 (E23K) gene polymorphism with susceptibility to type 2 diabetes in Iranian patients, *Adv Biomed Res*, 4:1
- Reis AF, Ye WZ, Daniele DL, Christine BC, Timsit J, dan Velho G, 2000, Association of a variant in exon 31 of the sulfonylurea receptor 1 (SUR1) gene with type 2 diabetes mellitus in French Caucasians, *Hum Genet*, 107: 138–44.
- Report of the Expert committee on the diagnosis and clasification of diabetes mellitus, 2003, *Diabetes Care*, 37 (Suppl 1): S5-S20.
- Riedel MJ, Boora P, Steckley D, de Vries G, dan Light PE, 2003, Kir6.2 Polymorphisms Sensitize β -Cell ATP-Sensitive Potassium Channels to Activation by Acyl CoAs : A Possible Cellular Mechanism for Increased Susceptibility to Type 2 Diabetes?, *Diabetes* 52(10): 2630–35.

- Rissanen J, Markkanen A, Karkkainen P, Pihlajamaki J, Kekalainen P, Mykkanen L, *et al*, 2000, Sulfonylurea receptor 1 gene variants are associated with gestational diabetes and type 2 diabetes but not with altered secretion of insulin, *Diabetes care*, 23(1): 70-73.
- Sakamoto Y, Inoue H, Keshavarz P, Miyawaki K, Yamaguchi Y, Moritani M, *et al*, 2007, SNPs in the KCNJ11-ABCC8 gene locus are associated with type 2 diabetes and blood pressure levels in the Japanese population, *J Hum Genet* 52: 781-93.
- Sastroasmoro S dan Ismael S, 2011, Dasar-Dasar Metodologi Penelitian Klinis, Ed 2, CV Sagung Seto, Yogyakarta.
- Schaid DJ, Batzler AJ, Jenkins GD, dan Hildebrandt MAT, 2006, Exact Tests of Hardy-Weinberg Equilibrium and Homogeneity of Disequilibrium across Strata, *Am. J. Hum. Genet.* 79:1071–80.
- Schwanstecher C, Meyer U, dan Schwanstecher M, 2002, Kir6.2 Polymorphism Predisposes to Type 2 Diabetes by Inducing Overactivity of Pancreatic β -Cell ATP-Sensitive K^+ Channels, *Diabetes* 51(3):875-79.
- Schwanstecher C dan Schwanstecher M, 2002, Nucleotida Sensitivity of Pancreatic ATP-Sensitive of Pottasium Channels and Type 2 Diabetes, *Diabetes*, 51, Suppl 3: S358-S362.
- Sesti G, Federici M, Hribal MI, Lauro D, Sbraccia P dan Lauro R, 2006, Defects of the insulin receptor substrate (IRS) system in human metabolic disorders, *FASEB J*, 15: 2099-111.
- Shaat N, Almgren P, Ekelund M, Berntorp K, Lernmark A, Groop L dan Ivarsson S, 2005, Association of the E23K polymorphism in the KCNJ11 gene with gestational diabetes mellitus, *Diabetologia* 48: 2544–51.
- Sunita R, Sadewa AH dan Farmawati A, 2015, Lower HOMA- β values are detected among individuals with variant of E23K polymorphism of potassium inwardly-rectifying channel, subfamily J, member 11 (*KCNJ11*) gene, *Egyptian J Med Hum Gen.* 16: 227-31.
- Thevenod F, 2008, Pathophysiology of diabetes mellitus type 2: Roles of obesity, Insulin resistance and β -cell dysfunction, *F. Diab.* Basel, 19 pp1-18.
- UPTD Diabetes Center, 2013, *Profil kerja UPTD Diabetes Center 2013*, Dinas Kesehatan Kota Ternate.
- Wigginton JE, Cutler DJ, dan Abecasis GR, 2005, A Note on Exact Tests of Hardy-Weinberg Equilibrium, *Am. J. Hum. Genet.* 76:887–883.
- Wild S, Roglic G, Green A, Sicre R, dan King H, 2004, Global Prevalence of Diabetes: Estimates for the year 2000 and projections for 2030, *Diabetes Care*, 27: 1047-53.
- World Health Organization, 2004, Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies, *Lancet*, 363: 157-63.
- Yokoi N, Kanamori M, Horikawa Y, Takeda J, Sanke T, Furuta H dan Nanjo K *et al.* 2006, Association Studies of Variants in the Genes Involved in Pancreatic β -Cell Function in Type 2 Diabetes in Japanese Subjects, *Diabetes* 55:2379-86.

Zhou D, Zhang D, Liu Y, Zhao T, Chen Z, Liu Z, Yu L et al. 2009, The E23K variation in the KCNJ11 gene is associated with type 2 diabetes in Chinese and East Asian population, *J Hum Genet*, 54: 433–35.