

## REFERENCES

- Angelidaki, I., Ellegaard, L., Ahring, B.K. 1999. A comprehensive model of anaerobic bioconversion of complex substrates to biogas. *Biotechnol. Bioeng.* 63: 363-372.
- Angelidaki, I., Ellegaard, L., Sorensen, A.H., Schmidt, J.E. (2002). *Anaerobic processes*. In: Angelidaki I, editor. Environmental biotechnology. Institute of Environment and Resources. Technical University of Denmark (DTU). pp. 1-114.
- Arias, C, R., Burns, J, K., Friedrich, L, M., Goodrich, R, M., Parish, E, P. 2002. Yeast Species Associated with Orange Juice: Evaluation of Different Identification Methods. *Applied and environmental microbiology*. 68(4): 1955-1961.
- Aslanzadeh, S., Rajendran, K., Jeihanipour, A., Taherzadeh, M. 2013. The effect of effluent recirculation in a semi-continuous two-stage anaerobic digestion system. *Energies*. 6: 2966-2981.
- Beccari, M., Bonemazzi, F., Majone, M., Riccardi, C. 1996. Interaction between acidogenesis and methanogenesis in the anaerobic treatment of olive oil mill effluents. *Water Res.* 30: 183-189
- Boluda-Aguilar, M., and López-Gómez, A. 2013. Production of bioethanol by fermentation of lemon (*Citrus limon L.*) peel wastes pretreated with steam explosion. *Ind.Crops Prod.* 41: 188-197.
- Bouallagui, H., Ben Cheikh, R., Marouani, L., Hamdi, M. 2003. Mesophilic biogas production from fruit and vegetable waste in a tubular digester. *Bioresour. Technol.* 86: 85-89.
- C, De Blas., G, G, Mateos., P, García-Rebollar. 2010. *FEDNA tables of composition and nutritive value of foods for the manufacturing of compound feeds (3<sup>rd</sup> Edition)*. Madrid: Fundación Española para el Desarrollo de la Nutrición Animal.
- Cohen, A., Breure, A.M., van Andel, J.G., van Deursen, A. 1982. Influence of phase separation on the anaerobic digestion of glucose, 2. Stability and kinetic responses to shock loadings. *Water Res.* 16: 449-455.
- Converti, A., Delborghi, A., Zilli, M., Arni, S., Delborghi, M. 1999. Anaerobic digestion of the vegetable fraction of municipal refuses. Mesophilic versus thermophilic conditions. *Bioprocess Eng.* 21: 371-376.
- Crawshaw, R. 2003. *Co-product feeds: animal feeds from the food and drinks industries*. Nottingham: Nottingham University Press.
- Demirel, B., and Yenigün, O. 2002. Two-phase anaerobic digestion processes: A review. *J. Chem. Technol. Biotechnol.* 77: 743-755.
- Deublein, D., and A, Steihauser. 2008. *Biogas from waste and renewable resources*. Weinheim: Wiley-VCH.
- Drosg, Bernhard. 2013. *Process monitoring in biogas plants*. IEA Bioenergy.
- FAO-Food and Agriculture Organization. 2016. *Citrus Fruit Statistics*. FAO. Rome.
- F. Bakkali, S. Averbek, D. Averbek, M. Idaomar. 2008. Biological effects of essential oils-a review, *Food Chem. Toxicol.* 46: 446-475.

- Forgács, G. 2012. *Biogas production from citrus waste and chicken feather: pretreatment and co-digestion*. Chalmers University of Technology: Gothenburg, Sweden.
- Forster-Carneiro, T., Pérez, M., Romero, L.I. 2008. Anaerobic digestion of municipal solid wastes: dry thermophilic performance. *Bioresour. Technol.* 99: 8180-8184.
- Garcia-Peña, E.I., Parameswaran, P., Kang, D.W., Canul-Chan, M., Krajmalnik-Brown, R. 2011. Anaerobic digestion and co digestion of vegetable and fruit residues: process and microbial ecology. *Bioresour. Technol.* 102: 9447–9455.
- Ghosh, S., Ombregt, J.P., Pipyn, P. 1985. Methane production from industrial wastes by two-phase anaerobic digestion. *Water Res.* 19: 1083–1088.
- Grohmann, K., Cameron, R.G., Buslig, B.S. 1995. Fractionation and pretreatment of orange peel by dilute acid hydrolysis. *Bioresour. Technol.* 54: 129–141.
- Hansen, T.L., Schmidt, J.E., Angelidaki, I., Marca, E., Jansen, J.I., Mosbaek, H., Christensen, T.H. 2004. Method for determination of methane potentials of solid organic waste. *Waste Manag.* 24(4): 393-400.
- Kaparaju, P, L, N., Rintala, J, A. 2006. Thermophilic anaerobic digestion of industrial orange waste. *Environ. Technol.* 27: 623–633.
- Kapdan, I.K., and Kargi Fikret. 2006. Bio-hydrogen production from waste materials. *Enzyme Microb. Technol.* 26: 569-582.
- Koppar, A., and Pullammanappallil, P. 2013. Anaerobic digestion of peel waste and wastewater for on site energy generation in a citrus processing facility. *Energy.* 60: 62–68.
- Kroeker, E.J., Schulte, D.D., Sparling, A.B., Lapp, H.M. 1979. Anaerobic treatment process stability. *J Water Pollut Control Fed.* 51:718–727.
- Landine, R.C., Brown, G.J., Cocci, A.A., Virara, H. 1983. Anaerobic treatment of high strength, high solids potato waste. *Agric. Wastes.* 17: 111–123.
- Lane, A.G. 1984. Anaerobic digestion of orange peel. *Food Technol.* 36: 125–127.
- Lin, J., Zuo, J., Gan, L., Li, P., Liu, F., Wang, K., Chen, L., Gan, H. 2011. Effects of mixture ratio on anaerobic co-digestion with fruit and vegetable waste and food waste of China. *J. Environ. Sci.* 23 (8): 1403-1408.
- Liu, Z. 1985. *Oxidation-reduction potential*. In: Yu T, editor. Physical chemistry of paddy soils. Beijing: Science Press. p 1-26.
- Lun, S.Y., Wu, J., Chen, J. 1995. Anaerobic waste treatment: efficient separation of the acid and methane forming phases using two UASB reactors. *Process Biochem.* 30: 523–529.
- Madigan, M., Martinko, J., Dunlap, P., Clark, D. 2008. Brockbiology of microorganisms (12<sup>nd</sup> edition). San Fransisco: Benjamin cummings.
- Mamma, D., Kourtoglou, E., Christakopoulos, P. 2008. Fungal multienzyme production on industrial by-products of the citrus-processing industry. *Bioresour. Technol.* 99: 2373–2383.
- Mamma, D., and Christakopoulos, P. 2014. Biotechnological potential of citrus peels. In: Microbes in Process. *Nova Science Publishers Inc:* 59–91.
- Marriott, P, J., Shellie, R., Cornwell, C. 2001. Gas chromatographic technologies for the analysis of essential oils. *J. Chromatogr. A.* 936: 1–22.

- Martín, M., A., Siles, J., A., Chica, A., F., Martín, A. 2010. Biomethanization of orange peel waste. *Bioresour. Technol.* 101: 8993–8999.
- Martín, M., A., Fernández, R., Serrano, A., Siles, J., A. 2013. Semi-continuous anaerobic co-digestion of orange peel waste and residual glycerol derived from biodiesel manufacturing. *Waste Manage.* 33: 1633–1639.
- Mizuki, E., T., Akao, T., Saruwatari. 1990. Inhibitory effect of *Citrus unshu* peel on anaerobic digestion. *Biol. Wastes.* 33(3): 161–168.
- Mota, V.T., Santos, F. S., Amaral, M, C, S. 2013. Two-stage anaerobic membrane bioreactor for the treatment of sugarcane vinasse: Assessment on biological activity and filtration performance. *Bioresource Technology.* 146: 494–503.
- NASS.2011. <[http://www.nass.usda.gov/Publications/Reports\\_by\\_Release\\_Day](http://www.nass.usda.gov/Publications/Reports_by_Release_Day)>. Accessable in August 2016.
- Negro, V., Mancini, G., Ruggeri, B., Fino, D. 2016. Citrus waste as feedstock for bio-based products recovery: Review on limonene case study and energy valorization. *Bioresource Technology.* 214: 806–815
- Pohland, F.G., and Ghosh, S. 1971. Developments in anaerobic stabilization of organic wastes-The two-phase concept. *Environ. Lett.* 1: 255–266.
- Pourbafrani, M., Forgács, G., Horváth, I.S., Niklasson, C., Taherzadeh, M.J., 2010. Production of biofuels, limonene and pectin from citrus wastes. *Bioresour. Technol.* 101: 4246–4250.
- Prawira, W., Murto, M., Read, J.S., Mattiason, B. 2005. Profile of hydrolases and biogas production during two-stage mesophilic anaerobic digestion of solid potato waste. *Process Biochem.* 40 (9): 2945–2952.
- Reineccius, G. 2006. Flavor Chemistry and Technology. Boca Raton, FL: CRC Press.
- Rieger, C, H., and Weiland, P. 2006. Prozessstörungen frühzeitig erkennen. *Biogas Journal.* 4: 18–20.
- Ruiz, B., and Flotats, X. 2014. Citrus essential oils and their influence on the anaerobic digestion process: an overview. *Waste Manage.* 34: 2063–2079.
- Ruiz, B., and Flotats, X. 2015. Effect of limonene on batch anaerobic digestion of citrus peel waste. *Biochemical Engineering Journal.* 109: 9–18.
- Ruynal, J., Delgenes, J.P., Moletta, R. 1998. Two phase anaerobic digestion of solid waste by a multiple liquefaction reactors process. *Bioresour. Technol.* 65: 97–103.
- Siles, J.A., Vargas, F., Gutiérrez, M.C., Chica, A.F., Martín, M.A., 2016. Integral valorisation of waste orange peel using combustion, biomethanisation and cocomposting technologies. *Bioresour. Technol.* 211: 173–182.
- Sluiter, A., Hames, B., Ruiz, R., Scarlata, C., Sluiter, J., Templeton, D. 2008. *Determination of ash in biomass.* Colorado: National Renewable Energy Laboratory.
- S, Sánchez-Segado, L, J, Lozano, A, P, de los Ríos, F,J, Hernández-Fernández, C, Godínez, D, Juan. 2012. Process design and economic analysis of a hypothetical bioethanol production plant using carob pod as feedstock. *Bioresour. Technol.* 104: 324–328.
- Verrier, D., Ray, F., Albagnac, G. 1987. Two-phase methanization of solid vegetable wastes. *Biol. Wastes.* 22: 163–167.

- Weiland, P. 2010. Biogas production: current state and perspectives. *Applied Microbiology and Biotechnology*. 85(4): 849-860.
- Wikandari, R., Gudipudi, S., Pandiyan, I., Millati, R., Taherzadeh, M.J. 2013. Inhibitory effects of fruit flavors on methane production during anaerobic digestion. *Bioresour. Technol.* 145: 188–192.
- Wikandari, R., Millati, R., Cahyanto, M.N., Taherzadeh, M.J. 2014. Biogas production from citrus waste by membrane bioreactor. *Membranes (Basel)*. 4: 596–607.
- Wikandari, Rahma. 2014. *Effect of fruit flavors on anaerobic digestion: inhibitions and solutions*. University of Borås: Borås, Sweden.
- Wilkins, M.R., Suryawati, L., Maness, N.O., Chrz, D. 2007. Ethanol production by *Saccharomyces cerevisiae* and *Kluyveromyces marxianus* in the presence of orange-peel oil. *World J Microbiol Biotechnol.* 23: 1161–1168.
- Zhang, R., El-Mashad, H.M., Hartman, K., Wang, F., Liu, G., Choate, C., Gamble, P. 2007. Characterization of food waste as feedstock for anaerobic digestion. *Bioresour. Technol.* 98: 929-935.