UNIVERSITAS GADJAH MADA
EVALUATION OF VOLATILE FATTY ACIDS IN THE FIRST STAGE OF THE TWO STAGE ANAEROBIC DIGESTION OF CITRUS WASTE
ERVINA DWI PERMANASARI, Dr. Ria Millati, S.T., M.T. dan Dr. Ir. M. Nur Cahyanto, M.Sc.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

REFERENCES

Angelidaki, I., Ellegaard, L., Ahring, B,K. 1999. A comprehensive model of anaerobic bioconversion of complex substrates to biogas. *Biotechnol. Bioeng*. 63: 363-372.

Angelidaki, I., Ellegaard, L., Sorensen, A.H., Schmidt, J.E. (2002). *Anaerobic processes*. In: Angelidaki I, editor. Environmental biotechnology. Institute of Environment and Resources. Technical University of Denmark (DTU). pp. 1-114.

Arias, C, R., Burns, J, K., Friedrich, L, M., Goodrich, R, M., Parish, E, P. 2002. Yeast Species Associated with Orange Juice: Evaluation of Different Identification Methods. *Applied and environmental microbiology*. 68(4): 1955-1961.

Aslanzadeh, S., Rajendran, K., Jeihanipour, A., Taherzadeh, M. 2013. The effect of effluent recirculation in a semi-continuous two-stage anaerobic digestion system. *Energies*. 6: 2966-2981.

Beccari, M., Bonemazzi, F., Majone, M., Riccardi, C. 1996. Interaction between acidogenesis and methanogenesis in the anaerobic treatment of olive oil mill effluents. *Water Res.* 30: 183–189

Boluda-Aguilar, M., and López-Gómez, A. 2013. Production of bioethanol by fermentation of lemon (*Citrus limon L*.) peel wastes pretreated with steam explosion. *Ind.Crops Prod.* 41: 188-197.

Bouallagui, H., Ben Cheikh, R., Marouani, L., Hamdi, M. 2003. Mesophilic biogas production from fruit and vegetable waste in a tubular digester. *Bioresour. Technol*. 86: 85–89.

C, De Blas., G, G, Mateos., P, García-Rebollar. 2010. *FEDNA tables of composition and nutritive value of foods for themanufacturing of compound feeds (3rd Edition).* Madrid: Fundación Espa˜nola para elDesarrollo de la Nutrición Animal.

Cohen, A., Breure, A,M., van Andel, J,G., van Deursen, A. 1982. Influence of phase separation on the anaerobic digestion of glucose, 2. Stability and kinetic responses to shock loadings. *Water Res*. 16: 449–455.

Converti, A., Delborghi, A., Zilli, M., Arni, S., Delborghi, M. 1999. Anaerobic digestion of the vegetable fraction of municipal refuses. Mesophilic versus thermophilic conditions. *Bioprocess Eng.* 21: 371–376.

Crawshaw, R. 2003. *Co-product feeds: animal feeds from the food and drinks industries.* Nottingham: Nottingham University Press.

Demirel, B., and Yenigün, O. 2002. Two-phase anaerobic digestion processes: A review. *J. Chem. Technol. Biotechnol.* 77: 743–755.

Deublein, D., and A, Steinhauser. 2008. Biogas from waste and renewable resources. Weinheim: Wiley-VCH.

Drosg, Bernhard. 2013. *Process monitoring in biogas plants*. IEA Bioenergy.

FAO-Food and Agriculture Organization. 2016. Citrus Fruit Statistics. FAO. Rome.

F. Bakkali, S. Averbeck, D. Averbeck, M. Idaomar. 2008. Biological effects of essentialoils-a review, Food Chem. Toxicol. 46: 446–475.

UNIVERSITAS GADJAH MADA
EVALUATION OF VOLATILE FATTY ACIDS IN THE FIRST STAGE OF THE TWO STAGE ANAEROBIC DIGESTION OF CITRUS WASTE
ERVINA DWI PERMANASARI, Dr. Ria Millati, S.T., M.T. dan Dr. Ir. M. Nur Cahyanto, M.Sc.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Forgács, G. 2012. *Biogas production from citrus waste and chicken feather: pretreatment and co-digestion*. Chalmers University of Technology: Gothenburg, Sweden.

Forster-Carneiro, T., Pérez, M., Romero, L.I. 2008. Anaerobic digestion of municipal solid wastes: dry thermophilic performance. *Bioresour. Technol*. 99: 8180-8184.

Garcia-Peña, E.I., Parameswaran, P., Kang, D.W., Canul-Chan, M., Krajmalnik-Brown, R. 2011. Anaerobic digestion and co digestion of vegetable and fruit residues: process and microbial ecology. *Bioresour. Technol*. 102: 9447–9455.

Ghosh, S., Ombregt, J,P., Pipyn, P. 1985. Methane production from industrial wastes by two-phase anaerobic digestion. *Water Res*. 19: 1083–1088.

Grohmann, K., Cameron, R.G., Buslig, B.S. 1995. Fractionation and pretreatment of orange peel by dilute acid hydrolysis. *Bioresour. Technol.* 54: 129–141.

Hansen, T,L., Schmidt, J,E., Angelidaki, I., Marca, E., Jansen, J,I., Mosbaek, H., Christensen, T,H. 2004. Method for determination of methane potentials of solid organic waste. *Waste Manag*. 24(4): 393-400.

Kaparaju, P, L, N., Rintala, J, A. 2006. Thermophilic anaerobic digestion of industrial orange waste. *Environ. Technol*. 27: 623–633.

Kapdan, I.K., and Kargi Fikret. 2006. Bio-hydrogen production from waste materials. *Enzyme Microb. Technol*. 26: 569-582.

Koppar, A., and Pullammanappallil, P. 2013. Anaerobic digestion of peel waste and wastewater for on site energy generation in a citrus processing facility. *Energy*. 60: 62–68.

Kroeker, E,J., Schulte, D,D., Sparling, A,B., Lapp, H,M. 1979. Anaerobic treatment process stability. *J Water Pollut Control Fed*. 51:718–727.

Landine, R.C., Brown, G.J., Cocci, A.A., Virara, H. 1983. Anaerobic treatment of high strength, high solids potato waste. *Agric. Wastes*. 17: 111–123.

Lane, A.G. 1984. Anaerobic digestion of orange peel. *Food Technol*. 36: 125–127.

Lin, J., Zuo, J., Gan, L., Li, P., Liu, F., Wang, K., Chen, L., Gan, H. 2011. Effects of mixture ratio on anaerobic co-digestion with fruit and vegetable waste and food waste of China. *J. Environ. Sci.* 23 (8): 1403-1408.

Liu, Z. 1985. *Oxidation-reduction potential*. In: Yu T, editor. Physical chemistry of paddy soils. Beijing: Science Press. p 1-26.

Lun, S,Y., Wu, J., Chen, J. 1995. Anaerobic waste treatment: efficient separation of the acid and methane forming phases using two UASB reactors. *Process Biochem*. 30: 523–529.

Madigan, M., Martinko, J., Dunlap, P., Clark, D. 2008. Brockbiology of microorganisms (12nd edition). San Fransisco: Benjamin cummings.

Mamma, D., Kourtoglou, E., Christakopoulos, P. 2008. Fungal multienzyme production on industrial by-products of the citrus-processing industry.*Bioresour.Technol*. 99: 2373–2383.

Mamma, D., and Christakopoulos, P. 2014. Biotechnological potential of citrus peels. In: Microbes in Process. *Nova Science Publishers Inc*: 59–91.

Marriott, P, J., Shellie, R., Cornwell, C. 2001. Gas chromatographic technologies for the analysis of essential oils. *J. Chromatogr. A.* 936: 1–22.

UNIVERSITAS
GADJAH MADA
EVALUATION OF VOLATILE FATTY ACIDS IN THE FIRST STAGE OF THE TWO STAGE ANAEROBIC DIGESTION OF CITRUS WASTE
ERVINA DWI PERMANASARI, Dr. Ria Millati, S.T., M.T. dan Dr. Ir. M. Nur Cahyanto, M.Sc.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Martín,M, A., Siles, J, A., Chica, A, F., Martín, A. 2010. Biomethanization of orange peel waste. *Bioresour. Technol*. 101: 8993–8999.

Martín, M, A., Fernández, R., Serrano, A., Siles, J, A. 2013. Semi-continuous anaerobic co-digestion of orange peel waste and residual glycerol derived from biodiesel manufacturing. *Waste Manage*. 33: 1633–1639.

Mizuki, E., T, Akao., T, Saruwatari. 1990. Inhibitory effect of *Citrus unshu* peel on anaerobic digestion. *Biol. Wastes.* 33(3): 161-168.

Mota, V,T., Santos, F, S., Amaral, M, C, S. 2013. Two-stage anaerobic membrane bioreactor for the treatment of sugarcane vinasse: Assessment on biological activity and filtration performance. *Bioresource Technology*. 146: 494–503.

NASS.2011. <http://www.nass.usda.gov/Publications/Reports_by_Release_Day>. Accessable in August 2016.

Negro,V., Mancini, G., Ruggeri, B., Fino, D. 2016. Citrus waste as feedstock for bio-based products recovery: Review on limonene case study and energy valorization. *Bioresource Technology*. 214: 806–815

Pohland, F.G., and Ghosh, S. 1971. Developments in anaerobic stabilization of organic wastes-The two-phase concept. *Environ. Lett.* 1: 255–266.

Pourbafrani, M., Forgács, G., Horváth, I.S., Niklasson, C., Taherzadeh, M.J., 2010. Production of biofuels, limonene and pectin from citrus wastes. *Bioresour. Technol*. 101: 4246–4250.

Prawira, W., Murto, M., Read, J.S., Mattiason, B. 2005. Profile of hydrolases and biogas production during two-stage mesophilic anaerobic digestion of solid potato waste. *Process Biochem*. 40 (9): 2945-2952.

Reineccius, G. 2006. Flavor Chemistry and Technology. Boca Raton, FL: CRC Press.

Rieger, C, H., and Weiland, P. 2006. Prozessstörungern frühzeitig erkennen. *Biogas Journal*. 4: 18-20.

Ruiz, B., and Flotats, X. 2014. Citrus essential oils and their influence on the anaerobic digestion process: an overview. *Waste Manage*. 34: 2063–2079.

Ruiz, B., and Flotats, X. 2015. Effect of limonene on batch anaerobic digestion of citrus peel waste. *Biochemical Engineering Journal.* 109: 9–18.

Ruynal, J., Delgenes, J.P., Moletta, R. 1998. Two phase anaerobic digestion of solid waste by a multiple liquefaction reactors process. *Bioresour. Technol*. 65: 97–103.

Siles, J.A., Vargas, F., Gutiérrez, M.C., Chica, A.F., Martín, M.A., 2016. Integral valorisation of waste orange peel using combustion, biomethanisation and cocomposting technologies. *Bioresour. Technol.* 211: 173–182.

Sluiter, A., Hames, B., Ruiz, R., Scarlata, C., Sluiter, J., Templeton, D. 2008. *Determination of ash in biomass*. Colorado: National Renewable Energy Laboratory.

S, Sánchez-Segado., L, J, Lozano., A, P, de los Ríos., F,J, Hernández-Fernández., C, Godínez., D, Juan. 2012. Process design and economic analysis of a hypotheticalbioethanol production plant using carob pod as feedstock. *Bioresour. Technol*. 104: 324–328.

Verrier, D., Ray, F., Albagnac, G. 1987. Two-phase methanization of solid vegetable wastes. *Biol. Wastes.* 22: 163–167.

UNIVERSITAS
GADJAH MADA
EVALUATION OF VOLATILE FATTY ACIDS IN THE FIRST STAGE OF THE TWO STAGE ANAEROBIC DIGESTION OF CITRUS WASTE
ERVINA DWI PERMANASARI, Dr. Ria Millati, S.T., M.T. dan Dr. Ir. M. Nur Cahyanto, M.Sc.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Weiland, P. 2010. Biogas production: current state and perspectives. *Applied Microbiology and Biotechnology*. 85(4): 849-860.

Wikandari, R., Gudipudi, S., Pandiyan, I., Millati, R., Taherzadeh, M.J. 2013. Inhibitory effects of fruit flavors on methane production during anaerobic digestion. *Bioresour. Technol*. 145: 188–192.

Wikandari, R., Millati, R., Cahyanto, M.N., Taherzadeh, M.J. 2014. Biogas production from citrus waste by membrane bioreactor. *Membranes (Basel)*. 4: 596–607.

Wikandari, Rahma. 2014. *Effect of fruit flavors on anaerobic digestion: inhibitions and solutions*. University of Borås: Borås, Sweden.

Wilkins, M.R., Suryawati, L., Maness, N.O., Chrz, D. 2007. Ethanol production by *Saccharomyces cerevisiae* and *Kluyveromyces marxianus* in the presence of orange-peel oil. *World J Microbiol Biotechnol.* 23: 1161–1168.

Zhang, R., El-Mashad, H,M., Hartman, K., Wang, F., Liu, G., Choate, C., Gamble, P.2007. Characterization of food waste as feedstock for anaerobic digestion. *Bioresour. Technol*. 98: 929-935.

.