

DAFTAR PUSTAKA

Aamir, S., Jahangir, S. F., & Farooq, S. (2009). Family Functioning Among Depressive and Dissociative (Conversion) Patients. *Journal of the College of Physicians and Surgeons Pakistan*, 19(5), 300–303.

Al-muzien, N. A., & Al-sowielem, A. L. S. (2014). Prevalence of depression in diabetics attending primary healthcare centers in the Eastern Province of Saudi Arabia. *Journal Bahrain Med Soc*, 25(1), 14–18.

American College of Endocrinology. (2002). American College of Endocrinology Consensus Statement on Guidelines for Glycemic Control. *Endocrine Practice* 8(1) : 5-11., 8(1), 5–11.

American Diabetic Association. (2010). Standard of Medical Care in Diabetes. *Diabetes Care*, 33(Suppl 1), 511–561.

American Diabetic Assosiation. (2015). Standard of Medical Care in Diabetes. *Diabetes Care*, 38(1), 510–93.

Anderson, R. J., Freedland, K. E., Clouse, R. E., & Lustman, P. J. (2001). The Prevalence of Comorbid Depression. *Diabetes Care*, 24(6), 1069–1078.

Apostol-Nicodemus, L., & Umali, M. (2014). Family assessment tools and their application in family practice. In *Textbook of family medicine, principle, concepts, practice, and context*. C&E Publishing, Inc and the Philipphine Academy of Family Physicians.

Asdie, A. H. . H. . (2000). *Patogenesis dan terapi diabetes melitus tipe 2*. Yogyakarta: Medika Fakutas Kedokteran UGM.

Asghar, S., Hussain, A., Ali, S. M., Khan, A. K., & Magnusson, A. (2007). Prevalence of depression and diabetes: a population-based study from rural Bangladesh. *Diabet Medicine*, 24(8), 872–7.

Azwar, A. (1997). *Pengantar Pelayanan Dokter Keluarga*. Yayasan Penerbitan Ikatan Dokter Indonesia.

Badan Penelitian dan Pengembangan Departemen Kesehatan RI. (2013). *Riset KESEHATAN DASAR*.

Bastyr EJ III, Stuart CA, Brodows RG, *et al* (IOEZ S. G. 2003. (2003). Therapy focused in lowering postprandial glucose, not fasting glucose, may be superior for lowering HbA1c. (23) : 1236-41. *Diabetes Care*, 23, 1236–41.

- Beaton, D., Bombardier, C., Guillemin, F., Ferraz, M. (2000). Guidelines for the process of cross-cultural adaptation of self-report measures. *SPINE Lippincott Williams & Wilkins, Inc*, 25(24), 3186–91.
- Bjelland, I., Dahl, A. A., Tangen, T., & Neckelmann, D. (2002). The validity of the Hospital Anxiety and Depression Scale An updated literature review, 52, 69–77.
- Black, S. A., Markides, K. S., & Ray, L. A. (2003). of Adverse Health Outcomes in Older Mexican Americans With Type 2. *Diabetes Care* 26, 26(10), 2822–2828.
- Bruce DG, Davis WA, Starkstein SE, D. T. 2005. (2005). A prospective study of depression and mortality in patients with type II diabetes : the fremantle diabetes study. *Diabetologia*, 48(12), 2532–9.
- Chan, J. C. N., Malik, V., Jia, W., Kadowaki, T., Yajnik, C. S., Yoon, K. H., & Hu, F. B. (2009). Diabetes in Asia: epidemiology, risk factors, and pathophysiology. *JAMA*, 301(20), 2129–2140.
- Chaves, C., Amaral, O., Nelas, P., Coutinho, E., & Dionisio, R. (2014). PL PL. *Procedia - Social and Behavioral Sciences*, 113, 62–67.
<http://doi.org/10.1016/j.sbspro.2014.01.011>
- Collins, M. M., Corcoran, P., & Perry, I. J. (2009). Psychology Anxiety and depression symptoms in patients with diabetes. *Diabetic Medicine* 26: 153-61, 26, 153–61.
<http://doi.org/10.1111/j.1464-5491.2008.02648.x>
- Davenport, Liam. (2016). J-shaped link between diabetes duration and depression. *Maturitas*. Published online January 11
- Dieren, S., J.W.J., B., Schouw, & Yvonne T. van der, D. E. G. and B. N. (2010). The global burden of diabetes and its complication: an emerging pandemic. *European Journal of Cardiovascular Prevention & Rehabilitation*, 17.
- Dimatteo, M. R., Lepper, H. S., & Croghan, T. W. (2000). Depression Is a Risk Factor for Noncompliance With Medical Treatment, *Arch Intern Med*. 160, 2101–7.
- Dinas Kesehatan Kabupaten Bantul. (2013). *Profil Kesehatan Kabupaten Bantul 2013*.
- de Groot, M., Anderson, R., Freedland, K. E., Clouse, R. E., Lustman, P. J. (2001). Association of depression and diabetes complications: a meta-analysis. *Psychosom. Med.* 63 (4) : 619-30.
- Eaton, W. W., Armenia, H., Gallo, J. (1996). Depression and risk for onset of type II diabetes. *Diabetes Care*, 19(10), 1097 – 102.

- Egede, L.E., Ellis, Charles. (2010). Diabetes and depression: Global perspectives. *Diabetes Research and Clinical Practice*. 87:302-312.
- Egede, L. E., Zheng, D., & Simpson, K. (2002). Comorbid Depression is Associated With Increased Health Care Use and and Expenditures in Individuals With Diabetes. *Diabetes Care*, 25(3).
- Egede, L.E., Zheng, D. (2003). Independent factors associated with major depressive disorder in a national sample of individuals with diabetes. *Diabetes Care*, 26, 104–11.
- Fauci, A. S., Kasper, D. L., Longo, D. L., E, B., Hauser, S. H., Jameson, J., Loscalzo. (2008). *Harrison's Internal Medicine, 17th ed.* Mc Grawhills Companies.
- Gan, G. L., Azwar, A., & Wonodirekso, S. (2004). *A Primer on Family Medicine Practice*. Singapore International Foundation.
- Gardner, W., Nutting, P. A., Kelleher, K. J., Werner, J. J., Tillman, F., Stewart, L., ... Orzano, A. J. (2001). Does the family apgar effectively measure family functioning? *The Journal of Family practice*2, 50(1), 19–25.
- Ghazali, M., Sastromiharjo, S., Soedjarwo, S., Soelaryo, T., Pramulyo, H. 2014. (2014). Studi cross sectional dalam Dasar-dasar metodologi penelitian klinis. *Sagung Seto*.
- Ghazanfari, Z., Haghdooost, A. A., Alizadeh, S. M., & Atapour, J. (2010). IJPM A Comparison of HbA1c and Fasting Blood Sugar Tests in General Population, 1(3), 187–194.
- Golden, S.H., Lazo, M., Carnrthon, M., Bertoni, A.G., Schreiner, P.J., Roux, A. V. *et al.* (2008). Examining a bidirectional association between depressive symptoms and diabetes. *JAMA*, 299(23), 2751–59.
- Goldenberg, I., & Goldenberg, H. (1980). *Family therapy*. Brooks/Cole publishing comp. Saint Louis.
- Guyen, S., Matfin, G., & Kuenzi, J. A. (2009). *Diabetes mellitus and metabolic syndromes. Pathophysiology: concept of altered health states 8th edition*. Lippincott Williams and Wilkins.
- International Diabetes Federation. (2009). *Diabetes atlas 4th edition*.
- Iwasaki, Y., Nishinyama, M., Taguchi, T., Asai, M., Yoshida, M., Kambayashi, M., Terada, Y., Hashimoto, K., (2009). Insulin exhibits short-term anti-inflammatory but longterm proinflammatory effects in vitro. *Mol.Cell.Endocrinol.*298: 25-32.

- Jeong, P., Roy, J., Saz, P., Marcos, G., Lobo, A., & Investigators, Z. (2006). Prevalent and incident depression in community-dwelling elderly persons with diabetes mellitus: result from ZARADEMP project. *Diabetologia*, 49, 2627–33.
- Juarez, D. T., Sentell, T., Tokumaru, S., Goo, R., Davis, J. W., & Marjorie, M. (2012). Factors Associated With Poor Glycemic Control or Wide Glycemic Variability Among Diabetes Patients in Hawaii , 2006 – 2009, 9, 1–10.
- Kementrian Kesehatan RI. (2009). Tahun 2030 prevalensi diabetes melitus di Indonesia mencapai 21,3 juta orang. Retrieved May 10, 2015, from www.depkes.go.id
- Khattab, M., Khader, Y. S., Al-khawaldeh, A., & Ajlouni, K. (2010). Factors associated with poor glycemic control among patients with Type 2 diabetes. *Journal of Diabetes and Its Complications*, 24(2), 84–89.
<http://doi.org/10.1016/j.jdiacomp.2008.12.008>
- Kisokanth, G., Prathapan, S., Indrakumar, J., & Joseph, J. (2013). Factors influencing self management of diabetes mellitus; a review article. *Journal of Diabetology*.
- Krabbe, K.S., Nielsen, A.R., Krogh-Madsen, R., Plomgard, P., Rasmussen, P., Erikstrup, C., Fischer, C.P., Lindegaard, B., Petersen, A.M.W., Taudorf, S., Secher, N.H., Pilegaard, H., Bruunsgaard, H., Pedersen, B.K., 2007. Brain-derived neurotrophic factor (BDNF) and type 2 diabetes. *Diabetologia* 50, 431—438.
- Lee, S., Kim, Y., Lee, S., Cho, Y., Choi, Y., Chun, C., & Chang, Y. (2004). Effect of diabetes education on diabetic management in non insulin dependent diabetics mellitus patients. *Journal of Thr Korean Dietetic Association*, 10, 300–08.
- Lubis, N.L. (2009). Depresi; Tinjauan psikologis. Kencana Jakarta:
- Luppino, E.S., de Wit, L.M., Bouvy, P.F., Stijnen, T., Cuijpers, P., Penninx, B.W., Zitman, F.G. (2010). Overweight, obesity, and depression: a systematic review and meta-analysis of longitudinal studies. *Arch. Gen. Psychiatry* 67: 220-9
- Lustman, P.J., Anderson, R.J., Freedland, K.E., De Groot, M., Carney, R., Clouse, R. E. (2000). Depression and poor glycemic control: a meta-analytic review of the literature. *Diabetes Care*, 23(7), 934–42.
- Lustman, P.J., Clouse, R.E., Freedland, K. E. 1998. M. of major depression in adult with diabetes: implication of recent clinical trials. *S. clinical neuropsychiatry* 3: 102-114. (1998). Management of major depression in adult with diabetes: implication of recent clinical trials. *Semin Clinical Neuropsychiatry*, 3, 102–114.
- Lustman, P.J., Griffith, L.S., Clouse, R.E., Freedland, K.F., Eise, R.A., Rubin, E.H., et al. 1997. (1997). Effect of nortriptyline on depression and glycemic control in

diabetes: result of a double-blind , placebo-controlled trial. *Psychosom Med*, 59, 241–250.

Madiyono, B., Moeslichan, S., Sastroasmoro, S., Budiman, I., & Purwanto, S. H. (2014). Perkiraan besar sampel dalam *Dasar-dasar metodologi penelitian klinis*.

Maslim, R. (2004). *Buku Saku Diagnosis Gangguan Jiwa (PPDGJ-III)*. FK Jiwa Unika Atmajaya. Jakarta:

Maurer, D.M, Darnall C.R. 2012. Screening for depression. *Am Fam Physician* 85(2):139-44. Texas

McWhinney, I. R. (1997). *A textbook of family medicine*. Oxford university press. New York.

Monnier, L., & Colette, C. (2009). Target for glycemic control. *Diabetes Care*, 32(2), 5199–204.

Mosby's medical dictionary. (2009). Retrieved from <Http://medical-dictionary.thefreedictionary.com/family+function>

Musselman, D.L., Betan, E., Larsen, H., Phillips, L. S. (2003). Relationship of depression to diabetes type 1 dan 2: epidemiology, biology, and treatment. *Biol. Psykaiatry*, 54(3), 317–79.

Nash,Jen. (2014). Understanding the psychological impact ofdiabetes and the roe of clinical psychology. *Journal of Diabetes Nursing* 18: 137-42.

Oliveira, S.C., Santos, A.A., Pavarini,S.C.I. (2014). The relationship between depressive symptoms and family functioning in institutionalized elderly. *Rev Esc Enferm USP* 48(1): 65-71

Perkeni. (2011). pedoman pengendalian dm n penykronis.depkes.pdf.

Rohlfing, C. L., Wiedmeyer, H., & Little, R. R. D. C. (2002). Defining the relationship between plasma glucose and HbA1c : analysis of glucose profile and HbA1c in the diabetes control and complication trial. *Diabetes Care*, 25(2) : 275-78.

Santos, A. A., & Pavarini, S. C. (2012). Family functionality regarding the elderly with cognitive impairments: the caretaker's perception. . *Rev Esc Enferm USP*, 46(5), 1140–6.

Sastroasmoro, S. (2014). Pemilihan subjek penelitian. In *Dasar-dasar metodologi penelitian klinis*.

- Shepard, L.A. (1979). Self-acceptance: the evaluative component of the self-concept construct. *American Educational Research Journal of Health Psychology*. Vol 6(6) pp.651-663
- Shortridge-baggett, L. M. (2004). Structured evaluation method for family assessment tools : evaluation of the family apgar. *Fam. Saúde Desenv.*, 6(2), 91–99.
- Smilkstein, G. (1978). Family APGAR : a proposal for a family function test and its use by physicians. *The Journal of Family Practice*, 6(6), 1231–9.
- Stuart, M.J. , Baune, B.T. (2012). Depression and type 2 diabetes: Inflammatory mechanisms of a psychoneuroendocrine co-morbidity. *Neuroscience and Biobehavioral Reviews* 36: 658-76.
- Sutikno, E. (2011). Hubungan antara Fungsi Keluarga dan Kualitas Hidup Lansia. *Jurnal Kedokteran Indonesia*, 2(1), 73–79.
- Takenaka, H., Sato, J., Suzuku, T., Ban, N. (2013). Family issues and family functioning of Japanese outpatients with type 2 diabetes: a cross-sectional study. *BioPsychoSocial Medicine* 7: 13, 7(13).
- Vaciana, M., Major, C., & Morgan, M. (1995). Postprandial versus preprandial blood glucose monitoring in women with gestational diabetes mellitus requiring insulin therapy. *New England J Med*, 333, 1237–41.
- Whitehead, A. L., Dimmock, M., & Place, M. (2013). Diabetes Control and The Influence of Family Functioning,. *Journal of Diabetes Research and Clinical Metabolism*, 2(16).
- Widoyoko, E.P. (2012). *Teknik penyusunan instrumen penelitian*. Pustaka Pelajar. Yogyakarta.
- Wild, S., Roglic, G., Green, A., Sicree, R., & King, H. (2004). Estimates for the year 2000 and projections for 2030. *Diabetes Care*, 27(5), 1047–53.
- Wonca. (2002). The European definition of general practice/family medicine.
- Yun, J., & Kim, K. (2009). Relationships of family support , diet therapy practice and blood glucose control in type II diabetic patients *, 3, 141–148.
<http://doi.org/10.4162/nrp.2009.3.2.141>.
- Zahid, N., Asghar, S., Claussen, B., & Hussain, A. (2008). Depression and diabetes in a rural community in Pakistan. *Diabetes Research and Clinical Practice*.
- Zigmond, A. S., & Snaith, R. P. (1983). The Hospital Anxiety and Depression Scale. *Acta Psykaiatr Scand*, 67, 361–70.