

REFERENCES

- Amagata, T., Minoura, K., and Numata, A., 1998, Cytotoxic Metabolites Produced by a Fungal Strain from a *Sargassum* Alga, *J. Antibiot.*
- Anonim, 2016, Global Health Observatory (GHO) data, World Health Organization (WHO), www.who.int
- Ariefta, Nanang Rudianto, 2015, Screening of Novel secondary Metabolites from Endophytic Fungi of Tropical Forest, *Thesis*, Department of Chemistry, Gadjah Mada University.
- Bills, G. F., Platas, G., Fillola A., Jimenez M.R., Collado, J., Vicente, F., Martin, J., Gonxzalez, A., Bur-Zimmermann, J., Tormo, J.R.n and Pelaez F., 2008, Enhancement of antibiotic and secondary metabolite detection from filamentous fungi by growth on nutritional arrays, *J. App. Microbiol.*, 104, 1644-1658.
- Bode, Helge Bjorn, Bethe, B., Hofs, Regina, and Zeeck, A., 2002, Big Effects from Small Changes: Possible Ways to Explore Nature's Chemical Diversity, *Chem. Bio. Chem.*, 3, 619-627.
- Chokpaiboon, Supichar, Sommit, D., Teerawatananond, T., Muangsing, N., Bunyapaiboonsri, and Pudhom, K., 2010, Cytotoxic Nor-chamigrane and Chamigrane Endoperoxides from a Basidiomycetous Fungus, *J. Nat. Prod.*, 73, 1005-1007.
- Ciavatta, Maria L., Lopez-Gresa, M. P., Gavagnin, M., Nicoletti, R., Manzo, E., Mollo, E., Guo, Y., and Cimino G., 2008, Cytosporin-related compounds from the marine-derived fungus *Eutypella scoparia*, *Tetrahedron*, 64, 5365-5369.
- Cole, Richard J. and Kirksey, Jerry W., 1974, A new tremorgenic metabolite from *Penicillium paxilli*. *Can. J. Microbiol.* Vol. 20
- Connolly, J. D. and Hill, R. A., 1991, Dictionary of Terpenoids Volume 1 Mono- and Sesquiterpenoids, UK: University of Glasgow.
- Cragg, Gordon M., Newman, David J., and Snader, Kenneth M., 1997, Natural Products in Drug Discovery and Development, *J. Nat. Prod.*, Vol. 60, No. 1.
- Deshmukh, Sunil Kumar, Prabhu, Mishra Dutt, Kulkarni-Almeida, A., Verekar, S., Sahoo, M. S., Periyasamy, G., Goswami, H., Khanna, A., Balakrishman A.,

and Vishwakarma, R., 2009, Anti-Inflammatory and Anticancer Activity of Ergoflavin Isolated from an Endophytic Fungus, *Chem. Biodiver.*, Vol 6.

- Dixon, Richard A., 2001, Natural products and plant disease resistance, *Nature*, Vol 411.
- Fallon, A., Booth, R.F.G., and Bell, L.D., 1987, Laboratory Techniques in Biochemistry and Molecular Biology Volume 17 *applications of HPLC in biochemistry*. New York: Elsevier.
- Firodiya, Anand Dilip and Tenguria, Rajesh Kumar, 2016, Endophytic Fungi: Treasure for Anti-cancerous Compounds, *Int. J. Pharm. Sci.*, Vol 8, Issue 8, 35-42.
- Gubiani, Juliana R., Zeraik, Maria L., Oliveira, Camila M., Ximenes, Valdecir F., Nogueira, Cludio R., Fonseca, Luiz M., Silva, Dulce H. S., Bolzani, Vonderlan S., Araujo, angela R., 2014, Biologically Active Eremophilane-Type Sesquiterpenes from *Camaropsi* sp., an Endophytic Fungus Isolated from *Albertia macrophylla*, *J. Nat. Prod.*, 77, 668-672.
- Gunatilaka, A. A. Leslie, 2006, Natural Products from Plant-Associated Microorganisms: Distribution, Structural Diversity, Bioactivity, and Implications of Their Occurrence, *J. Nat.* 69, 509-526.
- Gutierrez, R. M. P., Gonzalez A. M. N., and Ramirez, A. M., 2012, Compounds Derived from Endophytes: A Review of Phytochemistry and Pharmacology, *Curr. Med. Chem.*, 19, 2992-3030.
- Hajnos, Monika Waksmundzka., Sherma, Joseph., and Kowalska, Teresa, 2008, Thin Layer Chromatography in Phytochemical, *Chromatography Science Series* Volume 99, New York: CRC Press.
- Hajnos, Monika Waksmundzka and Sherma, Joseph., 2011, High Performance Liquid Chromatography in Phytochemical Analysis, *Chromatography Science Series*, New York: CRC Press.
- Hashida, J., Niitsuma, M., Iwatsuku, M., Mori, M., Ishiyama, A., Namatame, M., Nishihara-Tsukashima, A., Nonaka, K., Ui, H., Masuma, R., Otaguro, K., Yamada, H., Shiomi, K., and Omura, S., 2010, Pyrenocine I, a new pyrenocine analog produced by *Paecilomyces* sp. FKI-3573, *J. Antibiot.*, 53, 559-561.
- Hosoe, Tomoo., Nozawa, Koohei., Udagawa, Shun-ichi., Nakajima, Shoichi., and Kawai, Ken-ichi., 1990, Structure of New Indoloditerpenes, Possible Biosynthetic Precursors of The Tremorgenic Mycotoxins, Penitrem, from *Penicillium crustosum*. *Chem. Pharm. Bull.* 38 (2), 3473- 3475.

- Huang, Yong Fu, Qiao, L., Lv, A. Li, Pei, Yue Hu, and Tian, Li, 2008, Eremophilane sesquiterpenes from the marine fungus *Penicillium* sp. BL27-2, *Chin. Chem. Lett.*, 19, 562-564.
- Hyeon, Suong-Be., Ozaki, Akio., Suzuki, Akinori, and Tamura, Saburo., 1976, Isolation of $\alpha\beta$ -Dehydrocurvularin and β -Hydrocurvularin from *Alternaria tomato* as Sporulation-suppressing Factors, *Agr. Biol. Chem.*, 40 (8), 1663-1664.
- Isaka, Masahiko, Chinthanom, P., Boonruangprapa, T., Rungjindamai N., Pinruan, U., 2010, Eremophilane-Type Sesquiterpenes from the Fungus *Xylaria* sp. BCC 21097, *J. Nat.*, 73, 683-687.
- Joseph, Baby and Priya R. Mini, 2011, Bioactive Compounds from Endophytes and their Potential in Pharmaceutical Effect: A Review, *Am.J. Biochem. Mol. Biol.*, 1 (3): 291-309.
- Kobayashi, A., Hino, T., Yata, S., Itoh, Tomohiko, J., Sato, H., and Kawazu, K., 1988, Unique Spindle Poisons, Curvularin and Its Derivatives, Isolated from *Penicillium* Species, *Agric. Biol. Chem.*, 52 (12), 3119-3123.
- Kusano, M., Nakagami, K., Fujioka, S., Kawano, T., Shimada, A., and Kimura, Y., 2003, $\beta\gamma$ -Dehydrocurvularin and Related Compounds Nematicides of *Pratylenchus penetrans* from the Fungus *Aspergillus* sp., *Biosci. Biotechnol. Biochem.*, 67 (6), 1413-1416.
- Liu, Ling, Liu, S., Chen, X., Guo, L., 2009, Pestalofones A-E, Bioactive Cyclohexanone Derivatives from The Plant Endophytic Fungus *Pestalotiopsis fici*, *Bioorg. Med. Chem.*, 17, 606-613
- Lodge, D. Jean, Fisher, P.J., and Sutton, B. C., 1996, Endophytic fungi of *Manilkara bidentata* leaves in Puerto Rico, *Mycologia*, 88, 733-738.
- Motohashii, Keiichiro, et al., 2009, New sesquiterpenes, JBIR-27 and -28, isolated from a tunicate-derived fungus, *Penicillium* sp. SS080624SCf1, *The Journal of Antibiotics*, 62, 247-250.
- Newman, David J. and Cragg, Gordon M., 2007, Natural Products as Sources of New Drugs over the Last 25 Years, *J. Nat. Prod.*, 70, 461-477.
- Newman, David J. and Cragg, Gordon M. 2012, Natural Products As Sources of New Drugs over the 30 Years from 1981 to 2010, *J. Nat. Prod.*, 75, 311-335.

- Nozawa, K., Horie, Y., Udagawa, Shun-ichi., Kawai, Ken-ichi., and Yamazaki, Mikio., 1989, Isolation of a New Tremorgenic Indoloditerpene, 1'-O-Acetylpaxilline, from *Emericella striata* and Distribution of Paxilline in *Emericella* spp, *Chem. Pharm. Bull.*, 37 (5), 1387-1389.
- Pavia, Donald L., Lampman, Gary M., Kriz G.S., Vyvyan, J.R., 2001, Introduction to Spectroscopy Fourth Edition, *Book/ Cole*, USA.
- Pelaez, Fernando, 2006, The historical delivery of antibiotics from microbial natural products-Can you repeat?, *Biochemic. Pharmacol.*, 71. 981-990.
- Riss, Terry L., Moravec, Richard A., Niles, Andrew L., Duellman, S., Benink, Helene A., Worzella, Tracy J., Minor, L., 2016, Assay Guidance Manual Cell Viability Assays, National Library of Medicine, National Center for Advancing Translational Sciences.
- Schulz, B., Wanke U., Draeger S., Aust H-J., 1993, Endophytes from herbaceous plants and shrubs: effectiveness of surface sterilization methods, *Mycol. Res.*, 97 (12), 1447-1450.
- Siriwach, Ratklao, 2013, Screening of Novel Secondary Metabolites from Endophytic Fungi by Chemical Library Analysis, *Dissertation*, Department of Biotechnology, Osaka University.
- Shiono, Y., Hayasaka S., Murayama T., Koseki T., Uesugi S., Kimura K., 2012, A polyketide metabolite from the fungicolous *Nodulisporium* sp. SH-1, *Phytochem. Lett.*, 5, 549-552.
- Silva, Geraldo H., de Oliveira, Camila M., Teles, Helder L., Pauletti, Patricia M., Castro-Gamboa, I., Silva, Dulce H.S., Bolzani, Vanderlan S., Young, Maria C.M., Costa-Neto, Claudio M., Pfenning, Ludwig H., Berlinck, Roberto G.S., Araujo, Angela R., 2010, Sesquiterpenes from *Xylaria* sp., an endophytic fungus associated with *Piper aduncum* (Piperaceae), *Phytochem. Lett.*, 3, 164-167.
- Souza, Jucimar Jorgeane de, Vieira I.J.C., Rodrigues-Filho, E. and Braz-Filho, R., 2011, Terpenoid from endophytic Fungi, *Molecules*, 16, 10604-10618.
- Strobel, Gary and Daisy, Bryn, 2003, Bioprospecting for Microbial Endophytes and Their Natural Products, *Microbiol. Mol. Biol. Rev.*, 67(4): 491.
- Strobel, Gary, Daisy, B., Castillo, U., and Harper J., 2004, Natural Products from Endophytic Microorganisms, *J. Nat. Prod.*, 67, 257-268.

- Suryanarayanan, Trichur S. and Kumaresan, Vadivelu, 2001, Occurrence and distribution of endophytic fungi in a mangrove community, *Mycol. Res.* 105 (11): 1388-1391.
- Suryanarayanan, T. S., Murali, T. S., and Vankatesan G., 2002, Occurrence and distribution of fungal endophytes in tropical forests across a rainfall gradient, *Can. J. Bot.* 80: 818-826.
- Takagi, Ichiro., Tazuke, Yashuhito., and Naya, Keizo., 1977, The Components of *Petasites japonicas* Maxim. IV. The Structure of Petasitin, *Bull. Chem. Soc. Japan*, Vol. 50 (12), 3320-3323.
- Tan, R. X. and Zou W. X., 2001, Endophytes: a rich source of functional metabolites, *Nat. Prod. Rep.*, 18, 448-459.
- Tanaka, Seiichi., Mada, Kojiro., Morumo, Shingo., and Hattori, Hiroyuki, 1984, Structure of Sporogen-AO 1, A Sporogen Substance of *Aspergillus oryzae*, *Tetrahedron Lett.*, Vol. 25, 51, 5907-5910.
- Tejesvi, Mysore V., Mahesh. B., Nalini, M.S., Prakash, H.S., Kini, K.R., Subbiah V., Shetty, H.S., 2006, Fungal endophyte assemblages from ethnopharmacologically important medicinal trees, *Can. J. Microbiol.* 52: 427-435.
- Tejesvi, Mysore V., Kini, K.R., Prahash, H.S., Subbiah, V., Shetty, H.S., 2008, Antioxidant, antihypertensive, and antibacterial properties of endophytic *Pestalotiopsis* species from medicinal plants, *Can. J. Microbiol.* 54: 769-780.
- Tuntiwachwuttikul, Pittaya, Taechowisan, T., Wanbanjob, A., Thadaniti, S., Taylor, W. C., 2008, Lansai A-D, secondary metabolites from *Streptomyces* sp. SUC1, *Tetrahedron*, 64, 7583-7586.
- VanderMolen, Karen M., Raja, Huzefa A., El-Elimat, T., and Oberlies, Nicholas H., 2013, Evaluation of culture media for the production of secondary metabolites in a natural products screening program, *AMB Express*, 3: 71.
- Verma, Vijay C., Gond, S.K., Kumar, A., Kharwar, R.N., Boulanger, L., Strobel G. A., 2011, Endophytic Fungal Flora from Root and Fruits of an Indian Neem Plant *Azadirachta indica* A. Juss., and Impact of Culture Media on Their Isolation, *Indian. J. Microbiol.*, 51 (4): 469-476.
- Wani, M. C., Taylor, H.L., Wall, Monroe E., Coggon, P., McPhail, A.T., 1971, Plant Antitumor Agents. VI. The Isolation and Structure of Taxol, a Novel Antileukemic and Antitumor Agent from *Taxus breoifolia*, *J. Am. Chem. Soc.*, 93:9.

- Weber, Holly A., Baenziger, Norman C., and Gloer, James B., 1990, The Preussomerins: Novel Antifungal Metabolite from the Coprophilous Fungus *Preussia isomera* Cain, *J. Org. Chem.*, 56, 4355-4360.
- Weber, Holly A. and Gloer, James B., 1991, Structure of Preussomerin A: An Unusual New Antifungal Metabolite from the Coprophilous Fungus *Preussia isomera*, *J. Am.Chem. Soc*, Vol 112, No. 18.
- Xu, Ren, Wang, M., Lu, C., Zheng, Z., Shen, Y., 2009, Tuberculariols A – C, New Sesquiterpenes from the Mutant Strain M-741 of *Tubercularia* sp. TF 5, *Helvetica Chimica Acta*, Vol 92.
- Yu, Hongsheng, Zhang, L., Li, L., Zheng, C., Guo, Lei., Li, W., Sun, P., Qin, L., 2009, Recent developments and future prospects of antimicrobial metabolites produced by endophytes, *Microbiol. Res.*, 165, 437-449.
- Yurchenko, A. N., Smetanina, O.F., Kalinovskii, A.I., Kirichuk, N.N, Yurchenko, E.A., Afiyatullo, Sh., Sh., 2013. Biologically Active Metabolites of the Facultative Marine Fungus *Penicillium citrinum*, *Chem. Nat. Compd.*, Vol. 48, No. 6.