

DAFTAR PUSTAKA

- Adam, J.M., Soegondo, S., Semiardji, G., Adriansyah, H., 2006. *Petunjuk Praktis Penatalaksanaan Dislipidemia*. PB.PERKENI; Jakarta: 1-22.
- Ahmad, J., Ahmed, F., Siddiqui, M.A., Khan, A.R., Katyal, P., Hameed, B., 2007, Inflammatory Markers, Insulin Resistance and Carotid Intima-Media Thickness in North-Indian Type 2 Diabetic Subjects. *JAPI*; 55 : 693-700.
- Ahuja, A., Evans, R., 2000. *Carotid and Vertebral Ultrasonography. Practical head & Neck ultrasound*. Greenwich. Medical Media. London: 148-150.
- American Diabetes Association (ADA), 2006, American Diabetes Association: Clinical Practise Recommendations 2006. *Diabetes Care*: 29.
- American Diabetes Association (ADA). 2010. Position statement: Standar of medical care in diabetes 2010. *Diabetes Care*; 33: 1-47.
- Anonymous, 2010. *Cobas C system*. Roche; English V; 6:3.
- Antonio G.C, Luis E.S.M, Sandra E.A.2011. Elevated Triglycerides/HDL-cholesterol ratio associated with insulin resistnace. *Cir Cir*; 79:115-119.
- Bahar, Iriani., Murtala, Bachtiar., Ilyas, Muhammad., Liyadi, Frans., Aman Andi, Makbul., Bahar, Burhanuddin., 2013, Hubungan Ketebalan Intima Media Arteri Karotis Berdasarkan Pemeriksaan Ultrasonografi Dengan Fraksi Lipid Darah Penderita Dislipidemia. *Indonesian Journal of Clinical Pathology And Medical Laboratory*; 20:1-10.
- Basta, G., Schmidt, A.M., De Caterina, R., 2004. Advanced glycation end products and vascular inflammation: implications for accelerated atherosclerosis in diabetes. *Cardiovasc Res*; 63 : 582-592.
- Beckman, J.A., Creager, M.A., Libby, P., 2002, Diabetes and Atherosclerosis Epidemiology, Pathophysiology, and Management, *JAMA*; 287:2570-81.
- Belhassen, L., Carville, C., Pelle, G., *et al.* 2002. Evaluation of Carotid Artery and Aortic IntimaMedia Thickness Measurements for Exclusion of Signi cant Coronary Atherosclerosis in Patients Scheduled for Heart Valve Surgery. *J Am Coll Cardiol*; 39:1139-44.
- Bots, Michiel L., Hoes, Arno W., Peter J, Koudstaal., Albert, Hofman., Grobbee, Diederick E., 1997. *Common Carotid Intima-Media Thickness and Risk of Stroke and Myocardial Infarction*. American Heart Association, Inc, 96:1432-1437.
- Bravo, Mónica., Collado, Laura., Dardanelli, Esteban., Araujo, María Beatriz., Lipsich, José., Moguillansky, Silvia., 2012. Ultrasonography measurement of carotid intimamedia thickness in pediatric patients with obesity, familial hypercholesterolemia and type 1 diabetes. *RAR*; 76(1):1-7.
- Brown, CT, 1995. Penyakit aterosklerotik koroner. Dalam: Price SA, Wilson LM, editor. *Patofisiologi Konsep Klinis Proses-Proses Penyakit*. Edisi 4. volume 2. Jakarta: EGC.

- Burtis, C.A., Ashwood, E.R., Bruns, D.E., 2008. *Tietz Fundamental of Clinical Chemistry*. 6th ed. Saunders, Elsevier, Missouri.:201-228.
- Chan, Y.H., 2003, Biostatistics 101: Data Presentation, *Singapore Med. J*; 44(6): 280-285.
- Chodakauskas, T., Feinstein, S., 2006. Non-invasive measurement of surrogate markers of atherosclerosis: Carotid intima-media thickness (c-IMT). *Vivid.Ultrasound*;7:1.
- Da luz, P,L., Cesena, F,H., Favarato, D., Cerqueira, E,S., 2005. Comparison of serum lipid values in patients with coronary artery disease at <50, 50 to 59, 60 to 69, and >70 years of age. *Am J Cardiol*;96:1640-3.
- Dahlan, M.S., 2009. *Statistik untuk Kedokteran dan Kesehatan: Deskriptif, Bivariat, dan Multivariat Dilengkapi Aplikasi dengan Menggunakan SPSS*. Penerbit Salemba Medika, Jakarta.
- Dahlen, E.M., Lanne, T., Engvall, J., Lindstrom, T., Grodzinsky, E., Nystrom, F.H., et al., 2009. Carotid intima-media thickness and apolipoprotein B/apolipoprotein A-I ratio in middle-aged patients with Type 2 diabetes, Journal compilation. *Diabetic medicine*, 26: 834 – 390.
- De Giorgis., Tommaso., Marcovecchio, M., Loredana., Di Giovanni, Ilaria., Giannini, Cosimo., Chiavaroli, Valentina., Chiarelli, Francesco., Mohn, Angelika., 2014, Triglycerides-to-HDL ratio as a new marker of endothelial dysfunction in obese prepubertal children. *Europ J Endocrin*: 170, 173–180.
- Departemen Kesehatan Republik Indonesia (Depkes RI), 2005, *Pharmaceutical Care Untuk Penyakit Diabetes Melitus*, Direktorat Bina Farmasi Komunitas dan Klinik, Jakarta.
- Departemen Kesehatan Republik Indonesia (Depkes RI), 2007. Riset Kesehatan Dasar (RISKESDA): *Laporan Nasional tahun 2007 Badan Penelitian dan Pengembangan Departemen Kesehatan*. Depkes RI. Jakarta.
- Departemen Kesehatan Republik Indonesia (Depkes RI), 2009. *Profil Kesehatan Indonesia 2008*. Jakarta.
- Dharmalingam., Mala., Deshpande., Neeta, R., Vidyasagar., Sudha. 2004. Triglyceride Levels And Its Correlation With Carotid Intima-Media Thickness. *INT. J. Diab. Dev. Countries*; 1 24:19-24.
- Dinas Kesehatan Daerah Istimewa Yogyakarta (Dinkes DIY), 2013. *Laporan Surveilans Terpadu Penyakit (STP) Yogyakarta tahun 2012*. Yogyakarta.
- Djokomoeljanto, R., 2000, *Mengelola diabetes dalam rangka mencegah komplikasi kardiovaskuler*, Simposium Diabetes, Semarang.
- Dobiasova., M, Frohlich., J. 2001. The plasma parameter log(TG/HDL-C) as an atherogenic index: correlation with lipoprotein particle size and esterification rate in apoB-lipoprotein-depleted plasma (FERHDL). *Clinical Biochemistry*; 34:583-588.

- Fan, A.Z., Labrador, M.P., Bairey, C.N., 2006. Smoking status and common carotid artery intima-medial thickness among middle-aged men and women based on ultrasound measurement: a cohort study. *BMC*; 6:42.
- Frances, F., 2002. A Manual of Laboratory And Diagnostic Test. 6th ed. Philadelphia:Lippincott,467-75.
- Gaziano, J.M., Hennekens, C.H., O'Donnell, C.J., Breslow, J.L., Buring, J.E., 1997. Fasting triglycerides, high density lipoprotein, and risk of myocardial infarction. *Circulation*. 96:2520-5.
- Giannini, C., Santoro, N., Caprio, S., Kim, G., Lartaud, D., Shaw, M., 2011. The triglyceride-to-HDL cholesterol ratio: association with insulin resistance in obese youths of different ethnic backgrounds. *Diabetes Care*; 34: 1869-1874.
- Golemati, Tegos., 2003. Echogenicity of B-Mode Sonographic Image of the Carotid Artery. *J Ultrasound Med*; 23(5):659-69.
- Gosai, Falgun., Patel, Hetal., 2014. Relationship Between Carotid Intima Media Thickness And Lipid Profile In Type Two Diabetes Mellitus And Stage 2 Hypertension. *NJIRM*; 5(2) :39-45
- Gotthelf, S.J., 2014. TG/HDL Index in adolescents of Salta, *Argentina Journal*; 43(1):18-24.
- Hofer, M., 2004. *Cerebrovascular Imaging. Teaching Manual of Color Duplex Sonography*. Second edition. Theme; 28-30.
- Hu, Y.M., Tian, H.M., Liu, R., Chen, X., 2004. Atherogenic index of plasma is associated with carotid intima-media thickness in patients with type 2 diabetes mellitus. *Journal of Sichuan University*; 35(5):696-8, 707.
- International Diabetes Federation (IDF), 2005, *IDF Clinical Guidelines Task Force. Global guideline for Type 2 diabetes*. Brussels.
- International Diabetes Federation (IDF), 2013, *IDF Clinical Guidelines Task Force. Global guideline for Type 2 diabetes*. Brussels.
- Jameson, J., Larry., 2006. *Harrison's Endocrinology*. 16th Edition. The McGraw-Hill Companies, Inc. United States of America; p.283-293.
- Jandeleit, D.K., Cooper, M.E., 2008. The role of AGEs in cardiovascular disease. *Curr Pharm*; 14:979-986.
- Jegelevicius, D., Lukosevicius, 2002. An Ultrasonic Measurement of Human Carotid Artery Wall Intima Media Thickness. *Ultragarsas*; 2:43.
- Jeppesen, J., Hein, H.O., Suadicani, P., Gyntelberg, F.. 1998. Triglycerides concentration and ischemic heart disease: an eight-year follow-up in the Copenhagen Male Study. *Circulation*. 97:1029-36.
- Jungner, I., Marcovina, S.M., Walldius, G., Holme, I., Kolar, W., Steiner, E., 1998. Apolipoprotein B and A-I values in 147 576 Swedish males and females, standardized according to the World Health Organization-International Federation of Clinical Chemistry First International Reference Materials. *Clin Chem* 44: 1641-9.

- Juonala, M., Ka, H.N.M., Laitine, T., Jokinen, E., Taittonen, L., 2008. Effect of age and sex on carotid intima-media thickness, elasticity and brachial endothelial function in healthy adults: The Cardiovascular Risk in Young Finns Study. *Euro Heart J*; 29 : 1198–1206.
- Kader., Mohammad, A., 2015. Correlation Between Triglyceride/HDL-Cholesterol Ratio and The Severity of Atherosclerotic Lesions. Cardiology Department, El-Menia University. The Egyptian Journal of Hypertension And Cardiovascular Risk; 12 (1):1687-5338.
- Kaptoge, S., White, I.R., Thompson, S.G., Wood, A.M., 2007. Fibrinogen Studies Collaboration:Associations of plasma fibrinogen levels with established cardiovascular disease risk factors, inflammatory markers, and other characteristics: individual participant meta-analysis of 154,211 adults in 31 prospective studies: the fibrinogen studies collaboration. *Am J Epidemiol*; 166: 867–879.
- Kasami, R., Kaneto, H., Katakami, N., *et al.* 2011. Relationship Between Carotid Intima-Media Thickness and the Presence and Extent of Coronary Stenosis in Type 2 Diabetic Patients With Carotid Atherosclerosis but Without History of Coronary Artery Disease. *Diabetes Care*, 34:468-70.
- Khan, Sohail, Rafi., Ayub, Nishat., Nawab, Sohail., Shamsi, Tahir., 2008. Triglyceride Profile in Dyslipidaemia of Type 2 Diabetes Mellitus. *Journal of the College of Physicians and Surgeons*;18 (5): 270-273.
- Kementerian Kesehatan Republik Indonesia (Kemenkes RI). 2014. *Situasi dan Analisis Diabetes. Pusat Data dan Informasi.* Jakarta.
- Kerwin, W.S., 2007. Correlation of Carotid Artery Pathology and Morphology in imaging. *Springer Vienna.J*; 19:1
- Keulen, E.T.P., Kruijshoop, M., Schaper, N.C., Hoeks, A.P.G., de Bruin, T.W.A., 2002. Increased Intima-Media Thickness in Familial Combined Hyperlipidemia Associated With Apolipoprotein B. *Arterioscler Thromb Vasc Biol*; 22: 283-8.
- Kwiterovich, J.P.O., 2000. The metabolic pathways of high-density lipoprotein, low density lipoprotein, and triglycerides: A current review. *Am J Cardiol*; 86: 5-10.
- Lee, E.J., Kim, H.J., Bae, J.M., Kim, J.C., Han, H.J., Park, C.S., 2007. Relevance of Common Carotid Intima-Media Thickness and Carotid Plaque as Risk Factors for Ischemic Stroke in Patients with Type 2 Diabetes Mellitus. *AJNR*; 28: 916-9.
- Lina Y., 2007. High Sensitivity C-Reactive Protein (hsCRP). *Forum Diagnosticum*; 3:1-11.
- Magyar, M.T., Szikszai, Z., Balla, J., Valikovics, A., Kappelmayer, J., Imre, J., 2002, Early-onset carotid atherosclerosis is associated with increased intima-media thickness and elevated serum levels of inflammatory markers. *Stroke*; 34:58-63.

- Maiti, R., Agrawal, N.K., 2007. Atherosclerosis in diabetes mellitus: Role of inflammation. *Atherosclerosis in diabetes mellitus. Indian J.Med*; 61:292-306.
- Marieb., Elaine, N., Hoehn., Katja., 2007, *Human Anatomy and Physiology*, 7th edition [e-book]. San Francisco: Benjamin Cummings; Chap.19.
- Matsumoto, S., Gotoh, N., Hishinuma, S., Abe, Y., Shimizu, Y., Katano, Y., 2014. The Role of Hypertriglyceridemia in the Development of Atherosclerosis and Endothelial Dysfunction. *Nutrients*; 6: 1236-1250.
- Mayes, P.A., 1997. Pengangkutan dan Penyimpanan Lipid. Dalam: Murray R.K., Granner D.K., Mayes P.A., Rodwell V.W., ed: *Biokimia Harper*. Edisi 24. Penerbit EGC, Jakarta.
- Mohan, V., Ravikumar, S., Rani, S., Deepa, R., 2000. Intimal medial thickness of the carotid artery in South Indian diabetic and non-diabetic subjects: the Chennai Urban Population Study (CUPS). *Diabetologia*; 43: 494-9.
- Mooradian, A.D., 2008. Dyslipidemia in type 2 diabetes mellitus. *Nature Clinical Pract Endocrinol & Metab*; 5(3): 150 – 57.
- Packard, C.J., Shepherd, J., 1999. Lipoprotein heterogeneity and apolipoprotein B metabolism. *Arterioscler Thromb Vasc Biol*;19:2456–64.
- Perkumpulan Endrokinologi Indonesia (PERKENI). 2011. *Konsesus pengelolaan dan pencegahan diabetes mellitus tipe 2 di Indonesia*. PB PERKENI; Jakarta.
- Pischon, T., Girman ,C.J., Sacks, F.M., Rifai, N., Stampfer, M.J., Rimm, E.B., 2005. Non-HDL Cholesterol and Apolipoprotein B in the Prediction of Coronary Heart Disease in Men. *Circulation*; 112:3375-83.
- Ridker, P.M., Stampfer, M.J., Rifai, N., 2001. Novel risk factors for systemic atherosclerosis: a comparison of C-reactive protein, fibrinogen, homocysteine, lipoprotein(a), and standard cholesterol screening as predictors of peripheral arterial disease. *JAMA*; 285 : 2481-5.
- Rifai, N., Warnick, G.R., Remaley, A.T., 2008. Analytes : Lipid, Lipoproteins, Apolipoproteins, and Other Cardiovascular Risk Factors. In : Burtis CA, Ashwood ER, Brunis DE (editors) *Fundamentals of Clinical Chemistry*. Sixth Edition. Philadelphia : Saunders Elsevier:428.
- Rizzo, M., Corrado, E., Coppola, G., Muratori, I., Novo, G., Novo, S., 2009. Markers of inflammation are strong predictors of subclinical and clinical atherosclerosis in women with hypertension. *Coron Artery Dis*; 20: 15-20.
- Rochmah, W., 2009. Diabetes Mellitus Pada Usia Lanjut. In: Soedoyo, A.W., Setiyohadi, B., Alwi, I., Simadibrata, M.,Setiati, S. (Ed.): *Buku Ajar Ilmu Penyakit Dalam*, pp:1967-72. Interna Publishing, Jakarta.
- Rothwell., P.M., 2001. The Interrelation between carotid, femoral and coronary artery disease. *Euro Heart J*; 22: 11-14.
- Powers, A.C., 2008. Diabetes Mellitus. In: Fauci AS, Braunwald E, Kasper DL, editors. *Harrison's Principles of Internal Medicine*. 17th ed. McGraw-Hill Companies, Inc. : 2275-2304.

- Protasio L.L.I., Desiderio, F.I., Jose, R.F., Neto, J., Pedro, L., Antonio C.C., 2008. High Ratio of Triglycerides to HDL-Cholesterol Predicts Extensive Coronary Disease. *Clinics*; 63(4): 427-432.
- Sastroasmoro, S., 2008, Pemilihan Subyek penelitian. Dalam : S. Sastroasmoro, S. Ismael, *Dasar-dasar Metodologi Penelitian Klinis*, Sagung Seto, Jakarta, 67-76.
- Schiel, R., Beltschikow, W., Radon, S., Kramer, G., Perenthalen, T., Stein, G., 2007. Increased Carotid Intima-Media Thickness and Assosiations with Cardiovascular Risk Factors in obese and overweight Children and Adolescent, *Euro J Med Res*; 12: 503-508.
- Sheriff, Dhastagir, Sultan., Sachu, P., Elshaari F.A., 2013. HDL, apo B/apo A1 ratio, Diabetes Mellitus and Cardiovascular Disease. *INTECH*; 8: 172-194.
- Sibernagl, S., Lang, F., 2007. *Teks Dan Atlas Patofisiologi: Penyakit Jantung Koroner*. EGC; Jakarta:218-20, 286, 290.
- Siedel, J., 1993. AACC Meeting Abstract. *Clin Chem*; 39: 1127
- Sniderman, A.D., Williams, K., Contois, J.H., Monroe, H.M., McQueen, M.J., de Graaf, J., Furberg, C.D., 2011. A Meta-Analysis of Low-Density Lipoprotein Cholesterol, Non-High Density Lipoprotein Cholesterol, and Apolipoprotein B as Marker of Cardiovascular Risk. *Circulation*; 4:337-345.
- Soegondo, S., Soewondo, P., Subekti, I., 2002. *Petunjuk Praktis Pengelolaan Diabetes Melitus Tipe 2*. Perkumpulan Endokrinologi Indonesia (PERKENI). Jakarta.
- Solonen, R., Solonen, J.T., 1990. Progression of carotid atherosclerosis and its determinants: a population-based ultrasonography study. *Atherosclerosis*, 81: 33-40.
- Solonen, R., Solonen, J.T., 1993. Ultrasound B-mode imaging in observational studies of atherosclerotic progression. *Circulation*, 87(3):56-65.
- Spagnoli, L.G., Bonanno, E., 2007. Role of Inflammation in Atherosclerosis. *J. Nuclear Med*; 48(11):1800-15.
- Stein, J.H., Korcarz, C.E., Hurst, R.T., 2007. Use of Carotid Ultrasound to Identify Subclinical Vascular Disease and Evaluate Cardiovascular Disease Risk: A the Carotid Intima-Media Thickness Task Force Endorsed by the Society of Echocardiography Carotid Intima-Media Thickness Task Force Endorsed by the Society for Vascular Medicine. *JASE*; 21:93-111.
- Stompor, T., Krasniak, A., Sulowick., W, Dembin, A., Janda, K., Wojcik, K., 2007. Changes in common carotid artery intima-media thickness Over in 1 Patients On Peritoneal Dialysis. *Nephrol Dial Transplant*; 20(2):1495-500.
- Sugiuchi, H., Uji, Y., Okabe, H., Irie, T., 1995. Direct Measurement of High-Density Lipoprotein Cholesterol in Serum with Polyethylene Glycol-Modified Enzymes and Sulfated α -Cyclodextrin. *Clin Chem*; 41:717-723.
- Suhartono, T., 2008, *Hiperkolesterolemia dan Aterosklerosis*. BP UNDIP, Semarang;161-9.

- Suyono, S., 2009. *Diabetes Melitus Di Indonesia*, Departemen Ilmu Penyakit Dalam Fakultas Kedokteran Universitas Indonesia, Jakarta, 5 : 1874-1878.
- Tahara, N., Yamagishi, S., Matsui, T., Takeuchi, M., Nitta, Y., Kodama, N., *et al.* 2012. Serum levels of advanced glycation end products (AGEs) are independent correlates of insulin resistance in nondiabetic subjects. *Cardiovasc Ther*; 30:42-48.
- Tapp, R., Shaw, J., Zimmet, P., 2003. *Complications of Diabetes*. Diabetes Atlas. Edisi ke-2. Belgium: International Diabetes Federation:72-112.
- Teramoto, T., Sasaki, J., Ueshima, H., Egusa, G., Kinoshita, M., Shimamoto, K., 2007. Diagnostic Criteria for Dyslipidemia : Executive Summary of Japan Atherosclerosis Society (JAS) Guideline for Diagnosis and Prevention of Atherosclerotic Cardiovascular Diseases for Japanese. *J Atheroscler Thromb*. 14:155-158.
- Thevenod, F., 2008. Pathophysiology of Diabetes Mellitus Type 2 : Roles of Obesity, Insulin Resistance and β -cell dysfunction. *Journal of Clinical Endocrinology and Metabolism*; 19: 1-18
- Tomkin, G.H., Owens, D., 2012. The chylomicron: Relationship to atherosclerosis. *Int. J. Vasc. Med*; 10:1155
- Tugasworo, Dodik., 2010, *Patogenesis Aterosklerosis*. Semarang : Badan Penerbit Universitas Diponegoro; 3 – 24.
- Wild, S., Roglic, G., Green, A., Sicree, R., 2004. Global Prevalence of Diabetes, Estimates for the year 2000 and projections for 2030. *Diabetes Care*; 27:1047–1053.
- Williams, K.S., Wagenknecht, Haffner, S.M., 2003. Comparison of the Associations of Apolipoprotein B and Low-Density Lipoprotein Cholesterol With Other Cardiovascular Risk Factors in the Insulin Resistance Atherosclerosis Study (IRAS). *Circulation*; 108: 2312-6.
- World Health Organization (WHO), 1999, *Definition, Diagnosis and Classification of Diabetes Mellitus and its Complications*, 1-65.
- Yamada, N., 2001. Control of triglyceride. *Asian Med. J*; 44:1.
- Yamasaki, Y., Kodama, M., Sakamoto, K., Matsuhisa, M., Kajimoto, Y., Kosugi, K., Shimizu, Y., Kawamori, R., Hori, M., 2000. Carotid Intima-Media Thickness in Japanese Type 2 Diabetic Subjects. *Diabetes Care*; 23: 1310-1315.
- Yuliani, F., Oenzil, F., Iryani, D., 2014. Hubungan Berbagai Faktor Risiko Terhadap Kejadian Penyakit jantung Koroner Pada Penderita Diabetes Mellitus Tipe 2. *J Kes Andalas*; 3(1): 24-31.
- Yokoyama, H., Katakami, N., Yamasaki, Y., 2006, Recent Advances of Intervention to Inhibit Progression of Carotid Intima-Media Thickness in Patients With Type 2 Diabetes Mellitus, *Stroke Journal*; 37:2420-7.
- Zilvermit., D.B., 1999. A proposal linking atherogenesis to the interaction of endothelial lipase with triglyceride-rich lipoprotein. *Circ. Res*; 17: 413-418.