UNIVERSITAS GADJAH MADA
PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

CONTENTS

APPROVAL ii
ACKNOWLEDGMENTS iv
DECLARATION vi
ABSTRACT vii
SARI ix
ABBREVIATIONS AND SYMBOLS xi
CONTENTS xv
LIST OF FIGURES xx
LIST OF TABLES xxx

CHAPTER 1: INTRODUCTION 1
1.1 Background 1
1.2 Location and accessibility 4
1.3 Exploration history 4
1.4 Previous investigations 5
1.5 Objective of the research 7
1.6 Benefits of the research 7
1.7 Dissertation organization 8
1.7.1 Chapters 8
1.7.2 Abbreviations, symbols, figures, tables and appendix 9

CHAPTER 2: REGIONAL GEOLOGY 10
2.1 Indonesian magmatic arcs 10
2.2 Geological framework of Java island 11
2.2.1 Magmatism and volcanism 11
2.2.2 Geological structures 16
2.2.3 Mineralization 18
2.3 Geology of the Southern Mountains of East Java 20

CHAPTER 3: BACKGROUND THEORY 27
3.1 Porphyry Cu-Au ore type deposit 27
3.1.1 Characteristics 27
3.1.2 Tectonic setting 28
3.1.3 Intrusive rocks association 29
3.1.4 Age relationship 29
3.1.5 Hydrothermal alteration model 30
3.1.6 Alteration and mineralization pattern 31

UNIVERSITAS GADJAH MADA
PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

3.1.7 Porphyry vein/veinlet types 34
3.2 Epithermal deposit 37
3.2.1 Characteristics 37
3.2.2 Tectonic setting 37
3.2.3 Classification 37
3.2.4 Alteration types 41
3.3 Geochemical evolution of porphyry-epithermal fluids 41
3.3.1 Magmatism related hydrothermal fluid occurences 41
3.3.2 Early stage 42
3.3.3 Late stage 46

CHAPTER 4: RESEARCH METHODOLOGY 47
4.1 Desk study 47
4.2 Fieldwork 47
4.2.1 Equipment for fieldwork 47
4.2.2 Geological surface mapping 48
4.2.3 Drilling core logging 50
4.2.4 Rock sampling 51
4.3 Laboratory work 52
4.3.1 Petrographic and ore microscopic analysis 52
4.3.2 QEMSCAN analysis 53
4.3.3 Whole rock chemistry analysis 53
4.3.4 Mineral chemistry analysis 54
4.3.5 Fluid inclusion analysis 55
4.3.6 X-Ray diffraction (XRD) analysis 56
4.4 Data compilation and analysis 57

CHAPTER 5: GEOLOGY OF THE RANDU KUNING AREA 61
5.1 Geomorphology 61
5.2 Stratigraphy 63
5.2.1 Tuff unit 65
5.2.2 Andesitic lava unit 66
5.2.3 Pumice breccia unit 67
5.2.4 Calcareous sandstone unit 68
5.2.5 Intrusive rocks and hydrothermal breccia unit 69
5.2.5.1 Hornblende-pyroxene diorite 70
5.2.5.2 Hornblende microdiorite 73
5.2.5.3 Quartz diorite 73
5.2.5.4 Hydrothermal breccia 74

UNIVERSITAS GADJAH MADA
PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

5.2.6 Volcanic rock of Lawu volcano 75
5.2.7 Alluvial deposit 76
5.3 Geological structures 76
5.4 Summary 80

CHAPTER 6: PETROLOGY AND GEOCHEMISTRY OF THE INTRUSIVE ROCKS 81
6.1 Hornblende-pyroxene diorite 85
6.1.1 Petrology of the hornblende-pyroxene diorite 85
6.1.2 Bulk-rock geochemistry of the hornblende-pyroxene diorite 87
6.2 Hornblende microdiorite 88
6.2.1 Petrology of the hornblende microdiorite 88
6.2.2 Bulk-rock geochemistry of the hornblende microdiorite 90
6.3 Quartz diorite 91
6.3.1 Petrology of the quartz diorite 91
6.3.2 Bulk-rock geochemistry of the quartz diorite 91
6.4 Summary 93

CHAPTER 7: HYDROTHERMAL ALTERATION AND MINERALIZATION 96
7. 1 Hydrothermal alteration types 96
7. 1.1 Magnetite+biotite±K-feldspar±chlorite (potassic) zone 98
7. 1.1.1 Mineralogy of the potassic zone 98
7. 1.1.2 Geochemistry of the potassic zone 102
7. 1.2 Chlorite+magnetite±actinolite± carbonate (inner propylitic) zone 106
7. 1.2.1 Mineralogy of the inner propylitic zone 106
7. 1.2.2 Geochemistry of the inner propylitic zone 109
7. 1.3 Chlorite+epidote+carbonate (outer propylitic) zone 112
7. 1.4 Chlorite+sericite+magnetite±actinolite zone 113
7. 1.4.1 Mineralogy of the Chlorite+sericite+magnetite±actinolite zone 113
7. 1.4.2 Geochemistry of the Chlorite+sericite+magnetite±actinolite zone 114
7. 1.5 Sericite+quartz+pyrite (phyllic) zone 114
7. 1.5.1 Mineralogy of the sericite+quartz+pyrite zone 115
7. 1.5.2 Geochemistry of the sericite+quartz+pyrite zone 116
7. 1.6 Illite+kaolinite±smectite (intermediate argillic) zone 118
7. 1.7 Illite+kaolinite±pyrophyllite±alunite (advanced argillic) zone 121
7. 1.8 Quartz+chlorite (silisic) zone 123
7. 2 Elemental mass balance 124
7. 2.1 Element mass balance in hornblende microdiorite 128

UNIVERSITAS GADJAH MADA
PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

7. 2.2 Element mass balance in hornblende-pyroxene diorite..... 133
7. 2.3 Element mass balance in quartz diorite..... 134
7. 3 Vein types 136
7. 3.1 Porphyry Cu-Au vein types..... 136
7. 3.2 Epithermal Au vein types..... 138
7. 4 Breccia types 141
7. 4.1 Magmatic hydrothermal breccia..... 141
7. 4.2 Phreatomagmatic breccia 143
7. 5 Paragenetic sequence 146
7. 5.1 Porphyry stages sequence..... 146
7. 5.2 Epithermal stage sequence 147
7. 6 Mineralization prospects 148
7. 6.1 Randu Kuning prospect..... 149
7. 6.2 Randu Kuning South prospect 152
7. 6.3 Gawe prospect..... 152
7. 6.4 Bukit Piti-Tumbu prospect..... 153
7. 6.5 Geblag prospect..... 154
7. 6.6 Jangglengan prospect 155
7. 6.7 Lancip-Kepil prospect..... 156
7. 6.8 Melikan prospect..... 156
7. 7 Mineralization style..... 158
7. 8 Summary 163

CHAPTER 8: GEOTHERMOBAROMETRY 168
8.1 Introduction..... 168
8.2 Geothermometry 168
8.2.1 Amphibole-plagioclase thermometer..... 168
8.2.2 Calcic amphibole thermometer 170
8.2.3 Chlorite thermometer 170
8.2.4 Fluid inclusion 172
8.2.4.1 Petrography 174
8.2.4.2 Microthermometric analysis..... 178
8.3 Geobarometry..... 185
8.4 Summary 186

CHAPTER 9: DISCUSSION 187
9.1 Magmatism..... 187
9.1.1 Correlation of magmatism event to regional tectonism 187
9.1.2 Chronology and emplacement of the intrusive rocks 188

PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

9.1.3 Magma afinities 190
9.2 Hydrothermal alteration and mineralization 192
9.2.1 Origin of hydrothermal alteration and mineralization 192
9.2.2 Hydrothermal ore fluid evolution 193
9.2.2.1 Early porphyry stage 195
9.2.2.2 Midle porphyry stage 197
9.2.2.3 Late porphyry stage 198
9.2.2.4 Intermediate sulphidation epithermal stage 198
9.2.3 Genetic model for Randu Kuning hydrothermal system 198
9.2.3.1 Co-existing porphyry and epithermal system 198
9.2.3.2 Porphyry system 199
9.2.3.2.1 Tectonic setting 200
9.2.3.2.2 Intrusive rocks association 201
9.2.3.2.3 Age relationship 201
9.2.3.2.4 Hydrothermal alteration model 202
9.2.3.3 Epithermal system 205
9.2.3.3 Comparison to other well known Cu-Au porphyry deposits 207

CHAPTER 10: CONCLUSIONS AND RECOMMENDATIONS 210
10.1 Conclusions 210
10.1.1 Magmatism 211
10.1.2 Hydrothermal alteration and mineralization 209
10.1.3 Hydrothermal deposits genetic model 210
10.2 Recommendations 212
10.2.1 Next exploration taget 212
10.2.2 Next research 212

REFERENCES 214

APPENDIX
1. Sample location, Geological and Alteration Zone maps 230
2. List of petrographic and ore microscopic analysis 241
3. List of X-Ray diffractometer and Qemscan analysis 252
4. Rock geochemistry analysis 260
5. Mineral geochemistry analysis 263
6. Fluid inclusions analysis 289

UNIVERSITAS GADJAH MADA
PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

LIST OF FIGURES

Figure 1.1	Location map of the Randu Kuning area, Selogiri district, Wonogiri regency.	4
Figure 2.1	Magmatic arcs distributing in Indonesia (redrawn after Carlile and Mitchell, 1994) and different styles of mineral deposits in Indonesia (Carlile and Mitchell, 1994; Setijadji and Maryono, 2012).	12
Figure 2.2	The distribution of the Tertiary magmatic arcs, Tertiary volcanic rocks and intrusions, center of the Cenozoic volcanos as well as basement crust types of Java island..	15
Figure 2.3	The main structural patterns in Java island, compiled from Pulunggono and Martodjoyo (1994), Satyana (2007), Sribudiyani *et al*. (2003), Setijadji and Maryono (2012).	17
Figure 2.4	The different styles of mineral deposits distribution in Java island (Modified from Setiawan and Yudawinata, 2000; Setijadji *et al*., 2006; Setijadji and Maryono, 2012).	19
Figure 2.5	Simplified geological map of East Java, showing the main geological provinces (Southern Mountains Arc, Modern Arc, Kendeng Basin, Sunda Shelf) and stratigraphic units (Smith *et al*., 2008).	21
Figure 2.6	Regional geological map of the Southern Mountains of East Java in Wonogiri area and its vicinity (compiled and modified from Surono *et al*., 1992; Samodra *et al*., 1992; Sampurna and Samodra, 1997).	23
Figure 2.7	Simplified stratigraphic column of the Southern Mountain Arc, shows three synthems and indicates the phases of arc development recorded by the sedimentary succession (Smyth *et al*., 2008).	25
Figure 3.1	Tectonic setting models in porphyry ore deposit.	28
Figure 3.2	Alteration and mineralization schematic of the porphyry ore deposit of San Manuel Kalamazoo (Gilbert and Lowell, 1974)	34
Figure 3.3	Schematic sequence of the formation of the types of vein-veinlets in A) deposition of porphyry Cu-Mo and B) Cu-Au deposition associated with calc-alkaline intrusions.	35
Figure 3.4	Shows the chronology of the formation of hydrothermal fluids associated with magmatism, based on models drawn from Sillitoe and Hedenquist (2003) and Heinrich (2005).	45
Figure 4.1	The research methodology flowchart.	49
Figure 5.1	The aster imagery of the Selogiri area	61

PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Figure 5.2	Photograph the landscape of the Selogiri area. A). Top photo was taken from the Tiyaran hill in the northern part of the area, B). Bottom photo was taken from Randu Kuning hill to the south.	62
Figure 5.3	Stratigraphic column of the Selogiri area	63
Figure 5.4	Geological Map of Selogiri area and its vicinity (compiled and modified from Hartono, 2010)	64
Figure 5.5	Many kind of rock types within Tuff unit.	65
Figure 5.6	A). Propylitic altered andesitic lava cut by lateral slip fault. Location Melikan (ST. 145). B). Volcanic breccia with andesitic rock fragments supported by volcanic matrix.	67
Figure 5.7	A weak propylitic altered andesitic lava showing collumnar joint, vesiculer and amigdaloidal structures . Location Randubang.	67
Figure 5.8	Many rock types within Pumice Breccia rock unit. a) Faulted intercalation of pumice breccia, lapilli-tuff and tuff, b). Pumice lapilli-tuff intercalated by pumice breccia, contain much of pumice and andesite fragments.	68
Figure 5.9	Some rock types of calcareous sandstone rock unit. a). Contact betweeen calcareous tuff and nodule bearing calcareous lapilli-tuff. b). Inset nodule bearing calcareous lapilli-tuff. c). Calcareous tuffaceous sandstone.	69
Figure 5.10	Hornblende-pyroxene diorite of the Tumbu and Geblag hill intruded by hornblende-microdiorite of the Randu Kuning hill.	70
Figure 5.11	Intrusive rocks at the Randu Kuning area. A). Hornblende-pyroxene diorite have a medium grained size occured before mineralization stage (pre-mineralization). B). Horblende microdiorite is syn-mineralization intrusive rock and C). Quartz diorite as post-mineralization intrusive rock.	71
Figure 5.12	Geological map of the Randu Kuning area and geological cross section line B-B' based on geological map and drilling cross sections.	72
Figure 5.13	Andesitic intrusive rock of Gunung Kukusan. This intrusion is possible occured as co-magmatic with the hornblende-pyroxene diorite (pre-mineralized intrusive rock).	73
Figure 5.14	Contact between intrusive rocks in the Randu Kuning area.	74
Figure 5.15	Gold bearing magmatic hydrothermal breccia cross cut by quartz-pyrite (epithermal) vein at the Kaloran area.	75
Figure 5.16	Phreatomagmatic breccia level epithermal environment with clast diorite and juvenil (WDD 34-124.50).	75
Figure 5.17	Volcanic breccia of the Lawu volcano, showing many angular of	76

PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

andesite fragments. Insert figure is an andesite fragment comprising of hornblende and plagioclase phenocrysts set in the weathered goundmas).

Figure 5.18 Alluvial deposit used for agricultural land in front of pumice breccia's hills. Location Bulu. 76

Figure 5.19 Contact quartz diorite and altered intercalatin sandstone and claystone of Kebo-Butak formation cross-cut by NE-SW (N32°E left lateral-slip (sinistral) fault. 77

Figure 5.20 Intercalation of the volcanic sandstone and pumice breccia cross-cut by a righ lateral normal slip fault. Location Tiyaran village, Bulu. 77

Figure 5.21 Hornblende-pyroxene diorite cross-cut by a N-S (N358E/82) right lateral slip (dextral) fault. In the dilational jog fault zone found many of brecciation, carbonate-sphalerite±silica-galena-chlorite veins. Location River Melikan. 78

Figure 5.22 Strong altered (chlorite-epidote) brecciated volcanic rock cross cut by N-S (N330E/80- N358E/82-) right lateral-slip fault. In the fault zone found many of brecciation, carbonate- red sphalerite-silica-epidote-chlorite veins, and pyrite-chalcopyrite disseminations. Location River Melikan. 78

Figure 5.23 Illustration of the veins movement in part of normal and reverse slip fault sinistral. Bore hole WDD 19-157.25 and 179.30. 79

Figure 5.24 Illustration of the veins movement in part of normal slip fault sinistral. Bore hole WDD 45-360.40. 79

Figure 6.1 Harker variation diagram for major-oxide elements (in wt.%) vs. SiO2 (in wt.%) in least-altered rock types in the Randu Kuning area. 82

Figure 6.2 A). The SiO_2-(Na_2O+K_2O) rock classification diagram (Midlemost, 1975) and B) The Zr/TiO_2-Nb/Y rock classification diagram (after Winchhester and Floyd, 1977) for least altered dioritic intrusive rocks of the study area 83

Figure 6.3 A).The Th/Yb-La/Yb tectonic setting diagram (after Condie, 1989), and B) Zr-Sr/2-Ti/100 affinity discrimination diagram (after Cann, 1973) showing all the sample are plotted as an island arc setting and calc-alkaline basalt field. 83

Figure 6.4 A) N-type MORB normalized (Sun, 1980; Saunders and Tarney, 1984) spider diagram for least altered intrusive rocks in the studied area, and B) REE-chondrite normalized (Sun and McDonough, 1989) pattern for the same samples 84

UNIVERSITAS GADJAH MADA
PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Figure 6.5	Least altered hornblende-pyroxene diorite shows a grey colour, equigranular, comprise plagioclase, pyroxene and hornblende as primary mineral, medium grained size. Sample: WDD 02-128.50.	85
Figure 6.6	Photomicrograph of least altered hornblende-pyroxene diorite. A). Plane polarized light (PPL). B) Crossed polarized light (XPL).	85
Figure 6.7	Backscatter image showing co-exisiting plagioclase and hornblende in hornblende-pyroxene diorite and the distribution of microprobe measurment point both in hornblende and plagioclase.	86
Figure 6.8	Calcic amphibole classification Si-Mg/(Mg+Fe2) diagram. The amphiboles in this rock sample are ploted within edenite hornblende to pargasite hornblende field.	87
Figure 6.9	Photograph and photomicrograph of least altered hornblende microdiorite. A) Photograph of least altered hornblende microdiorite's hand spicement from Randu Kuning hill (ST.1), B) Photograph of drilling core sample of hornblende microdiorite (WDD 45-31.20). C) Plane polarized light photomicrograph and D) Crossed polarized light photomicrograph.	89
Figure 6.10	Backscatter image across the plagioclase showing oscilatory zoning structure. Point 1=$An_{43.7}$; 2= $An_{53.9}$; 3=$An_{52.8}$; 4=$An_{49.8}$ and point 5=$An_{49.5}$.	89
Figure 6.11	Photograph and photomicrograph of least altered quartz diorites	92
Figure 7.1	Alteration zone map of the Randu Kuning area and its vicinity.	97
Figure 7.2	A) Magnetite+biotite±K-feldspar±chlorite (potassic) altered microdiorite cut by quartz vein and quartz-carbonate veinlets with chlorite-silica-epidote halos. Bottom: Photomicrograph of potassic altered hornblende-pyroxene diorite. B) Plane polarized light. C) Crossed polarized light.	99
Figure 7.3	Photomicrograph of polished section under the reflected light of the oxides and sulphides assemblages. A) Magnetite-chalcopyrite-bornite. B) Magnetite-hematite-chalcopyrite, where mostly magnetites were replaced by hematites.	99
Figure 7.4	Qemscan image of the magnetite+biotite±K-feldspar±chlorite (potassic)-altered hornblende microdiorite was overprinted by inner propylitic and intermediate argillic mineral assemblages.	100
Figure 7.5	A). Crossed polarized light photomicrograph of magnetite+biotite± K-feldspar±chlorite (potassic) altered hornblende-pyroxene diorite. B). Photomicrograph of polished section under the reflected light of magnetite cross cut by pyrite and chalcopyrite.	101

PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Figure 7.6	Qemscan image of A). The magnetite+biotite±K-feldspar±chlorite (potassic)-altered hornblende-pyroxene diorite. B). Closed up the image of figure A.	102
Figure 7.7	Harker variation diagram for major-oxide elements (in wt.%) vs SiO_2 in least-altered and altered hornblende microdiorite rocks.	103
Figure 7.8	Harker variation diagram for major-oxide elements (in wt.%) vs SiO_2 in least-altered and altered hornblende-pyroxene diorite rocks.	104
Figure 7.9	A) N-Type MORB normalized (Sun, 1980; Saunders and Tarney, 1984) spider diagram for least, magnetite+biotite±K-feldspar±chlorite (potassic), chlorite+magnetite+±actinolit± carbonate (inner propylitic) and Sericite+quartz+pyrite (phyllic) altered hornblende microdiorite, and B) REE-chondrite normalized (Sun and McDonough, 1989) pattern for the same samples.	105
Figure 7.10	Photomicrograph of the chlorite-magnetite-silica-epidote(inner propylitic)) altered hornblende microdiorite.	107
Figure 7.11	Propylitic (chlorite+magnetite±actinolite±carbonate)-altered hornblende-pyroxene diorite at Gritan village.	107
Figure 7.12	A). Crossed polarized light photomicrograph of the chlorite+magnetite±actinolite±carbonate (inner propylitic)-altered hornblende-pyroxene diorite. B). Crossed polarized light. photomicrograph polished section under the reflected light of sphalerite-chalcopyrite-galena-pyrite of carbonate-sulphides vein within inne propylitic-altered hornblende-pyroxene diorite.	108
Figure 7.13	Photomicrograph of chlorite+magnetite±actinolite±carbonate (inner propylitic)-altered quartz diorite A). Crossed polars light B). Plane polarized light.	108
Figure 7.14	Qemscan image of the chlorite+magnetite±actinolite±carbonate (inner propylitic)-altered quartz diorite, consist of many secondary minerals such as chlorites, quartzes, kaolinites, calcites and hematites. Ore minerals are not found within this section.	109
Figure 7.15	A). N-Type MORB normalized (Sun, 1980; Saunders and Tarney, 1984) spider diagram for least altered and the chlorite+magnetite±actinolite±carbonate (inner propylitic) and sericite+quartz+pyrite (phyllic) -altered hornblende-pyroxene diorite as well as least altered and chlorite+epidote+carbonate (outer propylitic)-altered quartz diorite B). REE-chondrite normalized (Sun and McDonough, 1989) pattern for the same	111

PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

samples..

Figure 7.16 A). Photo of drilling core handspicement of the chlorite+epidote+carbonate (outer propylitic)-altered hornblende-pyroxene diorite, cut by epidote-chlorite-carbonate veinlet. Bottom: Photomicrograph of chlorite+epidote+carbonate (outer propylitic)-altered hornblende-pyroxene diorite. 113

Figure 7.17 A).Photograph of drilling core handspicement of the chlorite+sericite+magnetite±actinolite altered hornblende microdiorite. Bottom: photomicrograph of the sample above. 114

Figure 7.18 The sericite+quartz+pyrite (phyllic)-altered hornblende-pyroxene diorite cross cut by pyrite-quartz-carbonate vein and quartz veinlets. 115

Figure 7.19 A). Phyllic-altered hornblende microdiorite associated with pyrite-quartz-carbonate vein. Bottom: photomicrograph of phyllic altered hornblende microdiorite (silica+sericite+pyrite± carbonate±chlorite) of the same sample. 116

Figure 7.20 A). Intermediate argillic altered hornblende-pyroxene diorite from the southern part of Geblag hill (left) and B). Intermediate argillic-altered hornblende-pyroxene diorite crackel breccia. 119

Figure 7.21 XRD graphic analysis of illite+kaolinite ±smectite (intermediate argillic) altered hornblende-pyroxene diorite. A) Bulk powder analysis, B) Clay treatment EG, C). Clay treatment AD. 120

Figure 7.22 Qemscan image of the illite+kaolinite ±smectite (intermediate argillic)- altered hornblende-pyroxene diorite, comprise secondary minerals such as kaolinite, illite, dolomite, sericite, pyrite and pyrrhotite. 121

Figure 7.23 A). The Illite+kaolinite±pyrophyllite±alunite (advanced argillic)-altered hornblende-pyroxene diorite and breccia of Lancip hill (top). B). Hand spicements of advanced argillic altered-oxidized hydrothermal breccia (bottom left) and C). Contact of oxidized vuggy silica and pyrophyllite+alunite minerals assemblages. 122

Figure 7.24 Photomicrograph of the illite+kaolinite±pyrophyllite±alunite (advanced argillic)-altered brecciated rock (silica+pyrophyllite± carbonate±chlorite). 123

Figure 7.25 Hand spicements of the silisic altered rocks. A). Silica selvage and veins. B). Initial fluidized phreatomagmatic hydrothermal related silisic zone cut by chalcopyrite+sphalerite+pyrite vein. C). Silisic rock cut by pyrite-quartz vein with chlorite halos. 124

Figure 7.26 Photomicrograph of the quartz+chlorite (silisic)-altered rock 124

UNIVERSITAS GADJAH MADA
PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

(silica+sericite±carbonate±chlorite).
Figure 7.27 Diagram of the hydrothermal fluid-wall rocks interaction processes. 125
Figure 7.28 Mass balance plots between the least altered and magnetite+biotite±K-feldspar±chlorite (potassic) altered of hornblende microdiorite. 130
Figure 7.29 Mass balance plots between the magnetite+biotite±K-feldspar±chlorite (potassic) and sericite+quartz+pyrite (phyllic) altered of hornblende microdiorite. 131
Figure 7.30 Mass balance plots between the least altered and M vein within magnetite+biotite±K-feldspar±chlorite (potassic) zone of hornblende microdiorite. 132
Figure 7.31 Mass balance plots between the least altered and A vein type within magnetite+biotite± Kfeldspar± chlorite (potassic) zone of hornblende microdiorite. 133
Figure 7.32 Concentration change ($\Delta C/C_{0i}$) between the propylitic-altered hornblende-pyroxene diorite and the phyllic-altered hornblende-pyroxene diorite. 134
Figure 7.33 Concentration change ($\Delta C/C_{0i}$) between the least altered quartz diorite and the propylitic -altered quartz diorite. 135
Figure 7.34 Many porphyry vein types at the Randu Kuning area. 138
Figure 7.35 Many epithermal vein types at the Randu Kuning area. 139
Figure 7.36 Magmatic hydrothermal breccia stages. 142
Figure 7.37 A) Contact hornblende-pyroxene diorite with phreatomagmatic breccia. B) Phreatomagmatic breccia. C) Porphyry preatomagmamatic breccia cut by carbonate-pyrite veins. 145
Figure 7.38 A) Phreatomagmatic breccia epithermal environment with clast consist of diorite and juvenil. B) Phreatomagmatic breccia with diorite, juvenil, quartzite, schist clasts, cut by carbonate-sphalerite-pyrite-galena vein. C) Rebrecciated of phreatomagmatic breccia. 145
Figure 7.39 The landscape of the ore deposits prospect in the Randu Kuning porphyry Cu-Au and its vicinity. 149
Figure 7.40 The mineralization prospect area map and core drilling hole location. 149
Figure 7.41 Some veins type in the Randu Kuning prospect area. 150
Figure 7.42 Some veins type in the Gawe prospect area. 153
Figure 7.43 Some veins type in the Bukit Piti prospect area. 154
Figure 7.44 Some veins type in the Geblag prospect area. 155
Figure 7.45 Some veins type in the Jangglengan prospect area. 155

UNIVERSITAS GADJAH MADA
PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Figure 7.46	Some vein types in the Lancip-Kepil prospect.	156
Figure 7.47	A) Intercalation of tuffaceous sandstone and siltstone of Kebobutak formation were intruded by quartz diorite. B) Hand spicement of argillic altered hydrothermal berccia cut by many stages of sphalerite-carbonate+pyrite+chalcopyrite veins at the Melikan area.	157
Figure 7.48	Some barren quartz-magnetite veins in the deeper portion of hornblende microdiorite intrusion.	159
Figure 7.49	Two type of disseminated sulphides mineralization.	160
Figure 7.50	A) Propylitic-altered microdiorite cut by pyrite-chalcopyrite C style veinlets with silica-chlorite-carbonate halos. B Quartz-pyrite-carbonate (D style vein) with sericite selvage.	160
Figure 7.51	Photomicrograph polished section under the reflected light of some samples containing oxides, sulphides and gold. A) Free gold as inclusion in gangue mineral grains among magnetites. B). Bornite is replaced by chalcopyrites associated with magnetite. C). Magnetite grains are replaced by pyrite and chalcopyrite. D).Inclusion free gold within magnetite.	161
Figure 7.52	Epithermal veins type which contribute in Au-Ag base metals mineralization. A) Pyrite-chalcopyrite-silica-carbonate vein with sericite selvage. B) Pyrite-sphalerite-carbonate vein with sericite selvage.	163
Figure 7.53	Photomicrograph polished section under the reflected light of sample containing sulphides and gold. A) Sphalerite-galena-pyrite-chalcopyrite assemblage and free gold grain replaced galena. Sphalerite was relaced by chalcopyrite and galena. B). Magnification of image A.	163
Figure 8.1	Ten quartz vein samples were used for fluid inclusions analysis.	175
Figure 8.2	Petrographic fluid inclusion analysis scheme. A) Showing the area where fluid inclusions abundant on double polished thin section photo scan. B) Microscopic photo of fluid inclusions type within one of quartz vein spot (sample 41). Bottom: examples of the fluid inclusions occurences in the studied area. C). Showing pseudo-secondary and secondary fluid inclusions and D) primary fluid inclusions (sample 1436/WDD49-567.25).	176
Figure 8.3	Many types of fluid inclusion in the area. A). Type II : two phase inclusions, vapor rich and type III : multiphase inclusions, contain liquid, vapor and solid with halite as daughter minerals. B) Type I : two phase inclusions, liquid rich and type III.	177

UNIVERSITAS GADJAH MADA
PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Figure 8.4	Many types of fluid inclusion in the area.	177
Figure 8.5	Frequency plots of homogenisation temperature (Th) and salinity (wt.% NaCl equiv.) for sample: 1345 (WDD 09-141.00).	180
Figure 8.6	Frequency plots of homogenisation temperature (Th) and salinity (wt.% NaCl equiv.) for sample: 1340 (WDD 08-87.05).	180
Figure 8.7	Frequency plots of homogenisation temperature (Th) and salinity (wt.% NaCl equiv.) for sample: 1339(WDD 18-197.80).	181
Figure 8.8	Frequency plots of homogenisation temperature (Th) and salinity (wt.% NaCl equiv.) for sample: 1338(WDD 18-329.50).	181
Figure 8.9	Frequency plots of homogenisation temperature (Th) and salinity (wt.% NaCl equiv.) for. sample: 1337 (WDD 49-366.50).	182
Figure 8.10	Frequency plots of homogenisation temperature (Th) and salinity (wt.% NaCl equiv.) for sample: 1336 (WDD 49-567.25).	182
Figure 8.11	Frequency plots of homogenisation temperature (Th) and salinity (wt.% NaCl equiv.) for sample: 1344 (WDD 19-85.05).	183
Figure 8.12	Frequency plots of homogenisation temperature (Th) and salinity (wt.% NaCl equiv.) for sample: 1338 (WDD 18-329.50).	183
Figure 8.13	Frequency plots of homogenisation temperature (Th) and salinity (wt.% NaCl equiv.) for sample: 1342 (WDD 19-166.90).	184
Figure 8.14	Frequency plots of homogenisation temperature (Th) and salinity (wt.% NaCl equiv.) for sample: 1341 (WDD 19-212.50).	184
Figure 9.1	The north-south geological cross section on Early Miocene east Java arc. It show the correlation of the Early Miocene magmatism, hydrothermal and volcanism event in Randu Kuning, Selogiri to the regional tectonism	188
Figure 9.2	Schematic cross section of the chronology and emplacement of the intrusive rocks in the Randu Kuning area (unscale). A). The emplacement of the hornblende-pyroxene diorite and the early emplacement of hornblende microdiorite, B). The late emplacement of hornblende microdiorite and quartz diorite as well as the latest intrusion of andesite.	189
Figure 9.3	Diagram Sr/Y vs Y (Defant and Drummond, 1993) of the least altered intrusive rocks in Selogiri are compared to representing Cenozoic volcanic centre rock samples from Java island (regional data from Setijadji *et al*., 2006).	191
Figure 9.4	Range of the hydothermal system ore deposit fields the salinity-temperature diagram (Large *et al*., 1988 vide Pirajno, 2009).	194
Figure 9.5	Schematic of magmatism and hydrothermal fluid evolution, alteration zoning as well as veining development form the	196

UNIVERSITAS
GADJAH MADA
PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

porphyry Cu-Au to the epithermal Au-base metals in the Randu Kuning prospect.

Figure 9.6 Schematic model of intrusive rocks, alteration zoning as well as vein types distribution in the Randu Kuning porphyry Cu-Au and the epithermal Au-base metals deposits. 197

Figure 9.7 Randu Kuning porphyry Cu-Au among the porphyry deposits in the world within Au versus Cu grades diagram of Sinclair (2007). 199

Figure 9.8 Comparison the alteration zone of the Randu Kuning porphyry Cu-Au (bottom) with those the porphyry system alteration model (top). 203

UNIVERSITAS GADJAH MADA
PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

LIST OF TABLES

Table 3.1	Alteration and mineralization characteristic of porphyry type deposit (Lowell and Gilbert, 1970; Gilbert and Lowell, 1974; and other sources).	32
Table 3.2	The development of epithermal classification (Sillitoe and Hedenquist, 2003).	39
Table 3.3	Principal field oriented characteristic of epithermal type and subtype (Sillitoe and Hedenquist, 2003).	40
Table 3.4	Summary of hydrothermal alteration assemblages forming in epithermal environment (Simmons *et al.*, 2005).	41
Table 4.1	Relationship of analytical method, information obtained and objectives of analysis.	59
Table 6.1	Petrographic analysis (textures and compositions) and silica content of the intrusive rocks at the Randu Kuning area, Selogiri.	94
Table 6.2	The compatible elements concentrations (ppm) in the intrusive rocks of the Randu Kuning area and its vicinity.	94
Table 6.3	The LILE concentrations (ppm) in intrusive rocks in the Randu Kuning area and its vicinity.	95
Table 6.4	The HSFE elements concentrations (ppm) in the intrusive rocks of the Randu Kuning area and its vicinity.	95
Table 7.1	Isocon analysis of average propylitic altered hornblende-pyroxene diorite to average least altered hornblende-pyroxene diorite composition at Randu Kuning.	129
Table 7.2	Major porphyry and epithermal vein/veinlet types and their characteristic in the Randu Kuning area	140
Table 7.3	Characteristics of the hydrothermal breccia at the Randu Kuning area and its vicinity.	144
Table 7.4	Paragenetic sequence of the hydrothermal veins/veinlets and minerals in the Randu Kuning prospect.	147
Table 7.5	The Mineralization prospect area and their characteristic (Sources: this research; Suasta and Sinugroho, 2011; Muthi, *et al.*, 2012; Nightingale, 2014)	151
Table 8.1	The result of geothermometer based on Holland and Blundy (1994) equation using amphibole-plagioclase compostion in hornblende microdiorite.	170

UNIVERSITAS GADJAH MADA
PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND
EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/

Table 8.2	The result of geothermometer based on Ridolfi, *et al*. (2010) and Ridolfi and Rezulli (2012) using calcic amphibole mineral in hornblende microdiorite.	170
Table 8.3	The result of geothermometer mean based on Cathelineau (1988) and Jowett (1991) used chlorite mineral in hydrothermally altered hornblende microdiorite and hornblende-pyroxene diorite.	171
Table 8.4	List of doubly polished thin section and their fluid inclusion analysis results.	179
Table 8.5	The result of geotherbaromometer based on Holland and Blundy (1994) equation using amphibole-plagioclase compostion in hornblende microdiorite.	185
Table 8.6	The result of geotherbaromometer based on Ridolfi, *et al*. (2010) and Ridolfi and Rezulli (2012) using calcic amphibole mineral in hornblende microdiorite.	186
Table 9.1	Characteristics of the hydrothermal fluid evolution stages in the Randu Kuning porphyry Cu-Au and Epithermal Au-base metals deposits (metal elements grade data are from PT. Alexis Perdana Mineral).	194
Table 9.2	Comparison of the alteration zoning characteristic in Randu Kuning porphyry Cu-Au to the porphyry's alteration zoning standart model and other porphyry Cu-Au in the Eastern Sunda Arch.	204
Table 9.3	Characteristic of the porphyry Cu-Au and epithermal Au deposits in the Randu Kuning area.	205
Table 9.4	Comparison of the characteristic of the Randu Kuning porphyry Cu-Au to the other porphyry Cu-Au in the Eastern Sunda Arc.	208

UNIVERSITAS
GADJAH MADA
PETROLOGY, GEOCHEMISTRY AND HYDROTHERMAL FLUID EVOLUTION OF THE RANDU KUNING PORPHYRY Cu-Au AND EPITHERMAL Au DEPOSITS IN SELOGIRI, CENTRAL JAVA, INDONESIA
SUTARTO, Dr.rer.nat. Arifudin Idrus, S.T.,M.T.; Dr. Agung Harijoko, S.T., M.Eng.; Dr. Lucas Donny Setijadji, S.T., M.
Universitas Gadjah Mada, 2016 | Diunduh dari http://etd.repository.ugm.ac.id/