

Daftar Pustaka

- Ahmed, M., Reda, A. and Ibrahim, M. (2015). Right ventricular function and symptomatology in patients with isolated mitral stenosis: A Doppler tissue imaging study. *The Egyptian Heart Journal*, 67(1), pp.41-46.
- Alkhalifa, M. S., Ibrahim, S. A., & Osman, S. H. (2008). Pattern and severity of rheumatic valvular lesions in children in Khartoum , Sudan, *14*(5), pp.1015–1021.
- Baumgartner, H., Hung, J., Bermejo, J., Chambers, J., Evangelista, A., Griffin, B., Iung, B., Otto, C., Pellikka, P. and Quinones, M. (2009). Echocardiographic assessment of valve stenosis: EAE/ASE recommendations for clinical practice. *European Journal of Echocardiography*, 10(1), pp.1-25.
- Carapetis, J. R. 2008. Rheumatic Heart Disease in Asia. *Circulation*, 118, pp.2748-2753.
- Clerico, A., Passino, C. and Emdin, M. (2011). When Gonads Talk to the Heart. *Journal of the American College of Cardiology*, 58(6), pp.627-628.
- Dahlan, S.M Statistik untuk Kedokteran dan Kesehatan, Jakarta: Salemba Medika, 2009
- Daniels, L. and Maisel, A. (2007). Natriuretic Peptides. *Journal of the American College of Cardiology*, 50(25), pp.2357-2368.
- El Zayat, A. (2014). Potential use of Brain Natriuretic Peptide in patients with asymptomatic significant mitral stenosis. *The Egyptian Heart Journal*, 66(3), pp.269-275.
- Gai, M., Piccoli, G. B., & Segoloni, G. P. (n.d.). coefficient; SPSS). The concordance in detecting pathologic samples was good for erythrocytes, pp.1560–1562.
- Galasko, G. (2005). What is the normal range for N-terminal pro-brain natriuretic peptide? How well does this normal range screen for cardiovascular disease?. *European Heart Journal*, 26(21), pp.2269-2276.
- Iltumur, K., Karabulut, A., Yokus, B., Yavuzkir, M., & Taskesen, T. (2005). N-terminal proBNP plasma levels correlate with severity of mitral stenosis N-Terminal proBNP Plasma Levels Correlate with Severity of Mitral Stenosis, (June 2016).
- Inci, S., Erol, M., Bakirci, E., Hamur, H., Degirmenci, H., Duman, H. and Karakelleoglu, S. (2015). Effect of percutaneous mitral balloon valvuloplasty on right ventricular functions in mitral stenosis: Short- and mid-term results. *Anadolu Kardiyoloji Dergisi/The Anatolian Journal of Cardiology*, 15(4), pp.289-296.

- Kumar, V., Jose, V., Pati, P. and Jose, J. (2014). Assessment of right ventricular strain and strain rate in patients with severe mitral stenosis before and after balloon mitral valvuloplasty. *Indian Heart Journal*, 66(2), pp.176-182.
- Lang, R., Bierig, M., Devereux, R., Flachskampf, F., Foster, E., Pellikka, P., Picard, M., Roman, M., Seward, J. and Shanewise, J. (2006). Recommendations for chamber quantification☆. *European Journal of Echocardiography*, 7(2), pp.79-108.
- Lisy, M., & Babal, P. (2007). Brain natriuretic peptide – the biological marker in the diagnosis of overt congestive heart failure and myocardial ischemia, *108*, pp.170–173.
- Mann, D.L., Zipes, D.P., Libby, P., Bonow. R.O., (2014). Braunwald’s Heart Disease : A Textbook of Cardiovascular Medicine, Elsevier.
- Maulana, G.H. (2013). Perbedaan Rerata Kadar *N-Terminal proB-Type Natriuretic Peptide* (NT-ProBNP) Serum Pada Pasien Hierptensi Dengan Dan Tanpa Hipertrofi Ventrikel kiri Di Poliklinik Jantung RSUP Dr. Sardjito Yogyakarta.
- Morello, A., Lloyd-Jones, D., Chae, C., van Kimmenade, R., Chen, A., Baggish, A., O'Donoghue, M., Lee-Lewandrowski, E. and Januzzi, J. (2007). Association of atrial fibrillation and amino-terminal pro-brain natriuretic peptide concentrations in dyspneic subjects with and without acute heart failure: Results from the ProBNP Investigation of Dyspnea in the Emergency Department (PRIDE) study. *American Heart Journal*, 153(1), pp.90-97.
- Ozkan, A. A.-. (2005). Serum N-Terminal pro-BNP Levels Correlate with Symptoms and Echocardiographic Findings in Patients with Mitral Stenosis, 22(6), pp.473–478.
- Pratiwi, D. I. N. (2010). Uji Diagnostik Pemeriksaan NT pro Natriuretic Peptide (NT proBNP) Untuk Diagnosis Gagal Jantung Kongestif. Yogyakarta: Universitas Gadjah Mada.
- Rudski, L., Lai, W., Afilalo, J., Hua, L., Handschumacher, M., Chandrasekaran, K., Solomon, S., Louie, E. and Schiller, N. (2010). Guidelines for the Echocardiographic Assessment of the Right Heart in Adults: A Report from the American Society of Echocardiography. *Journal of the American Society of Echocardiography*, 23(7), pp.685-713.a
- Sastroasmoro, S., Budiman, I., Madiyono, Bambang., Moeslichan, S., Purwanto, S. (2014). Dasar-dasar Metodologi Penelitian Klinis, edisi 5, Sagung Seto, Jakarta-Bab 17.
- Sharma, V., Stewart, R., Zeng, I., Raffel, C. and Kerr, A. (2011). Comparison of Atrial and Brain Natriuretic Peptide for the Assessment of Mitral Stenosis. *Heart, Lung and Circulation*, 20(8), pp.517-524.

Wafa, A. A., Daoud, E. M., Abofotoh, A., El-sadek, A. M., & Hakim, H. (2015). Correlation Between N-terminal Pro Brain Natriuretic Peptide and Right Ventricular Performance Measured by Doppler Echocardiography after Successful Percutaneous Balloon Mitral Valvuloplasty, pp.1–11.

Willerson, J. (2007). *Cardiovascular medicine*. 1st ed. London: Springer, pp.397-430.