

DAFTAR PUSTAKA

- Abdo, Z.M.A., R.A.Hassan, A.A. El-Salam and S.A.Helmy. 2010. Effect of adding green tea and its aquaeus extract as natural antioxidants to laying hen diet on productive, reproductive performance and egg quality during storage and its content of cholesterol. *Egyrt. Poultry Sci.* vol (30) (IV): 1121-114
- Abdulkarimi, R. And M. Daneshyar. 2012. Teh effects of thyme (*Thymus vulgaris*) extract supplementation in drinking water on iron metabolism in broiler chickens. *J.of Med. Plants Res.* Vol. 6(5), pp. 645-650, 9 February, 2012
- Abe, I., 2002. Inhibition of cholesterol biogenesis by tea polyphenols . *Foods Food Ingredients J. Japan* No. 204. Abstrac
- Adisakwattana,S., J. Intrawangso, A. Hemrid, B. Chanathong, and K. Makynen. 2012. Extract of edible plant inhibit pancreatic lipase, cholesterol esterase and cholesterol micellization and bind bile acid. *Food Technol. Biotechnol.* 50 (1) : 11-16
- Aengwanich, W., M. Suttajit, and N.A. Narkkong. 2009. Effects of polyphenols extracted from tamarind (*tamarindus indica* l) seed coat on difeferential white blood cell count in broilers (*Gallus domesticus*) exposed to high environmental temperature. *Intern. J. Of Poultry Sci.* 8 (10) : 957 - 962
- Ahadi, F., S. Chekani-Azar, H. A. Shahryar, A. Lotfi, N.H. Mansoud, and Y. Bahrawi . 2010. Effect of Dietary Supplementation with fish oil with Selenium or Vitamin E on Oxidative Stability and Consumer Acceptability of Broiler Meat. *Global Vet.* 4 (3): 216-221
- Ajuyah, A.O., D.U. Ahn, R.T. Hardin, and J.S.Sim. 1993. Dietary antioxidants and storage affect chemical characteristics of ω -3 fatty acid enriched broiler chicken meats. *J. of Food Sci.,* 58 (1) : 43-46
- Al-Daraji, H.J.,H.A. Al-Mashadani, A.S. Al-Hassani, H.A. Mirza, and W.K. Al-Hayani. 2012. The influence of parsley (*Petroselinum crispum*) as feed additive on hematological traits of local Iraqi geese. . *Adv. Nutr. Res.* 1(1): 1-5.
- Amarowicz,R., M. Naczka and F. Shahidi. 2000. Antioxidant activity of Crude Tannin of Canola and Rapeseed Hulls. *JAOCS*, 77: 957-961.
- Al-Najdawi, R . and B. Abdullah . 2002. Proximate composition, selected minerals, cholesterol content and lipid oxidation of mechanically and hand deboned chickens from teh Jordanian Market . *J. Meat Sci.* 61, 243- 247
- Anesini., C., G.E.Ferraro and R. Filip. 2008. Total polyphenol content and antioxidant capacity of commonly available tea (*Camelia sinensis*) in Argentina. *J. Agric. Food. Chem.* (56): 9225-9229

- AOAC. 1995. Official Method of Analysis of The Association of Chemical Analytic Chemist. Inc. Arlington Virginia. USA
- Arbiati, R., T.S. Utami, A. Kusuma, A. Sinaga. 2007. Comparative of antioxidant activity and total phenolic content of dilliena indica leaves extracts obtained using various technique. Proc. 14th Regional Symposium on Chemical Engineering. Chemical Engineering Depart.Gadjah Mada University. Yogyakarta 4-5 Desember 2007. Indonesia
- Ariana, M., A. Samie, M. A. Edriss, and R. Jahanian. 2011. Effects of powder and extract form of green tea and marigold, and α -tocopheryl acetate on performance, egg quality and egg yolk cholesterol levels of laying hens in late phase of production. J. of Med. Plants Res. Vol. 5(13), pp. 2710-2716, 4 July, 2011
- Azar, S.C., H.A.Shahriar, Sis, N.M., A.R. Ahmadzadeh, and T. Vahdatpoor. 2008. Omega -3 fatty acids enrichment and organoleptic characteristics of broiler meat. Asian J. of Anim. and Vet. Adv. 3 (2): 62-69
- Azman, M.A., Übrahim H. .ER.Ü, and Nurg.I BÜRBEN. 2005. Effects of various dietary fat sources on performance and body fatty acid composition of broiler chickens. Turk J Vet. Anim. Sci. (29) 811-819. © TUBITAK
- Bajerskaa, J., M. Wozniewiczza, J. Jeszkaa, S. Drzymala-Czyzb, and J. Walkowiaka.2011. Green tea aqueous extract reduces visceral fat and decreases protein availability in rats fed with a high-fat diet. Nutr. Res. 31 : 157–164
- Baolu, Z. 2003. Antioxidant effect of Green tea Polyphenols. Chinese Sci. Bulletin (48) no. 4315319
- Basu,A., K. Sanchez, M. J. Leyva, ,M. Wu, , N. M. Betts, C. E. Aston, and T. J. Lyons. 2010. Green Tea Supplementaion Affects Body Weight, Lipids and Lipid Peroxidation in Obese Subyects with Metabolic Syndrome. J. of teh Amer College of Nutri. Vol. 29, No. 1, 31–40
- Berges E.1999. Importance of vitamin E in teh oxidative stability of meat: Organoleptic qualities and consequences. In : Brufau J. (ed.), Tacon A. (ed.). Feed manufacturing in teh Mediterranean region: Recent advances in research and technology. Zaragoza : CIHEAM, 1 9 9 9 . p. 347 -363 (Cahiers Options Méditerranéennes; n .37)
- Betti,M., T.I. Perez, M. J. Zuidhof, and R.A.Renema. 2009. Omega-3-enriched broiler meat: 3. Fatty acid distribution between triacylglycerol and phospholipid classes. Poultry Sci. 88 :1740–1754
- Biswas, M.A.H. and M. Wakita, 2001. Effect of dietary japanese green tea powder supplementation on feed utilization and carcass profiles in broilers. J. Poult. Sci., 38: 50-57.

- Bose, M., J.D.Lambert, J. Ju, K.R.Reuhl, S.A. Shapes, and C.S.Yang. 2008. Teh Major Green Tea Polyphenol, (-)-Epigallocatechin-3-Gallate, Inhibits Obesity, Metabolic Syndrome, and Fatty Liver Disease in High-Fat–Fed Mice. *J. Nutr.* 138: 1677–1683,
- Brewer, M.S. 2011. Natural antioxidant : sources, compounds, mechanisms of action and potential application. *Comprehensive Reviews in Food Science and Food Safety* vo. 10 :211-246
- Bursill, C.A., M. Abbey, and P.D. Roach. 2007. A green tea extract lowers plasma cholesterol by inhibiting cholesterol synthesis and upregulating the LDL receptor in the cholesterol-fed rabbit. *Atherosclerosis* 193 : 86–93
- Bou, R., R. Codony, A. Tres, E.A.Decker, and F. Guardiola. 2009. Dietary strategies to improve nutritional value, oxidative stability, and sensory properties of poultry products. *Critical Reviews. Food Sci and Nutr.* 49:800–822
- Clarkson, M.P. and H.S. Thompson. 2000. Antioxidants : What role do they play in physical activity and health ? . *Amer J. Clin Nutr.* 72 (suppl) : 637S-646S
- Chi-Hua Lu and Lucy Sun Hwang. 2008. Polyphenol contents of Pu-Erh teas and their abilities to inhibit cholesterol biosynthesis in Hep G2 cell line. *Food Chem.* 111 (2008) 67–71
- Christaki E. 2012. Naturally derived antioxidant in poultry nutrition. *Research Journal of Biotechnology* Vol. 7 (3) August : 109 - 112
- Cortinas, L., C. Villaverde, J. Galobart, M.D. Baucells, R.Codony, and A.C. Barroeta. 2004. Fatty acid content in chicken thigh and breast as affected by dietary polyunsaturation level. *Poultry Sci.* 83:1155–1164
- Cuboni, J.,P. Hascik, I. Elimam, J. Garlik, M. Kacaniova, and H.A. Mohammed. 2013. The Influence of Bee Pollen on the Meat Chemical Composition for Broiler Ross 308 Muscles. *Journal of Microbiology, Biotech. and Food Sci.* 2 (Special issue on BQRMF) 1128-1137
- Cunningham JG. 2002. *Textbook of veterinary physiology.* USA: Saunders Company.
- Dauqan, E.M.A., A. Abdullah, and H.A.Sani. 2011. Natural antioxidants, lipid profile, lipid peroxidation, antioxidant enzymes of different vegetable oils. *Adv J. of Food Sci. and Tech.* 3(4): 308-316
- Daneshyar, M., M. Alizadeh Ghandkanlo, F.Sabzi Bayeghra, F. Farhangpajhoh, and M. Aghaei. 2011. Effects of dietary turmeric supplementation on plasma lipoproteins, meat quality and fatty acid composition in broilers. *S. Afr. J. Anim. Sci.* 41 (no. 4)

- D'Archivio, M., C. Filesi, R.D. Benedetto, R. Gargiulo, C. Giovannini, and R. Masella. 2007. Polyphenols, dietary sources and bioavailability. *Ann Ist Super Sanità* Vol. 43, No. 4: 348-361
- D'Mello, J.P.F. 2000. *Farm Metabolism and Nutrition*. CABI Publishing Cambridge. USA
- Dufresne, C.J. and E.R. Farnworth. 2001. A review of latest research findings on teh health promotion properties of tea. *J. Nutr. Biochem.* 12 (2001): 404-421
- Dullo, G.A., C. Duret, D. Rohrer, L. Girardier, N. Mensi, M. Fathi, P. Chantre, and J. Vandermander. 1999. Efficacy of a green tea extract rich in catechin polyphenols and caffeine in increasing 24-h energy expenditure and fat oxidation in humans. *Am J Clin Nutr* ;70:1040–5.
- El-Damrawy, S.Z. 2014. Effect of Grape seed extract on some physiological chane in broiler under heat stress. *Egypt. Poultr. Sci.* Vol (34) (I): (333-343)
- Erol, N.T., F.Sari, G. Polat, and Y.S.Velioglu. 2009. Antioxidant and antibacterial activities of various extracts and fractions of fresh tea leaves and green tea. *Ankara Universitesi Ziraat Fakultesi. Tarim Bilimleri Dergisi*, 15 (4) 371-378
- Esterbauer, H. 1993. Cytotoxicity and genotoxicity of lipid-oxidation product. *Teh Amer. J. Clin. Nutr.* 57 (suppl.) : 779S-786S
- Fellenberg, M.A., and H. Speisky. 2006. Antioxidants : Tehir effect on broiler oxidative stress and its meat oxidative stability. *World Poultry Sci. J.* 62 : 53-70
- Fki, I., M. Bouaziz, Z. Sahnoun, and S. Sayadi. 2005. Hypocholesterolemic effects of phenolic-rich extracts of chemlali olive cultivar in rats fed a cholesterol-rich diet. *Bioorg. & Med Chem.* 13: 5362–5370.
- Flis, M., W. Sobotka, Z. Antoszkiewicz, K. Lipiński, and Z. Zduńczyk. 2010. Teh effect of grain polyphenols and teh addition of vitamin E to diets enriched with α -linolenic acid on teh antioxidant status of pigs. *.lr. Anim. and Feed Sci.* 19, 539–553
- Frei, B. and J.V. Higdon. 2003, Antioxidant activity of tea polifenols In Vivo: Evidence from animal studies. *Teh J. Nutr.* 133 (10) : 3275S – 3284S
- Gladine, C., C. Morand, E. Rock, D. Bauchart, and D. Durand. 2007. Plant extracts rich in polyphenols (PERP) are efficient antioxidants to prevent lipoperoxidation in plasma lipids from animals fed n-3 PUFA supplemented diets. *Anim. Feed Sci and Tech.* 136 : 281–296

- Gladine, C., C. Morand, E. Rock, D. Gruffat, D. Bauchart, and D. Durand. 2007. Teh antioxidative effect of plant extracts rich in polyphenols differs between liver and muscle tissues in rats fed n-3 PUFA rich diets. *Anim. Feed Sci. and Tech.* 139 : 257–27
- Gobert, M., B. Martin, A. Ferlay , Y. Chilliard , B. Graulet , P. Pradel , D. Bauchart, and D. Durand. 2009. Plant polyphenols associated with vitamin E can reduce plasma lipoperoxidation in dairy cows given n-3 polyunsaturated fatty acids. *American Dairy Science Association*, 2009 . *J. Dairy Sci.* 92 :6095–6104.
- Gomikawa, S., Y. Ishikawa, W. Hayase, Y. Haratake, N. Hirano, H. Matuura, A. Mizowaki. A. Murakami, and M. Yamamoto. 2008. Effect of ground green tea drinking for 2 weeks on teh susceptibility of plasma and ldl to teh oxidation ex vivo in healthy volunteers. *Kobe J. Med. Sci.*, Vol. 54, No. 1, pp. E62-E72
- Gropper, S.S., J.L. Smith, and J.L. Groff. 2005. *Advanced nutrition and human metabolism*. Fourth edition. USA: Wardsworth
- Grau, A., F. Guardiola, S. Grimpa, A.C. Barroeta and R. Codony. (2001). Oxidative stability of dark chicken meat through frozen storage: influence of dietary fat and α -tocopherol and ascorbic acid supplementation. *Poultry Sci.* 80:1630–1642
- Guyton A.C and Hall J.E. 1997. *Buku ajar fisiologi kedokteran*. Edisi ke-9. Diterjemahkan oleh Irawati Setiawan. EGC. Jakarta
- Haidari, F., K. Omidian, H. Rafei, M. Zarei, and M.M. Shahi. 2013. Green tea (*camellia sinensis*) supplementation to diabetic rats improves serum and hepatic oxidative stress markers. *I. J. Pharm. Res.* 12 (1): 109-114.
- Handa, S.S., S.P.S.Kanuja, G. Longo and D.D.Rakesh. 2008. *Extraction technology for medicinal and aromatic plant*. isc-unido international center for science and high technology. Trieste, Italy
- Harborne, J.B, 1987, *Metode fitokimia penuntun cara modern menganalisa tumbuhan*, a.b. : Kosasih Padmawinata, ITB, Bandung
- Hoffbrand A.V, JE Pettit. 1996. *Kapita selekta hematologi*. Ed ke-2. Iyan D, penerjemah. Jakarta : Penerbit Buku Kedokteran, EGC. Terjemahan dari : *Essential Hematology*.
- Hsu, Y.W., C.F. Tsai, W.K. Chen, C. F. Huang and C.C. Yen. 2011. A subacute toxicity evaluation of green tea (*Camellia sinensis*) extract in mice. *Food and Chem. Toxic.* 49 : 2624–2630

- Ikeda, I., Y. Imasato and E. Sasaki. 1992. Tea catechins decrease micellar solubility and intestinal absorption of cholesterol in rats. *Biochim. Biophys. Acta.* 1127:141–146.
- Ikuro Abe.2002.Inhibition of cholesterol biogenesis by tea polyphenols. *foods food ingredients J. Japan* No. 204. Abstrak
- Jain, N. C. 1993. *Essentials of veterinary haematology.* Lea and Fabiger, Philadelphia.
- Jawad, A.A.H. 2007. Some hematological and biochemical effect of garlic on broiler chicken. *Bas.J.Vet.Res.Vol.6,No.2* : 56-63
- Jeong, J.Y., G.D. Kim, E.Y. Jung, Y.H. Hwang, S.H. Kim, K.H. Kang Y.H. Choi, S.T. Joo, and G.B. Park. 2009. Effects of dietary green tea on meat quality and physico-chemical properties of broiler during cold storage. *Korean J. Poult. Sci.* Vol. 36, No.1, 77~84
- John,K.M.M., D. Vijayan, R.R.Kumar and R.Premkumar. 2006. Factors influencing teh efficiency of extraction of polyphenols from young tea leaves. *asian journal of plant sci.* 5(1): 123-126
- Kahkonen, M.P. Hopia, A.I. dan Heinonen,(2001),“Berry phenolics and teh antioxidant activity”,*Journal of Agricultural and Food Chemistry*,49, hal4076-4082
- Karaca, T., F. Bayiroglu, M. Cemek, B. Comba, A. Ayaz, and I. Karaboga. 2013. . Effects of green tea extract and lactobacillus casei strain shirota on levels of serum minerals, cholesterol, triglycerides, glucose and lactate in rats fed on high-carbohydrate and high-lipid diets. *Kafkas J Med Sci* 2013; 3(1):1–7
- Karre, E.A. 2009. An overview of some natural antioxidants used in meat and poultry products. *Tehsis. Master of Science.* Kansas State University. Manhattan. Kansas. <https://krex.k-state.edu/dspace/handle/2097/1508>
- Khan, R.U., S. Naz, Z. Nikousefat, V. Tufarelli, M. Javdani, M.S. Qureshi, and V. Laudadio. 2012. Potensial application of ginger (*zingiber officinale*) in poultry diets. *World's Poultry Sci. J.* Vol. 68 : 245-25
- Kojima, S. And Y. Yoshida. 2008. Effects of green tea powder feed supplement on performance of hens in teh late stage of laying. *international J. Poultry Sci.* 7 (5): 491-496
- Koo, M.W.L. and C.H. Cho. 2004. Pharmacological effects of green tea on teh gastrointestinal system. *Europ. J. of Pharm.* Volume 500, Issues 1–3,,Pages177-185

- Koo, S.I. and S K. Noh. 2007. Green Tea as Inhibitor of Intestinal Absorption of Lipids: Potential Mechanism for its Lipid-Lowering Effect. *J Nutr Biochem.* 18(3): 179–183
- Kubo, I., N. Masuoka, P. Xiao, and H. Haraguchi. 2002. Antioxidant activity of dodecyl gallate. *J. Agric. Food Chem.* 50 : 3533-3539.
- Lu, C.H., and L.S. Hwang. 2008. Polyphenol contents of Pu-Erh teas and their abilities to inhibit cholesterol biosynthesis in Hep G2 cell line. *Food Chem.* 111 : 67–71
- Markham, K.R., 1982. *Technique of Flavonoid Identification*, diterjemahkan oleh Padmawinata, K. 1988, Penerbit ITB, Bandung
- Maroon, D.J., G.P.Lu, N.S. Cai, Z.G.Wu, Y.H. Li, H. Chen, J.Q.Zhu, X.J.Jin, B.C.Wouter, and J. Zhao. 2003. Cholesterol lowering effect of a theaflavin-enriched green tea extract. A Randomized Controlled Trial. *Arch Intern Med* 163 : 1448-1453
- Meyer D J and Harvey JW. 2004. *Veterinary Laboratory Medicine Interpretation & Diagnosis*. Third edition. USA: Saunders.
- Min, B. and D.U. Ahn. 2005. Mechanism of lipid peroxidation in meat and meat products -a review. *Food Sci. Biotechn.* Vol. 14, No. 1, pp. 152 – 163.
- Mirshekar, R., B. Dastar and B. Shabanpour. 2009. Effect of rosemary, echinacea, green tea extract and ascorbic acid on broiler meat quality. *Pakistan Journal of Biological Sciences* 12 (15): 1069-1074
- Mitruka, H.M. and S.K. Rawnsley. 1997. *Chemical, biochemical and haematological reference in moral experimental animal*. Mason, N.Y. PP. 287- 380.
- Morrisey, P.A., S.Brandon, D.J. Buckley, P.J.A. Sheehy, and M. Friegg. 1997. Tissue content of alpha tocopheryl acetate supplement for various periods pre-slaughter. *B. Poultry Sci.* 38 : 84-88
- Mukhtar, H and N.I Ahmad. 2000. Tea polyphenols: prevention of cancer and optimizing health. *Am J Clin Nutr*;71(suppl):1698S–1702S., Downloaded from www.ajcn.org by on April 16, 2010
- Ngamukote, S., K. Mäkynen, T. Thilawech and S. Adisakwattana. 2011. Cholesterol-lowering activity of tea major polyphenols in grape seed. *molecules* 16, 5054-5061
- Nagao, T., Y. Komine, S.Soga, S. Meguro, T. Hase, Y. Tanaka, and I. Tokimitsu. 2005. Ingestion of a tea rich in catechins leads to a reduction in body fat and malondialdehyde-modified LDL in men. *Am J Clin Nutr*; 81:122–129

- Nakagawa, T., T. Yokozawa, K. S. Terazawa, K., S. Shu, and L. R. Juneja, L. R., 2002. Protective activity of green tea against free radical and glukosemediated protein damage. *J. Agric. Food Chem.* 50 : 2418-2422
- Nelson, M and J. Poulter. 2004. Impact of tea drinking on iron status in teh UK: a review. *Teh British Dietetic Association Ltd 2004.* *J. Hum. Nutr. Dietet.* 17, pp. 43–54
- Nobar, R.S.D., K. Nazeradi, A. Gorbani, H. Aghdamshahriar and J. Gheyasi. 2007. Incorporation of DHA and EPA fatty acid in to broiler meat lipids. *J. Anim and Vet Adv* 6 (10) : 1199 – 1203.
- Osada, K., M. Funayama, S. Fuchi, S. Sami, Y. Ohta, T. Kanda, and M. Ikeda. 2006. Effect of dietary procyanidin and polyphenols on adipose tissue mass and fatty acid metabolism in rats on a higt fat diet . *J. Oleo Sci.* vol 55, no 2 , 79-89
- Orhan, F and M. Eren. 2011. Effect of herbal mixture supplementation to fish oiled layer diets on lipid oxidation of egg yolk, hen performance and egg quality. *Ankara Univ. Vet Fak Derg* 58, 33-39
- Owen, O.J. and A.O. Amakiri. 2011. Serological and haematological profile of broiler finishers' fed graded levels of Bitter Leaf (*Vernonia amygdalina*) Meal. *Adv. Agric. Biotech.* 1 : 77-81. *Journal homepage: www.woaj.org/AAB.*
- Saleh, H., S. Rahimi, M.A.K. Torshizi, and A. Golian. 2010. Effect of dietary fish oil on oxidative stability and lipid composition of broiler chickens breast and thigh meat. *J. Anim. and Vet. Adv.* 9 (22) : 2877 – 2882.
- Samman, S., B. Sandström, M.B. Toft, K. Bukhave, M. Jensen, S.S. Sorensen, and M. Hansen. 2001. Green tea or rosemary extract added to foods reduces nonheme-iron absorption. *Am J Clin Nutr* (73) :607–12.
- Singh, N., P. Rani, M. Gupta, and N. Tandan. 2013. Role of green tea on cadmium toxicity on haematological profile of albino rats. *amer. phytomed. and clin. Tehrapeutics.* (1)(5): 537-542.
- Singh, R.P., K.N.C. Murthy, and G.K. Jayaprakhasa. 2002. Studies on antioxidant activity of Pomegranate (*Punicum granatum*) peel and seed extracts using in vitro models. *J. Agric Food Chem.* 50 : 81-86
- Sahin, K., C. Orhan, M. Tuzcu, S. Ali, N. Sahin, and A. Hayirli. 2010. Epigallocatechin-3-gallate prevents lipid peroxidation and enhances antioxidant defense system via modulating hepatic nuclear transcription factors in heat-stressed quails. *Poultry Sci.* 89 :2251–2258

- Sarker, M.S.K., K.Seok-Yong, K. Gwi-Man, and Y. Chul-Ju. 2010. Effects of camellia *sinensis* and mixed probiotics on teh growth performance and body composition in broiler. J. Med. Plants Res. Vol. 4(7), pp. 546-550
- Smet, K., K. Raes, G. Huyghebaert, L. Haak,S. Arnouts, and S. De Smet. 2008. Lipid and protein oxidation of broiler meat as influenced by dietary natural antioxidant supplementation. Poultry Sci. 87:1682–1688
- Sohaib, M., F.M.Anjum, M.I.Khan, M.S. Arshad, and M. Shahid. 2012. Enhancement of lipid stability of broiler breast meat and meat products fed on alpha lipoic acid and alpha tocopherol acetate supplemented feed. Lipid in Health and Disease 11 : 57. <http://www.lipidworld.com/content/11/1/57>
- Souza XR, PB Faria, and MC Bressan. 2011. Proximate composition and meat quality of broilers reared under different production systems . Braz. J. Poultry Sci. Jan - Mar 2011 / v.13 / n.1 / 15-20
- Sudibya, 2003. Penggunaan kulit kedelai terhidrolisis dan minyak ikan lemuru dalam ransum terhadap kadar asam lemak Omega-3, Omega-6 dan kolesterol telur itik. Majalah Ilmiah UNSOED no. 1 th. XXIX. Universitas Djendral Soedirman. Purwokerto.
- Swenson .1984. Duke's Physiology of Domestic Animals. Tenth edition. London: Cornel university Press.
- Tan, C. Y., R. Z. Zhong, Z. L. Tan, X. F. Han, S. X. Tang, W. J. Xiao, Z. H. Sun, and M. Wang. 2011. Dietary inclusion of tea catechins changes fatty acid composition of muscle in goats. Lipids 46:239–247. <http://dx.doi.org/10.1007/s11745-010-3477-1>.
- Tang, S. Z., J. P. Kerry, D. Sheehan, D. J. Buckley, and P. A. Morrissey. 2000. Dietary tea catechins and iron-induced lipid oxidation in chicken meat, liver and heart. Meat Sci. 56 : 285–290.
- Tang, S.Z., J.P. Kerry, D. Sheehan, and D.J. Buckley. 2002. Antioxidative mechanisms of tea catechins in chicken meat systems. Food Chem., 76: 45-51.
- Tarladgis, B. G., B. M. Watts, and M. T. Younathan. 1960. A distillation method for teh quantitative determination of malonaldehyde in rancid foods. J. Am. Oil Chem. Soc. 37:44–48.
- Toghyani, M., M. Tohidi, A. A. Gheisari, and S.A. Tabeidian. 2010. Performance, immunity, serum biochemical and hematological parameter in broiler chicks fed dietary thyme as alternative for an antibiotic growth promotor. Afr. J. Biotech. Vol. 9(40), pp. 6819-6825.

- Terry, P.D., J.B. Terry, and T.E. Rohan. 2004, Long-Chain (n-3) Fatty Acid Intake and Risk of Cancers of teh Breast and teh Prostate. *Teh J. Nutr.* 134: 3412S–3420S.
- Turkmen, N., Y.S.Velioglu, F. Sari and G. Polat. 2007. Effect of extraction condition on measured total polyphenol contents and antioxidant and antibacterial activities of black tea. *Molecules* 12 : 484-496
- Uganbayar, D., I. S. Shin, and C. J. Yang. 2006. Comparative performance of hens fed diets containing korean, japanese and chinese green tea. *Asian-Aust. J. Anim. Sci.* Vol. 19, No. 8 : 1190 – 1196
- Uganbayar, D. (2004). A study on teh utilization of green tea for laying hens and broiler chicks. A Dissertation for teh degree of doctor of philosophy, Sunchon National University. Sunchon.
- Velasco, V. and P. Williams. 2011. Review. Improving meat quality through natural antioxidants. *Chil. J. Agric. Res.* 71 (2) : 313-322
- Wijeratne, S.S.K., S.L. Cuppett, and V. Schlegel. 2005. Hidrogen peroxide induced oxidation stress damage and oxidant enzyme response in Caco-2 human colon cell. *J. Agric Food Chem.* 53: 8768-8774
- Winarsi, H. 2007. *Antioksidan Alami dan Radikal Bebas. Potensi dan Aplikasinya dalam Kesehatan.* Penerbit Kanisius. Yogyakarta
- Vossen, E., M. Ntawubizi, K. Raes, K. Smet, G. Huyghebaert, S. Arnouts, and S. De Smet. 2011. Effect of dietary antioxidant supplementation on teh oxidative status of plasma in broilers. *J. Anim. Physiol. and Anim. Nutr.* 95 : 198–205
- Yang, C.J., I.Y. Yang, D.H. Oh, I.H. Bae, S.G. Cho, I.G. Kong, D. Uganbayar, and K.S. Choi. 2003. Effect of green tea by-product on performance and body composition in broiler chicks. *Asian-Aust. J. Anim. Sci.* (16)6: 867-872.
- Yosef, T.A., M.Z. Al-Julaifi, and Kandeel. 2012. Teh Effects of Green Tea (*Camellia Sinensis*) Probiotics on Broilers Exposed to Lead-Induced Oxidative Stress. *J. Amer. Sci.* 8 (4) :628-635
<http://www.americanscience.org>
- Zaveri, N.T., 2006. Green tea and its polyphenolic catechins : medicinal uses in cancer and noncancer applications. *Life Sci.* (78): 2073–2080
- Zuprizal, C.T.Noviandi, Indratiningsih, T. Yuwanta, S. Harimurti. 2002. Teh addition effect of sardine oil in quail ration on omega-3 fatty and composition of eggs. Faculty of Animal Science, Gadjah Mada University. Yogyakarta