



DAFTAR PUSTAKA

- Ahmed, K. R., Karim, M. N., Bukht, M. S., Bhowmik, B., Acharyya, A., Ali, L., & Hussain, A. (2011). Risk factors of diabetic retinopathy in Bangladeshi type 2 diabetic patients. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 5(4), 196–200.
<http://doi.org/10.1016/j.dsx.2012.02.020>
- American Academy of Ophthalmology. (2010). International Clinical Diabetic Retinopathy Disease Severity Scale.
- American Diabetes Association. (2004). Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care*, 27(Supplement 1), 55–60.
<http://doi.org/10.2337/diacare.27.2007.S5>
- Deshpande, A. D., Harris-Hayes, M., & Shootman, M. (2008). Epidemiology of diabetes and diabetes-related complications. *Physical Therapy*, 88, 1254–1264. <http://doi.org/10.2522/ptj.20080020>
- Kaku, K. (2010). Pathophysiology of Type 2 Diabetes and Its Treatment Policy. *Journal of the Japan Medical Association*, 53(1), 41–46.



Ozouguw, O. (2013). The pathogenesis and pathophysiology of type 1 and type 2 diabetes mellitus. *Journal of Physiology and Pathophysiology*, 4(4), 46–57. <http://doi.org/10.5897/JPAP2013.0001>

Powers, A. C. (2011). Diabetes Mellitus. In D. L. Longo, D. L. Kasper, J. L. Jameson, A. S. Fauci, S. L. Hauser, & J. Loscalzo (Eds.), *Harrison's Principles of Internal Medicine*, 18th Edition (pp. 2968–3002). United States: The McGraw-Hill Companies.

Soewondo, P., Soegondo, S., Suastika, K., Pranoto, A., & Soeatmadji, D. W. (2010). The DiabCare Asia 2008 study – Outcomes on control and complications of type 2 diabetic patients in Indonesia. *Med J Indones*, 19(4), 235–244.

Tarr, J. M., Kaul, K., Chopra, M., Kohner, E. M., & Chibber, R. (2013). Pathophysiology of Diabetic Retinopathy. *ISRN Ophthalmology*, 2013, 1–13. <http://doi.org/10.1155/2013/343560>

Tikellis, G., Anuradha, S., Klein, R., & Wong, T. Y. (2011). Association between Physical Activity and Retinal Microvascular Signs: The Atherosclerosis



Risk in Communities (ARIC) Study. *Microcirculation*, 17(5), 381–393. <http://doi.org/10.1111/j.1549-8719.2010.00033.x>. Association

Varis, E., & J. Savolainen, M. (2012). Alcohol Consumption, but Not Physical Activity or Smoking, Contribute To Advanced Forms of Diabetic Retinopathy: A Case-Control Study. *Journal of Clinical & Experimental Ophthalmology*, 01. <http://doi.org/10.4172/2155-9570.S5-005>

Whiting, D. R., Guariguata, L., Weil, C., & Shaw, J. (2011). IDF Diabetes Atlas: Global estimates of the prevalence of diabetes for 2011 and 2030. *Diabetes Research and Clinical Practice*, 94(3), 311–321. <http://doi.org/10.1016/j.diabres.2011.10.029>

World Health Organization. (2006). Definition and Diagnosis of Diabetes Mellitus and Intermediate Hyperglycemia. *Who2*, 50. <http://doi.org/ISBN 92 4 159493 4>

World Health Organization/Fao. (2004). Human energy requirements. *Agriculture and Consumer Protection*, 0, 107. Retrieved from <http://www.fao.org/docrep/007/y5686e/y5686e08.htm>



Hubungan Aktivitas Fisik dengan Kejadian Retinopati Diabetika pada Pasien Diabetes Mellitus Tipe 2:

Jogjakarta Eye Diabetic Study in the Community (JOGED.COM)

BUNGA BELLA ANNISA, Prof. dr. Suhardjo, SU, Sp.M(K); dr. M. Bayu Sasongko, M.Epi., Ph.D.

Universitas Gadjah Mada, 2015 | Diunduh dari <http://etd.repository.ugm.ac.id/>

UNIVERSITAS
GADJAH MADA

World Health Organization. (2005). Prevention of

blindness from diabetes mellitus. Geneva: WHO, 1-48.

<http://doi.org/ISBN9789242547123>