

## References

- Ahmed, N., Wahlgren, N.G., High Initial Blood Pressure After Acute Stroke is Associated With Poor Functional Outcome. *Journal of Internal Medicine*. 2001;249:467-473.
- Boli, P., 2006. Hipertensi Tidak Terkontrol Sebagai Faktor Resiko Utama Terjadinya Stroke Berulang. *Disertation*. Fakultas Kedokteran Universitas Gadjah Mada.
- Carlberg, B., Asplund, K., Hägg, E., The Prognostic Value of Admission Blood Pressure in Patients with Acute Stroke. *Stroke*. 1993;24:1372-1375.
- Caskie, G.I., Suttton, M.C., Margaret, J.A., The Relation of Hypertension to Changes in ADL/IADL Limitations of Mexican American Older Adults. *J Gerontol B Psychol Sci Soc Sci*. 2010; 65B(3):296-305.
- Chindaprasirt, J., Sawanyawisuth, K., Chattakul, P., Limpawattana, P., Age Predicts Functional Outcome in Acute Stroke Patients with rt-PA Treatment. *ISRN Neurology*. 2013;
- Doehner, W., Schenkel, J., Anker, S.D., Springer, J., Audebert, H.J., Overweight and Obesity are Associated with Improved Survival, Functional Outcome, and Stroke Recurrence After Acute Stroke or Transient Ischaemic Attack: Observations from The TEMPis Trial. *European Heart Journal*. 2012; 2013(34):268-277.
- Goldstein, L., 2009. *A Primer on Stroke Prevention Treatment: An Overview Based on AHA / ASA Guidelines*. 1st ed. Durham North Carolina USA: Wiley Blackwell.
- Goldstein, L., Amarenco, P., Zivin, J., Messig, M., Alfatullah, I., Callahan, A., Hennerici, M., Statin Treatment and Stroke Outcome in The Stroke Prevention by Aggressive Reduction in Cholesterol Levels (SPARCL) Trial. *Stroke*. 2009;40(11):3256-31.
- Lamsudin, R., Pengendalian Hipertensi Sebagai Faktor Risiko Stroke Dan Manajemen Hipertensi Pada Penderita Stroke Akut. *Berkala NeuroSains*. 2000; 1(3):127-31.
- Markus, H.S., *Stroke: Causes and Clinical Features*. The Medicine Publishing Company Ltd. 2004:57-61.

- Megherbi, S.E., Milan, C., Minier, D., Couvreur, G., Osseby, G.V., Tilling K., Carlo, A., C., Association Between Diabetes and Stroke Subtype on Survival and Functional Outcome 3 Months After Stroke : Data From the European BIOMED Stroke Project. *Stroke*. 2003;34:688-694.
- Nakayama, H., Jørgensen, H.S., Raaschou, H.O., Olsen, T.S., The Influence of Age on Stroke Outcome. *The Copenhagen Stroke Study*. 1994;25:808-813.
- Obviegale, B., Weir, C.J., Saver, J.L., Muir, K.W., Lees, K.R., Effect of Smoking Status on Outcome After Ischemic Stroke. *Cerebrovascular Dis*. 2006;21(4):260-5.
- Petrea, R.E., Beiser, A.S., Shadri, S., Kelly-Hayes, M., Kase, M.S., Wolf, P.A., Gender Differences in Stroke Incidence and Poststroke Disability in the Framingham Heart Study. *Stroke*. 2009;40:1032-1037.
- Sacco, R., 2001. *Stroke Prevention*. Edited by John Norris and Vladimir Hachinski. 1st ed. London: Oxford University Press.
- Schretzman, D., Acute Ischemic Stroke. *Dimensions of Critical Care Nursing*. 2001;20(2).
- Setyopranoto, I., 2012. *Odem Otak pada Pasien Stroke Iskemik Akut*, Yogyakarta: Badan Penerbit Fakultas Kedokteran UGM.
- Setyopranoto, I., Stroke: Gejala Dan Penatalaksanaan. *CDK 185*. 2011;38(4): 247-50.
- Supraptiningsih, 1992. Reliabilitas Modifikasi Indeks Barthel Pada Penderita Stroke. *Disertation*. Fakultas Kedokteran Universitas Gadjah Mada.
- Supraptiningsih, Lamsudin, R., Was'an, M., Sutanto, Reliabilitas Modifikasi Indeks Barthel Pada Penderita Stroke. *Berkala NeuroSains*. 2012;3(2): 1-10.