

## 6. References

- Aprilia, A., Tezuka, T., Spaargaren, G., 2012. Household Solid Waste Management in Jakarta , Indonesia : A Socio-Economic Evaluation. *Waste Manag.* 70–95. doi:<http://dx.doi.org/10.5772/51464>
- Belostotskiy, D.E., Ziganshina, E.E., Siniagina, M., Boulygina, E. a., Miluykov, V. a., Ziganshin, A.M., 2015. Impact of the substrate loading regime and phosphoric acid supplementation on performance of biogas reactors and microbial community dynamics during anaerobic digestion of chicken wastes. *Bioresour. Technol.* 193, 42–52. doi:10.1016/j.biortech.2015.06.066
- Blackwood, C.B., Marsh, T., Kim, S., Paul, E.A., 2003. Terminal Restriction Fragment Length Polymorphism Data Analysis for Quantitative Comparison of Microbial Communities Terminal Restriction Fragment Length Polymorphism Data Analysis for Quantitative Comparison of Microbial Communities. doi:10.1128/AEM.69.2.926
- Borneman, J., Triplett, E.W., 1997. Molecular microbial diversity in soils from eastern Amazonia: evidence for unusual microorganisms and microbial population shifts associated with deforestation. *Appl. Environ. Microbiol.* 63, 2647–53.
- Cahyari, K., Putra, R.A., 2010. Design of Biogas Plant from Fruit Market Waste in Indonesia. *Renew. Energy Conf.*
- Code, C., Fac, B., Science, F., 2006. THE FIRST DECADE OF TERMINAL RESTRICTION FRAGMENT LENGTH POLYMORPHISM (T-RFLP) IN MICROBIAL ECOLOGY *Blaž STRES* 88, 65–73.
- Díez, B., Pedrós-Alió, C., Marsh, T.L., Massana, R., 2001. Application of denaturing gradient gel electrophoresis (DGGE) to study the diversity of marine picoeukaryotic assemblages and comparison of DGGE with other molecular techniques. *Appl. Environ. Microbiol.* 67, 2942–51. doi:10.1128/AEM.67.7.2942-2951.2001
- Freguia, S., Teh, E.H., Boon, N., Leung, K.M., Keller, J., Rabaey, K., 2010. Microbial fuel cells operating on mixed fatty acids. *Bioresour. Technol.* 101, 1233–1238. doi:10.1016/j.biortech.2009.09.054



- Hafez, H., Nakhla, G., El. Naggar, M.H., Elbeshbishy, E., Baghchehsaraee, B., 2010. Effect of organic loading on a novel hydrogen bioreactor. *Int. J. Hydrogen Energy* 35, 81–92. doi:10.1016/j.ijhydene.2009.10.051
- Hughes, A.D., Kelly, M.S., Black, K.D., Stanley, M.S., 2012. Biogas from Macroalgae: is it time to revisit the idea? *Biotechnol. Biofuels* 5, 86. doi:10.1186/1754-6834-5-86
- Jang, S., Kim, D.-H., Yun, Y.-M., Lee, M.-K., Moon, C., Kang, W.-S., Kwak, S.-S., Kim, M.-S., 2015. Hydrogen fermentation of food waste by alkali-shock pretreatment: Microbial community analysis and limitation of continuous operation. *Bioresour. Technol.* 186, 215–222. doi:10.1016/j.biortech.2015.03.031
- Kim, D.H., Kim, S.H., Shin, H.S., 2009. Hydrogen fermentation of food waste without inoculum addition. *Enzyme Microb. Technol.* 45, 181–187. doi:10.1016/j.enzmictec.2009.06.013
- Kirk, J.L., Beaudette, L. a, Hart, M., Moutoglis, P., Klironomos, J.N., Lee, H., Trevors, J.T., 2004. Methods of studying soil microbial diversity. *J. Microbiol. Methods* 58, 169–88. doi:10.1016/j.mimet.2004.04.006
- Levin, D.B., Pitt, L., Love, M., 2004. Biohydrogen production : prospects and limitations to practical application 29, 173–185. doi:10.1016/S0360-3199(03)00094-6
- Lin, X., Xia, Y., Yan, Q., Shen, W., Zhao, M., 2014. Acid tolerance response (ATR) of microbial communities during the enhanced biohydrogen process via cascade acid stress. *Bioresour. Technol.* 155, 98–103. doi:10.1016/j.biortech.2013.12.097
- Nielson, D.L., 2004. Aligning & analyzing TRFLP data 1–4.
- Nissilä, M.E., Tähti, H.P., Rintala, J. a., Puhakka, J. a., 2011. Effects of heat treatment on hydrogen production potential and microbial community of thermophilic compost enrichment cultures. *Bioresour. Technol.* 102, 4501–4506. doi:10.1016/j.biortech.2010.12.072
- Nitipan, S., Mamimin, C., Intrasungka, N., Birkeland, N.K., O-Thong, S., 2014. Microbial community analysis of thermophilic mixed culture sludge for biohydrogen production from palm oil mill effluent. *Int. J. Hydrogen Energy* 9. doi:10.1016/j.ijhydene.2014.05.139

- Oh, S.-E., Iyer, P., Bruns, M.A., Logan, B.E., 2004. Biological hydrogen production using a membrane bioreactor. *Biotechnol. Bioeng.* 87, 119–27. doi:10.1002/bit.20127
- Park, J.H., Lee, S.H., Yoon, J.J., Kim, S.H., Park, H.D., 2014. Predominance of cluster I Clostridium in hydrogen fermentation of galactose seeded with various heat-treated anaerobic sludges. *Bioresour. Technol.* 157, 98–106. doi:10.1016/j.biortech.2014.01.081
- Ren, N., Guo, W., Wang, X., Xiang, W., Liu, B., Wang, X., Ding, J., Chen, Z., 2008. Effects of different pretreatment methods on fermentation types and dominant bacteria for hydrogen production 33, 4318–4324. doi:10.1016/j.ijhydene.2008.06.003
- Sciences, A., 2010. Technical Report on the PCR-DGGE Analysis of Soil Nematode Community Table of Contents.
- Shih, J.C.W.C.S., Wen, S.Y.J.L.F., 2006. Molecular detection of the clostridia in an anaerobic biohydrogen fermentation system by hydrogenase mRNA-targeted reverse 598–604. doi:10.1007/s00253-005-0106-7
- Sikora, A., Błaszczuk, M., Jurkowski, M., Zielenkiewicz, U., 2013. Lactic Acid Bacteria in Hydrogen-Producing Consortia : On Purpose or by Coincidence ? Lactic Acid Bacteria - R & D for Food, Health and Livestock Purposes, Dr. J. Marcelino Kongo (Ed.), ISBN: 978-953-51-0955-6, InTech, Available from: <http://www.intechopen.com>.
- Sivagurunathan, P., Sen, B., Lin, C.Y., 2014. Batch fermentative hydrogen production by enriched mixed culture: Combination strategy and their microbial composition. *J. Biosci. Bioeng.* 117, 222–228. doi:10.1016/j.jbiosc.2013.07.015
- Stratford, J.P., Beecroft, N.J., Slade, R.C.T., Grüning, A., Avignone-Rossa, C., 2014. Anodic microbial community diversity as a predictor of the power output of microbial fuel cells. *Bioresour. Technol.* 156, 84–91. doi:10.1016/j.biortech.2014.01.041
- Suzuki, R., Shimodaira, H., 2004. An application of multiscale bootstrap resampling to hierarchical clustering of microarray data: How accurate are these clusters. *Fifteenth Int. Conf. ...* 0–1.
- Tawfik, A., El-Bery, H., Kumari, S., Bux, F., 2014. Use of mixed culture bacteria for photofermentative hydrogen of dark fermentation effluent. *Bioresour. Technol.* 168, 119–126. doi:10.1016/j.biortech.2014.03.065



- Vijayaraghavan, K., Ahmad, D., Soning, C., 2007. Bio-hydrogen generation from mixed fruit peel waste using anaerobic contact filter. *Int. J. Hydrogen Energy* 32, 4754–4760. doi:10.1016/j.ijhydene.2007.07.001
- Weimer, P.J., Nerdahl, M., Brandl, D.J., 2015. Production of medium-chain volatile fatty acids by mixed ruminal microorganisms is enhanced by ethanol in co-culture with *Clostridium kluyveri*. *Bioresour. Technol.* 175, 97–101. doi:10.1016/j.biortech.2014.10.054
- Wiesemüller, B., Rothe, H., 2006. Interpretation of bootstrap values in phylogenetic analysis. *Anthropol. Anz.* 64, 161–5.
- Yang, Z., Xu, X., Guo, R., Fan, X., Zhao, X., 2015. Accelerated methanogenesis from effluents of hydrogen-producing stage in anaerobic digestion by mixed cultures enriched with acetate and nano-sized magnetite particles. *Bioresour. Technol.* 190, 132–139. doi:10.1016/j.biortech.2015.04.057
- Zhi, W., Ge, Z., He, Z., Zhang, H., 2014. Methods for understanding microbial community structures and functions in microbial fuel cells: A review. *Bioresour. Technol.* 171, 461–468. doi:10.1016/j.biortech.2014.08.096