

## DAFTAR PUSTAKA

- Anna, F. & Wibisono, M.J., 2010. Manajemen Sepsis Pada Pneumonia. *Majalah Kedokteran Respirasi*, 1(2), p.52
- Baudouin, S., 2008. 'Sepsis: Introduction and Epidemiology' in *Sepsis*, Springer, London, pp.1-4
- Beth, D.S. & Robert, G.T., 2004. *Basic and Clinical Biostatistics*, 4<sup>th</sup> edn, The Mc-Graw Hill Companies, New York
- Braun, E. et al., 2014. Is elevated red cell distribution width a prognostic prediktor in adult patients with community acquired pneumonia? *BMC Infectious Disease*, 14(1), p.129
- Braun, E. et al., 2011. Elevated red cell distribution width predicts poor outcome in young patients with community acquired pneumonia. *Critical Care*, 15(4), p.R194
- Cengiz, M. et al., 2013. Is increased red cell distribution width an indicating marker of non alcoholic steatohepatitis and fibrotic stage? *World Journal of Gastroenterology*, 19(42), pp.7412-8
- Dahlan, Z., 2009. 'Pneumonia' in *Buku Ajar Ilmu Penyakit Dalam*, eds Sudoyo A.W. et al., Internal Publishing, Jakarta Pusat, pp.2196-2206
- Fandos, S. et al., 2015. Red blood cell distribution width (RDW) and long term mortality after community-acquired pneumonia. A comparison with proadrenomodullin. *Respiratory Medicine Journal*, 109(6), pp.1193-1206
- Goyette, et al., 2004. Hematologic Changes in Sepsis and Their Therapeutic Implications. *Respiratory and Critical Care Medicine*, 25(6), pp.645-658

- Hu, Z. et al., 2013. Red blood cell distribution width is a potential index for liver disease. *Clinical Chemistry and Laboratory Medicine*, 51(7), pp. 1403-8
- Ishiguro, T. et al., 2013. Etiology and Factors Contributing to the Severity and Mortality of Community-acquired Pneumonia. *Internal Medicine*, 54, pp.317-324
- Jo, Y.H. et al., 2013. Red cell distribution width is a prognostic factor in severe sepsis and septic shock. *The American journal of emergency medicine*, 31(3), pp.545-8
- Kee, J.L., 2014. *Laboratory And Diagnostic Test With Nursing Implications*, 10<sup>th</sup> edn, Pearson, New Jersey, pp.371-374
- Kim, C.H. et al., 2013. An Increase in red blood cell distribution width from baseline predicts mortality in patients with severe sepsis or septic shock. *Critical Care*, 17(6), p.R282
- Lam, A.P. et al., 2013. Multiplicative interaction between mean corpuscular volume and red cell distribution width in predicting mortality of elderly patients with and without anemia. *American Journal of Hematology*, 88, pp.245-9
- Lee, J.H. et al., 2013. Red cell distribution width as a prognostic marker in patients with community-acquired pneumonia. *Am J Emerg Med*, 31(1), pp.72-79
- Loebinger, M.R. & Wilso, R., 2009. Bacterial Pneumonia. *Medical Progress*, 39(6), pp.275-279
- Lutfiyya, M.N., Henley, E. & Chang, L.F., 2006. Diagnosis and Treatment of Community-acquired Pneumonia. *Am Fam Physician*, 73(3), pp. 442-450
- Mandel, L.A. & Wunderink, R., 2010. 'Pneumonia' in *Harrison's Pulmonary and Critical Care Medicine*, eds

- Loscalzo, J., The McGraw-Hill Companies, New York, pp.99-108
- Marrie, T.J. et al., 2014. 'Pneumonia' in *Harrison's Principle of Internal Medicine*, 16<sup>th</sup> edn, eds Longo, D.L. & Fauci, A.S., The McGraw-Hill Companies, New York, pp.1528-1540
- Martin, S.W. & Al-Haddad, M., 2013. Community-acquired pneumonia. *Anaesthesia & Intensive Care Medicine*, 14(10), pp.457-459
- Mason, C.M. & Summer, W.R., 2010. 'Resporatory Infections' in *Pulmonary Pathophysiology*, eds Ali, J., Summer, W.R. & Levitzky, M.G., The McGraw-Hill Companies, New York, pp. 166-174
- Miyamoto, K. et al., 2015. Relationships among red cell distribution width, anemia, and interleukin-6 in adult congenital heart disease. *Circulation Journal*, 79, pp. 1100-6
- Munford, R.S., 2004. 'Severe sepsis and septic shock' in *Harrison's Principle of Internal Medicine*, 16<sup>th</sup> edn, eds Longo, D.L. & Fauci, A.S., The McGraw-Hill Companies, New York, pp. 1606-1611
- Niederman, M.S., Feldman, C. & Richards, G. a, 2006. Combining information from prognostic scoring tools for CAP: an American view on how to get the best of all worlds. *The European respiratory journal*, 27(1), pp.9-11
- Oehadian, A., 2012. Pendekatan klinis dan diagnosis anemia. *CDK-194*, 39(6), pp. 407-12
- Pascual-Fiagal, D.A. et al., 2009. Red blood cell distribution width predicts long term outcome regardless of anemia states in acute heart failure patients. *European Journal of Heart Failure*, 11, pp.840-6

- Pereira, J.M. et al., 2013. Can we predict pneumococcal bacteremia in patients with severe community-acquired pneumonia? *Journal of critical care*, 28(6), pp.970-4
- Pereira, J.M., Paiva, J.A. & Rello, J., 2012. Severe sepsis in community-acquired pneumonia--early recognition and treatment. *European journal of internal medicine*, 23(5), pp.412-9
- Straat, M. et al., 2012. Red Blood Cell Clearance in Inflammation. *Transfusion Medicine and Hemotherapy*, 39, pp. 353-360
- WHO., 2011. Haemoglobin concentration for the diagnosis of anemia and assesment of severity. Geneva, *World Health Organization*. Available at <http://www.who.int/vmnis/indicators/haemoglobin.pdf>
- Xu, W.S. et al., 2015. Red blood cell distribution width levels correlate with liver fibrosis and inflammation. *Medicine Journal*, 94(10), pp.1-7