

## DAFTAR PUSTAKA

- Andrew NC. 1999. Disorder of iron metabolism. *N Engl J Med*. 341:1986-1995.
- Anonim<sup>1</sup>. 2012. Reticulocyte haemoglobin content-adding value to anemia diagnostic. Sysmex educational Enhancement and Development.
- Anonim<sup>2</sup>. 2009. Retikulosit dan Ret-He. PT Sysmex.
- Anonim<sup>3</sup>. 2010. Reticulocyte Hemoglobin in Anemia Diagnosis and Management. Sysmex America, Inc.
- Anonim<sup>4</sup>. 2010. Reticulocyte and Their Significance. Sysmex Xtra Online. Available from : URL:  
[http://www.sysmex.ru/files/articles/Xtra\\_online\\_reticulocytes.pdf](http://www.sysmex.ru/files/articles/Xtra_online_reticulocytes.pdf)
- Ayu NP, Suega K, Widiani GR. 2010. Hubungan antara beberapa parameter anemia dan laju filtrasi glomerulus pada penyakit ginjal kronik pradialisis. *J Peny Dalam*. 11(3):140-148.
- Bandiara Ria, 2003. Penatalaksanaan Anemia Defisiensi Besi pada Pasien yang menjalani Hemodialisis. Diajukan pada Workshop “ Registrasi Unit Dialisis se Jawa Barat” PPGI Papandayan Bandung.
- Besarab A., Frinak S., Yee J. 1999. An indistinct balance: the safety and efficacy of parenteral iron therapy. *J Am Soc Nephrol*. 10:2029-2043.
- Brugnara C, Schiller B, Moran J. 2006. Reticulocyte hemoglobin equivalent (Ret He) and assessment of iron-deficient states. *Clin Lab Haematol*. 28:303–8.
- Buttarelo M, et al, 2010. Diagnosis of Iron Deficiency in Patients Undergoing Hemodialysis.
- Chiang, W.C., Tsai, T.J., Chen, Y.M., Lin, S.L., Hsieh, B.S. 2002. Serum soluble transferrin receptor reflects erythropoiesis but not iron availability in erythropoietin-treated chronic hemodialysis patients. *ClinNephrol*58: 363–369.
- Cook JD. 2005 Diagnosis and management of iron deficiency anemia. *Best Pract Res Clin Haematol*. 18(2):319-32.
- Coyne D. 2006. Iron indices: What do they really mean?. *Kidney Int*. 69(S):4-8.
- Dallalio, G., Fleury, T., Means, R.T. 2003. Serum hepcidin in clinical specimens. *Br J Haematol*122: 996–1000.

- Eschbach JW, Abdulhadi MH, Browne JK, et al. 1989. Recombinant human erythropoietin in anemic patients with end-stage renal disease. Results of a phase III multicenter clinical trial. *Ann Intern Med.* 111:992–1000.
- Eschbach JW, Adamson JW. 1999. Iron overload in renal failure patients: changes since the introduction of erythropoietin therapy. *Kidney Int.* 69(suppl 1):S35–S43.
- Eschbach JW, Haley NR, Egrie JC, Adamson JW. 1992. A comparison of the responses to recombinant human erythropoietin in normal uremic subjects. *Kidney Int.* 42:407–416.
- Eschbach JW. 2005. Iron requirements in erythropoietin therapy. *Best Pract Res Clin Haematol.* 18(2):347-361.
- Fishbane S, Pollack S, Feldman HI, Joffe MM. 2009. Iron indices in chronic kidney disease in the National Health and Nutritional Examination Survey 1988–2004. *Clin J Am Soc Nephrol.* 4:57-61.
- Ganz T. 2003. Hepcidin, a key regulator of iron metabolism and mediator of anemia of inflammation. *Blood.* 102:783-788.
- Garzia M, Di Mario A, Ferraro E, Tazza L, Rossi E, Luciani G, Zini G. 2007. Reticulocyte hemoglobin equivalent: an indicator of reduced iron availability in chronic kidney diseases during erythropoietin therapy. *Lab Hematol.* 13(1):6-11.
- Goodnough LT., Skikne B., Brugnara C. 2000. Erythropoietin, iron, and erythropoiesis. *Blood.* 96:823-833.
- Howayda A. Et al., 2012. Iron Deficiency among Anemic Pre-Dialysis Chronic Kidney Disease Patients. Department of Internal Medicine, Faculty of Medicine, Ain Shams University.
- Hsu CY., McCulloch CE., Curhan GC. 2002. Epidemiology of anemia associated with chronic renal insufficiency among adult in the United States: results from the third national health and nutrition examination survey. *J Am Soc Nephrol.* 13:504-510.
- Ifudu O., Feldman J., Friedman EA. 1996. The intensity of hemodialysis and the response to erythropoietin in patients with end-stage renal disease. 334:420-425.

Jonckheere S. et al, 2010. Erythrocyte Indices in the Assessment of Iron status in Dialysis-Dependent patients with end-stage Renal Disease on continuous erythropoietin Receptor activator versus epoetin  $\beta$  Therapy. Departements of clinical Chemistry and Nephrology, Algemeen Ziekenhuis Maria Middelaars, and Departmen of clinical chemistry, Ghent University Hospital, Ghent Belgium.

Kale V., Aftab AR. 2014. Diagnostic evaluation of anemia. Diunduh dari [www.intechopen.com](http://www.intechopen.com) tanggal 9 Desember 2014.

Kaneko Y., Miyazaki S., Hirasawa Y., Gejyo F. 2003. Transferrin saturation versus reticulocyte hemoglobin content for iron deficiency in Japanese hemodialysis patients. *Kidney International*. 63:1086-1093.

Koca E., Cetiner DA., Buyukasik Y., Uner A., Sayinalp N., Haznedaroglu IC., et al. 2013. Bone marrow iron staining is reliable test for elimination of iron deficiency anemia rather than its diagnosis. *International Journal of Hematology and Oncology*. 23: 260-263.

Lederer, E., Ouseph, R. 2007. Chronic kidney disease. Core curriculum in nephrology. *Am J Kidney Disease*. 9:162-171.

Locatelli F, Aljama P, Barany P, et al. 2004. Revised European best practice guidelines for the management of anaemia in patients with chronic renal failure. *Nephrol Dial Transplant*. 19(suppl 2):1-47.

Ly J., Marticorena R., Donnelly S. 2004. Red blood cell survival in chronic renal failure. *Am J Kidney Dis*. 44:715-719.

Ma JZ, Ebben J, Xia H, Collins AJ. 1999. Hematocrit level and associated mortality in hemodialysis patients. *J Am Soc Nephrol*. 10:610-619.

Macdougall IC., Cooper AC. 2002. Erythropoietin resistance: the role of inflammation and pro-inflammatory cytokine. *Nephrol Dial Transplant*. 17:39-43.

Marsh JT, Brown WS, Wolcott D, et al. 1991. rHuEPO treatment improves brain and cognitive function of anemic dialysis patients. *Kidney Int*. 39:155-163.

Mast AE., Blinder M., Dietzen D. 2008. Reticulocyte hemoglobin content. *Am. J. Hematol*. 83:307-310.

Miwa N, Akiba T, Kimata N, Hamaguchi Y, Arakawa Y, Tamura T, et al. 2009. Usefulness of measuring reticulocyte hemoglobin equivalent in the management of haemodialysis patients with iron deficiency. *Int Jnl Lab Hem.* 1-8.

M.Garzia. et al, 2006. Evaluation of the new RET-He parameter versus the CHr reference method in a cohort of 57 dialysed patients in a five month follow-up. International society for laboratory Hematology. Cathollic University of Sacred Heart, Rome- ReCAMH Hematology DPt.

Moreno F, Sanz-Guajardo D, Lopez-Gomez JM, Jofre R, Valderrabano F. 2000. Increasing the hematocrit has a beneficial effect on quality of life and is safe in selected hemodialysis patients. Spanish Cooperative Renal Patients Quality of Life Study Group of the Spanish Society of Nephrology. *J Am Soc Nephrol.* 11:335–342.

National Kidney Foundation. 2002. Clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis.* 39(2 Suppl 1):1-266.

National Kidney Foundation. IV. 2001. NKF-K/DOQI Clinical Practice Guidelines for Anemia of Chronic Kidney Disease: update 2000. *Am J Kidney Dis.* 37(suppl 1):182–238.

Nurko S. 2006. Anemia in chronic kidney disease: causes, diagnosis, treatment. *Cleveland Clinic Journal of Medicine.* 73:289-297.

Pagana K, Pagana T. 2010. Iron level. In: Pagana K, Pagana T, editors. *Mosby's manual of diagnostic and laboratory test.* 4 ed. St Louis: Mosby Elsevier. 336-340.

Perhimpunan Nefrologi Indonesia (Pernefri). 2011. Konsensus manajemen anemia pada penyakit ginjal kronik. Jakarta-Indonesia.

Rafi A., Karkar A., Abdelrahman M. 2007. Monitoring iron status in end stage reanal disease pasients on hemodialysis. *Saudi J Kidney Dis Transpant.* 18(1):73-78.

Reddy C.G, Devaki R and Rao P, 2003. Iron Indices in Patients with functional anemia in Chronic Kidney Disease. Department of Biochemistry, Kamineni Institute of Medical Sciences, Sceepuram, Narketpally,Nalgonda-508 254.

Robertson HT, Haley NR, Guthrie M, Cardenas D, Eschbach JW, Adamson JW. 1990. Recombinant erythropoietin improves exercise capacity in anemic hemodialysis patients. *Am J Kidney Dis.* 15:325–332.

- Schaefer RM, Schaefer L. 1998. Iron monitoring and supplementation: how do we achieve the best results? *Nephrol Dial Transplant*. 13(2):9-12.
- Tarng, D.C., Huang, T.P. 2002. Determinants of circulating soluble transferrin receptor level in chronic haemodialysis patients. *Nephrol Dial Transplant* 17: 1063–1069.
- Tessitore. N., Solero, G.P., Lippi, G., Bassi, A., Faccini, G.B., Bedogna, V., Gammara, L., Brocco, G., Restivo, G., Bernich, P., Lupo, A., Maschio, G. 2001. The role of iron status markers in predicting response to intravenous iron in haemodialysis patients on maintenance erythropoietin. *Nephrol Dial Transplant* 16: 1416–1423.
- Tong EM., Nissenson AR. 2001. Erythropoietin and anemia. *Semin Nephrol*. 21:190-203.
- Ullrich, C., Wu, A., Armsby, C., Rieber S., Wingerter S., Brugnara C., 2005. Healthy Infants for Iron Deficiency Using Reticulocyte Hemoglobin Content. *JAMA* 294 (8): 924–30.
- Urrechaga E, Borque L and Escanero F.J, 2012. Assessing Iron Status in CKD Patients : New Laboratory Parameters. Department of Pharmacology and Physiology, Faculty of Medicine University of Zaragoza, Spain
- VandenBosch G, VandenBossche J, Wagner C, Schouwer PD, VanDeVyvere M, Neels H. 2001. Determination of iron metabolism-related reference values in a healthy adult population. *Clin Chem*. 47(8):1465-1467.
- Vani Pali D, Moeis Emma Sy dan Rotty A.W.Linda. Gambaran Anemia Pada Penderita Penyakit Ginjal Kronik di BLU.RSUP.Prof.DR.R.D.Kandou. Bagian Ilmu Penyakit Dalam Fakultas Kedokteran Universitas Sam Ratulangi Manado- Sulawesi Utara.
- Widiana, I.G.K. 2007. Distribusi penyakit ginjal kronik di Bali : komparasi formula cockcroft-gault dan formula modification of diet in renal. *J Peny Dalam*. Vol: 8.
- Wirawan, R., 2006. Uji Ketelitian Ketepatan dan Nilai Rujukan parameter Retikulosit Orang Indonesia Dewasa Di Jakarta menggunakan alat Hitung Sel darah Otomatik Sysmex XT-2000i. *Balai penerbit FK UI*, Jakarta.
- Wish, J.B. 2006. Assessing iron status: Beyond serum Ferritin and Transferrin Saturation. *Clin J Am Soc Nephrol* 1 : S4 – S8.
- Xia H, Ebben J, Ma JZ, Collins AJ. 1999. Hematocrit levels and hospitalization risks in hemodialysis patients. *J Am Soc Nephrol*. 10:1309–1316.