



## DAFTAR PUSTAKA

- 2320: Alkalinity (Titration Method), 2011. doi: 10.2105/SMWW.2882.023
- 5220 Chemical Oxygen Demand (COD), 2017. APHA. In: Standard Methods For the Examination of Water and Wastewater. America Public Health Association, Washington, DC, USA.
- 5560C Organic and Volatile Acids (2017) *Standard Methods for The Examination of Water and Wastewater*. Washington, DC, USA: America Public Health Association. doi: 10.2105/SMWW.2882.111.
- Adekunle, K. F. and Okolie, J. A. (2015) ‘A Review of Biochemical Process of Anaerobic Digestion’, *Advances in Bioscience and Biotechnology*, 06(03), pp. 205–212. doi: 10.4236/abb.2015.63020.
- Ahring, Birgitte. K. and Westermann, Petter. (1988) ‘Product Inhibition of Butyrate Metabolism by Asetate and Hydrogen in a Thermophilic Coculture’, *Applied and Environment Microbiology*, (Oct), pp. 2393-2397.
- Amin, F. R. et al. (2021) ‘Functions of bacteria and archaea participating in the bioconversion of organic waste for methane production’, *Science of the Total Environment*, 763, p. 143007. doi: 10.1016/j.scitotenv.2020.143007.
- Atasoy, M. et al. (2018) ‘Bio-based volatile fatty acid production and recovery from waste streams: Current status and future challenges’, *Bioresource Technology*. Elsevier, 268(May), pp. 773–786. doi: 10.1016/j.biortech.2018.07.042.
- Atasoy, M.; Eyice, O.; Schnürer, A.; Cetecioglu, Z. Volatile fatty acids production via mixed culture fermentation: Revealing the link between pH, inoculum type and bacterial composition. *Bioresour. Technol.* 2019, 292, 121889.
- Batstone, D. J. et al. (2002) ‘The IWA Anaerobic Digestion Model No 1 (ADM1).’, *Water science and technology : a journal of the International Association on Water Pollution Research*, 45(10), pp. 65–73. doi: 10.2166/wst.2002.0292.
- Beccari, M., Bonemazzi, F., Majone, M., and Riccardi, C. (1996) ‘Interaction between acidogenesis and methanogenesis in the anaerobic treatment of olive oil mill effluents’, *Water Research*, 30(1), pp. 183–189.
- Bermúdez-Penabad, N.; Kennes, C.; Veiga, M.C. Anaerobic digestion of tuna waste for the production of volatile fatty acids. *Waste Manag.* 2017, 68, 96–102.
- Carrillo, D., Karimi, K. and Sárvári, I. (2011) ‘Improvement of biogas production from oil palm empty fruit bunches ( OPEFB )’, *Industrial Crops & Products*, 34(1), pp. 1097–1101. doi: 10.1016/j.indcrop.2011.03.022.



- Chae, K. J., Jang, A., Yim, S. K., and Kim, I. S. (2008) ‘The effects of digestion temperature and temperature shock on the biogas yields from the mesophilic anaerobic digestion of swine manure’, *Bioresource Technology*, 99(1), pp. 1–6.
- Chang, S. H. (2014) ‘An overview of empty fruit bunch from oil palm as feedstock for bio-oil production’, *Biomass and Bioenergy*. Elsevier Ltd, 62, pp. 174–181. doi: 10.1016/j.biombioe.2014.01.002.
- Cheah, Y.-K.; Vidal-Antich, C.; Dosta, J.; Mata-Álvarez, J. Volatile fatty acid production from mesophilic acidogenic fermentation of organic fraction of municipal solid waste and food waste under acidic and alkaline pH. *Environ. Sci. Pollut. Res.* 2019, 26, 35509–35522.
- Cho, H. U. *et al.* (2015) ‘Influence of temperature on volatile fatty acid production and microbial community structure during anaerobic fermentation of microalgae’, *Bioresource Technology*, 191, pp. 475–480. doi: 10.1016/j.biortech.2015.03.009.
- Christia, A. *et al.* (2016) ‘Ethanol production from alkali-pretreated oil palm empty fruit bunch by simultaneous saccharification and fermentation with mucor indicus’, *International Journal of Green Energy*. Taylor and Francis Inc., 13(6), pp. 566–572. doi: 10.1080/15435075.2014.978004.
- Coral Medina, J. D. *et al.* (2015) ‘Lignin preparation from oil palm empty fruit bunches by sequential acid/alkaline treatment - A biorefinery approach’, *Bioresource Technology*. Elsevier Ltd, 194, pp. 172–178. doi: 10.1016/j.biortech.2015.07.018.
- Deublein, D. and A. S. (2011) *Biogas from Waste and Renewable Resources*. Second. Deggendorf: Wiley-VCH.
- Dirar, H. A. and Amin, H. B. El (1988) ‘Methane fermentation of water hyacinth: effect of solids concentration and inoculum source’, *Applied Microbiology and Biotechnology* 4 (3), pp. 299–312. Available at: <https://doi.org/10.1007/BF01096135>.
- Ditjenbun (2020) *Statistik Perkebunan Indonesia 2015-2020: Kelapa Sawit*.
- Fogler, H. S. (1999) *Elements of Chemical Reaction Engineering*, Upper Saddle River, N. J.: Prentice Hall PTR
- Fukuhara, S. (2012) *A basis for the space of modular forms*, *Acta Arithmetica*. doi: 10.4064/aa151-4-5.
- Gerardi H. Michael (2007), *ORP Management in Wastewater as an Indicator of Process Efficiency*
- Girotto, F. *et al.* (2018) ‘Effect of Aeration Applied During Different Phases of



- Anaerobic Digestion', *Waste and Biomass Valorization*, 9(2), pp. 161–174. doi: 10.1007/s12649-016-9785-9.
- Goberna, M. et al. (2015) 'Start-up strategies in manure-fed biogas reactors : Process parameters and methanogenic communities', *Biomass and Bioenergy*, 5, pp. 46–56. doi: 10.1016/j.biombioe.2015.02.003.
- Ishola, M. M. and Taherzadeh, M. J. (2014) 'Bioresource Technology Effect of fungal and phosphoric acid pretreatment on ethanol production from oil palm empty fruit bunches ( OPEFB )', *Bioresource Technology*. Elsevier Ltd, 165, pp. 9–12. doi: 10.1016/j.biortech.2014.02.053.
- Kahar, A. et al. (2017) 'Perpindahan massa fase cair pada pengolahan lindi tpa sampah kota dalam bioreaktor anaerobik'.
- Karki, A. B., Shrestha, J. N. and Bajgain, S. (2005) *Biogas as Renewable Source of Energy in Nepal: Theory and Development*. Kathmandu: BSP-Nepal.
- Khan, M. A. et al. (2016) 'Optimization of process parameters for production of volatile fatty acid, biohydrogen and methane from anaerobic digestion', *Bioresource Technology*, pp. 738–748. doi: 10.1016/j.biortech.2016.08.073.
- Kim, M. et al. (2003) 'Hydrolysis and acidogenesis of particulate organic material in mesophilic and thermophilic anaerobic digestion', *Environmental Technology*. Taylor & Francis Group , 24(9), pp. 1183–1190. doi: 10.1080/09593330309385659.
- Kinanthi et al., (2020)' Recovery of high purity lignin and digestible cellulose from oil palm empty fruit bunch using low acid-catalyzed organosolv pretreatment
- Krayzelova, L. et al. (2015) 'Microaeration for hydrogen sulfide removal during anaerobic treatment: a review', *Reviews in Environmental Science and Bio/Technology*, 14(4), pp. 703–725. doi: 10.1007/s11157-015-9386-2.
- Lee, W. S. et al. (2014) 'A review of the production and applications of waste-derived volatile fatty acids', *Chemical Engineering Journal*. Elsevier B.V., 235, pp. 83–99. doi: 10.1016/j.cej.2013.09.002.
- Lim, P. N. et al. (2015) 'A potential bioconversion of empty fruit bunches into organic fertilizer using Eudrilus eugeniae', *International Journal of Environmental Science and Technology*. Center for Environmental and Energy Research and Studies, 12(8), pp. 2533–2544. doi: 10.1007/s13762-014-0648-2.
- Mandiri, Manual Pelatihan Teknologi Energi Terbarukan, Jakarta, 2012, 61.
- Meegoda, J. N. et al. (2018) 'A review of the processes, parameters, and optimization of anaerobic digestion', *International Journal of Environmental Research and Public Health*, 15(10). doi: 10.3390/ijerph15102224.



- Moraes, B. S. *et al.* (2014) 'Anaerobic digestion of vinasse from sugarcane biorefineries in Brazil from energy, environmental, and economic perspectives: Profit or expense?', *Applied Energy*. Elsevier Ltd, 113, pp. 825–835. doi: 10.1016/j.apenergy.2013.07.018.
- Mosier *et al.*, (2005) 'Features of promising technologies for pretreatment of lignocellulosic biomass'. *Bioresour Technol* 96:673-686.
- Myers, (2019) 'Innovations in Monitoring with Water-Quality Sensors With Case Studies on Floods, Hurricanes, and Harmful Algal Blooms'. *Separation Science and Technology*, Volume 11. <https://doi.org/10.1016/B978-0-12-815730-5.00010-7>.
- Najafpour, G. *et al.* (2005) 'Effect of organic loading on performance of rotating biological contactors using Palm Oil Mill effluents', *Process Biochemistry*, 40(8), pp. 2879–2884. doi: 10.1016/j.procbio.2005.01.002.
- Nghiem *et al.* (2014) 'Oxidation reduction potential as a parameter to regulate micro-oxygen injection into anaerobic digester for reducing hydrogen sulphide concentration in biogas', *Bioresource Technology* 173 (2014) 443–447. doi.org/10.1016/j.biortech.2014.09.052
- Nguyen, D. and Khanal, S. K. (2018) 'A little breath of fresh air into an anaerobic system: How microaeration facilitates anaerobic digestion process', *Biotechnology Advances*. Elsevier, 36(7), pp. 1971–1983. doi: 10.1016/j.biotechadv.2018.08.007.
- O-Thong S, Boe K, Angelidaki I (2012) 'Thermophilic anaerobic co-digestion of oil palm empty fruit bunches with palm oil mill effluent for efficient biogas production. *Appl Energ* 93:648-654
- Osman NB, Syamsuddin N, Uemura Y, (2016) 'Activation carbon of oil palm empty fruit bunch (EFB); core and shaggy', *Procedia Engineering* 148 (2016) 758-764.
- Rasit, N., Idris, A., Harun, R., Ghani, W. A. W. A. K. (2015) 'Effects of lipid inhibition on biogas production of anaerobic digestion from oily effluents and sludges: An overview', *Renewable and Sustainable Energy Reviews*. Elsevier, 45, pp. 351–358.
- Reichert, P. (1994) AQUASIM – A tool for simulation and data analysis of aquatic systems, *Wat. Sci. Tech.*, 30(2), 21-30
- Ruan, D. *et al.* (2019) 'Enhancing methane production of anaerobic sludge digestion by microaeration: Enzyme activity stimulation, semi-continuous reactor validation and microbial community analysis', *Bioresource Technology*. Elsevier, 289(June), p. 121643. doi: 10.1016/j.biortech.2019.121643.



Satpathy, S., Steinigeweg, S., Uhlenhut, F., & Siefert, E. (2013). Application of Anaerobic Digestion Model 1 (ADM1) for Prediction of Biogas Production. International Journal of Scientific & Engineering Research, 4(12), 86-89. Available:<http://www.ncbi.nlm.nih.gov/pubmed/16180468>

Sawatdeenarunat, Sung, dan Khanal, 2017 ' Enhanced volatile fatty acids production during anaerobic digestion of lignocellulosic biomass via micro-oxygenation' Bioresource Technology 237 (2017) 139–145

Seadi, T. A., Rutz, D., Prassl, H., Köttner, M., Finsterwalder, T., Volk, S., and Janssen, R. (2008) *Biogas Handbook*, University of Southern Denmark Esbjerg.

Shuler, M. L. and Kargi, F. (2002) *Bioprocess engineering: Basic concepts*. 2nd edn, 2nd edn. Prentice Hall.

Sun, W., Yu, G., Louie, T., Liu, T., Zhu, C., Xue, G., Gao, P., 2015. From mesophilic to thermophilic digestion: the transitions of anaerobic bacterial, archaeal, and fungal community structures in sludge and manure samples. Appl. Microbiol. Biotechnol. 99 (23), 10271–10282.

Trisakti, B. *et al.* (2018) 'Production of oil palm empty fruit bunch compost for ornamental plant cultivation', in *IOP Conference Series: Materials Science and Engineering*. Institute of Physics Publishing. doi: 10.1088/1757-899X/309/1/012094.

Tsapekos *et al*,2017 'Effect of micro-aeration and inoculum type on the biodegradation of Lignocellulosic substrate' Bioresource Technology 225 (2017) 246–253

Tzun-Wen Shaw, G. *et al.* (2017) 'Inferring microbial interactions in thermophilic and mesophilic anaerobic digestion of HOG waste', *PLoS ONE*, 12(7), pp. 1–22. doi: 10.1371/journal.pone.0181395.

Vavilin, V. A., Rytov, S. V., Lokshina, L. Ya. (2008) 'Hydrolysis kinetics in anaerobic degradation of particulate organic material: An overview', *Waste Management*, 28(6), pp. 939–951.

Wachenheim, D. E., Patterson, J. A. and Ladisch, M. R. (2003) 'Analysis of the logistic function model: Derivation and applications specific to batch cultured microorganisms', *Bioresource Technology*, 86(2), pp. 157–164. doi: 10.1016/S0960-8524(02)00149-9.

Wainaina, S. *et al.* (2019) 'Bioengineering of anaerobic digestion for volatile fatty acids, hydrogen or methane production: A critical review', *Bioengineered*. Taylor & Francis, 10(1), pp. 437–458. doi: 10.1080/21655979.2019.1673937.

Walker, M., Zhang, Y., Heaven, S., Banks, C. (2009) 'Potential errors in the quantitative evaluation of biogas production in anaerobic digestion



- processes', *Bioresource Technology*. Elsevier Ltd, 100(24), pp. 6339–6346.
- Wang, K. *et al.* (2014) 'Anaerobic digestion of food waste for volatile fatty acids (VFAs) production with different types of inoculum: Effect of pH', *Bioresource Technology*, 161, pp. 395–401. doi: 10.1016/j.biortech.2014.03.088.
- Weinrich, S. and Nelles, M. (2021) 'Systematic simplification of the Anaerobic Digestion Model No. 1 (ADM1) – Model development and stoichiometric analysis', *Bioresource Technology*. doi: 10.1016/j.biortech.2021.125124.
- Westerholm, M. and Schnürer, A. (2019) 'Microbial Responses to Different Operating Practices for Biogas Production Systems', *Anaerobic Digestion*. doi: 10.5772/intechopen.82815.
- Zhang, K., Woodruff, A.P., Xiong, M., Zhou, J., Dhande, Y.K., 2011. A synthetic metabolic pathway for production of the platform chemical isobutyric acid. *ChemSusChem* 4 (8), 1068–1070.
- Zhang, Q., Hu, J. and Lee, D. J. (2016) 'Biogas from anaerobic digestion processes: Research updates', *Renewable Energy*. Elsevier Ltd, 98, pp. 108–119.