

DAFTAR PUSTAKA

1. Gonzales. Legal Medicine Pathology and Toxilogy .second edition,
Appleton-Century-Croft Inc USA, 1954 ; p. 381 – 452.
2. Knight B. Forensik Pathology , Edward Arnold Great Britain, 1991 :
p. 222 – 251.
3. Simpson K . Forensic Medicine . eight edition, Edward Arnold Ltd
Great Britain, 1979; p. 74 – 90.
4. Moetrarsi dkk. Epidemiologi Bunuh Diri di Kabupaten Gunung Kidul.
DIY, Berita Kedokteran Masyarakat FK. UGM, 1996;
5. Gradwohl's. Legal Medicine. third edition , John Wright & Son Ltd
Great Britain, 1976; p.191 – 219.
6. Polson CJ. The Essentiale of Forensic Medicine. third edition, Pergamon
Press New York, 1973; p. 204 – 283.
7. Kerr D. Forensic Medicine. fourth edition Adam & Charles Black Great
Britain, 1946; p. 139 – 144.
8. Gordon I. Forensic Medicine A Guide To Principle, Churchill Livingstone
London, 1975; p. 292 – 306.
9. Idries AM. Pedoman Ilmu Kedokteran Forensik. edisi pertama, Binarupa
Aksara, 1989; p.103 – 130.
10. Hamdani N. Ilmu Kedokteran Kehakiman. edisi kedua, PT. Gramedia
Pustaka Utama, 1992; p. 122 – 128.





UNIVERSITAS
GADJAH MADA

**Korban kejahatan dengan senjata api di bagian ilmu kedokteran forensik FK. UNSRI / RSMH
Palembang
tahun 1996-2001**

WIDJAJA, Junus, dr. IBG. Surya Putra Pidada, SpF

Universitas Gadjah Mada, 2003 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Lampiran

Crosstabs

Tahun * Luka Crosstabulation

		Luka		Total	
		Luka tembak	Bukan luka tembak		
Tahun	1996	Count	1	133	134
		% within Luka	3,3%	13,8%	13,5%
1997	Count	2	169	171	
		% within Luka	6,7%	17,5%	17,2%
1998	Count	7	166	173	
		% within Luka	23,3%	17,2%	17,4%
1999	Count	2	155	157	
		% within Luka	6,7%	16,0%	15,8%
2000	Count	10	138	148	
		% within Luka	33,3%	14,3%	14,9%
2001	Count	8	205	213	
		% within Luka	26,7%	21,2%	21,4%
Total	Count	30	966	996	
		% within Luka	100,0%	100,0%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	14,107 ^a	5	,015
Likelihood Ratio	14,232	5	,014
Linear-by-Linear Association	5,791	1	,016
N of Valid Cases	996		

a. 3 cells (25,0%) have expected count less than 5. The minimum expected count is 4,04.

Symmetric Measures

		Value	Asymp. Std. Error ^a	Approx. T ^b	Approx. Sig.
Nominal by Nominal	Contingency Coefficient	,118			,015
Interval by Interval	Pearson's R	-,076	,027	-2,412	,016 ^c
Ordinal by Ordinal	Spearman Correlation	-,075	,027	-2,378	,018 ^c
N of Valid Cases		996			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

Lokasi luka * TKP

Crosstab

Lokasi luka			TKP			Total
			PLG	OKI	MUBA	
Dada Pinggang Perut	Count	1			1	
	% of Total	3,3%			3,3%	
Punggung	Count	3	2		5	
	% of Total	10,0%	6,7%		16,7%	
Dada	Count	3	3	4	10	
	% of Total	10,0%	10,0%	13,3%	33,3%	
Perut	Count		2	1	3	
	% of Total		6,7%	3,3%	10,0%	
Kepala	Count	4		2	6	
	% of Total	13,3%		6,7%	20,0%	
Kepala, Dada	Count		1		1	
	% of Total		3,3%		3,3%	
Leher	Count	1			1	
	% of Total	3,3%			3,3%	
Dada Punggung	Count	1	1		2	
	% of Total	3,3%	3,3%		6,7%	
Kepala Pinggang	Count			1	1	
	% of Total			3,3%	3,3%	
Total	Count	13	9	8	30	
	% of Total	43,3%	30,0%	26,7%	100,0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	16,765 ^a	16	,401
Likelihood Ratio	21,824	16	,149
Linear-by-Linear Association	,217	1	,641
N of Valid Cases	30		

a. 27 cells (100,0%) have expected count less than 5. The minimum expected count is ,27.

Symmetric Measures

		Value	Asymp. Std. Error ^a	Approx. T ^b	Approx. Sig. ^c
Nominal by Nominal	Contingency Coefficient	,599			,401
Interval by Interval	Pearson's R	,087	,180	,460	,649 ^c
Ordinal by Ordinal	Spearman Correlation	,108	,182	,575	,570 ^c
N of Valid Cases		30			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.



PL/PD * TKP

Crosstab

			TKP			Total
			PLG	OKI	MUBA	
PL/PD	PL	Count	7	7	5	19
		% of Total	23,3%	23,3%	16,7%	63,3%
	PL/PD	Count	6	2	3	11
		% of Total	20,0%	6,7%	10,0%	36,7%
Total		Count	13	9	8	30
		% of Total	43,3%	30,0%	26,7%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,315 ^a	2	,518
Likelihood Ratio	1,365	2	,505
Linear-by-Linear Association	,281	1	,596
N of Valid Cases	30		

a. 3 cells (50,0%) have expected count less than 5. The minimum expected count is 2,93.

Symmetric Measures

		Value	Asymp. Std. Error ^a	Approx. T ^b	Approx. Sig.
Nominal by Nominal	Contingency Coefficient	,205			,518
Interval by Interval	Pearson's R	-,098	,187	-,523	,605 ^c
Ordinal by Ordinal	Spearman Correlation	-,111	,188	-,592	,559 ^c
N of Valid Cases		30			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

Umur * TKP

Crosstab

			TKP			Total
			PLG	OKI	MUBA	
Umur 10-19 tahun	Count	2			2	
	% of Total	6,7%			6,7%	
20-29 tahun	Count	7	5	2	14	
	% of Total	23,3%	16,7%	6,7%	46,7%	
30-39 tahun	Count	4	4	5	13	
	% of Total	13,3%	13,3%	19,7%	46,3%	
40-49 tahun	Count			1	1	
	% of Total			3,3%	3,3%	
Total	Count	13	9	8	30	
	% of Total	43,3%	30,0%	26,7%	100,0%	

Chi-Square Tests

	Value	df	Asymp Sig (2-sided)
Pearson Chi-Square	7,620 ^a	6	,267
Likelihood Ratio	8,365	6	,213
Linear-by-Linear Association	6,626	1	,019
N of Valid Cases	30		

a. 10 cells (83,3%) have expected count less than 5. The minimum expected count is ,27.

Symmetric Measures

		Value	Asymp. Std. Error ^a	Approx. T ^b	Approx. Sig.
Nominal by Nominal	Contingency Coefficient	,450			,267
Interval by Interval	Pearson's R	,437	,142	2,567	,016 ^c
Ordinal by Ordinal	Spearman Correlation	,411	,157	2,383	,024 ^c
N of Valid Cases		30			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.