

DAFTAR PUSTAKA

- AYDOGDU, B., TIRELI, G., DEMIRALI, O., GUVENC, U., BESIK, C., KAVUNCUOGLU, S., ... SANDER, S. (2016). Factors affecting early mortality and morbidity in congenital duodenal obstruction: summary of a 20-year experience. *International Surgery*, (August 2018). <https://doi.org/10.9738/int Surg-d-15-00284.1>
- Bethell, G. S., Long, A. M., Knight, M., & Hall, N. J. (2019). Congenital duodenal obstruction in the UK: A population-based study. *Archives of Disease in Childhood: Fetal and Neonatal Edition*, 21, 1–6. <https://doi.org/10.1136/archdischild-2019-317085>
- Bishay, M., Lakshminarayanan, B., Arnaud, A., Garriboli, M., Cross, K. M., Curry, J. I., ... Eaton, S. (2013). The role of parenteral nutrition following surgery for duodenal atresia or stenosis. *Pediatric Surgery International*, 29(2), 191–195. <https://doi.org/10.1007/s00383-012-3200-9>
- Darnifayanti, D., Tjipta, G. D., Rusdidjas, R., & Lubis, B. M. (2015). Immature-to-total neutrophil ratio as an early diagnostic tool of bacterial neonatal sepsis. *Paediatrica Indonesiana*, 55(3), 153. <https://doi.org/10.14238/pi55.3.2015.153-7>
- Delano, M. J., & Ward, P. A. (2016). The immune system's role in sepsis progression, resolution, and long-term outcome. *Immunological Reviews*, 274(1), 330–353. <https://doi.org/10.1111/imr.12499>
- Escobar, M. A., Ladd, A. P., Grosfeld, J. L., West, K. W., Rescorla, F. J., Scherer, L. R., ... Billmire, D. F. (2004). Duodenal atresia and stenosis: Long-term follow-up over 30 years. *Journal of Pediatric Surgery*, 39(6), 867–871. <https://doi.org/10.1016/j.jpedsurg.2004.02.025>
- Fabio Chiarenza, S., Bucci, V., Luisa Conighi, M., Zolpi, E., Costa, L., Fasoli, L., & Bleve, C. (2017). Duodenal Atresia: Open versus MIS Repair-Analysis of Our Experience over the Last 12 Years. *BioMed Research International*, 2017. <https://doi.org/10.1155/2017/4585360>
- Ghrahani, R., Yuniati, T., Judistiani, R. T. D., & Setiabudiawan, B. (2019). Strong Positive Correlation between Neutrophil-to-lymphocyte Ratio and C-reactive Protein in Early Onset Sepsis Korelasi Positif Kuat antara Neutrophil-to-lymphocyte dan C-reactive Protein pada Sepsis Awitan Dinifile:///Users/SaskiaPrathana/Desktop/Thesis. *Majalah Kedokteran Bandung*, 51(38).
- Gomella T, Cunningham M, & Eyal F. (2013). *Lange Neonatology 7th Edition. Neonatology: Management, procedures, on call problems, disease, and drugs*.
- Gutiérrez-carrillo, M. P., Zertuche-coindreau, J. M., Santana-, C. L., Esparza-ponce, C., Sánchez-rodríguez, Y. B., & León, J. C. B. (2013). atresia in the

neonatal period, 81(6), 490–495.

- Gutierrez, I. M., Kang, K. H., Calvert, C. E., Johnson, V. M., Zurakowski, D., Kamin, D., ... Duggan, C. (2012). Risk factors for small bowel bacterial overgrowth and diagnostic yield of duodenal aspirates in children with intestinal failure: A retrospective review. *Journal of Pediatric Surgery*, 47(6), 1150–1154. <https://doi.org/10.1016/j.jpedsurg.2012.03.019>
- Holcomb III, G. W., Murphy, J. P., & St. Peter, S. D. (2020). *Holcomb and Ashcraft's Pediatric Surgery*. (G. W. Holcomb III, J. P. Murphy, & S. D. St. Peter, Eds.), Elsevier (Sevent, Vol. 1). [https://doi.org/10.1016/0011-3840\(82\)90048-X](https://doi.org/10.1016/0011-3840(82)90048-X)
- Liu, X., Shen, Y., Wang, H., Ge, Q., Fei, A., & Pan, S. (2016). Prognostic Significance of Neutrophil-to-Lymphocyte Ratio in Patients with Sepsis: A Prospective Observational Study. *Mediators of Inflammation*, 2016. <https://doi.org/10.1155/2016/8191254>
- Ni, J., Wang, H., Li, Y., Shu, Y., & Liu, Y. (2019). Neutrophil to lymphocyte ratio (NLR) as a prognostic marker for in-hospital mortality of patients with sepsis: A secondary analysis based on a single-center, retrospective, cohort study. *Medicine*, 98(46), e18029. <https://doi.org/10.1097/MD.00000000000018029>
- Rattan, K. N., Singh, J., & Dalal, P. (2016). Neonatal Duodenal Obstruction: A 15-Year Experience. *Journal of Neonatal Surgery*, 5(2), 13. <https://doi.org/10.7860/JCDR/2016/17204.7268>
- Riedemann, N. C., Guo, R. F., & Ward, P. A. (2003). Novel strategies for the treatment of sepsis. *Nature Medicine*, 9(5), 517–524. <https://doi.org/10.1038/nm0503-517>
- Sadler, T. W. (2012). Langman's Essential Medical Embryology. *Langmans Essential Medical Embryology*, 384.
- Sastiono, S., & Rahmaania, J. C. (2019). The relationship between sepsis and timing achievement of peristaltic function in congenital duodenal obstruction. *The New Ropanasuri Journal of Surgery*, 4(1), 24–26. <https://doi.org/10.7454/nrjs.v4i1.73>
- Shah, B. A., & Padbury, J. F. (2014). Neonatal sepsis an old problem with new insights. *Virulence*, 5(1), 170–178. <https://doi.org/10.4161/viru.26906>
- Shames, B. (2019). *Anatomy and Physiology of the Duodenum. Shackelford's Surgery of the Alimentary Tract, 2 Volume Set* (Eighth Edi). Elsevier Inc. <https://doi.org/10.1016/B978-0-323-40232-3.00068-6>
- Singh, R., Zogg, H., Wei, L., Bartlett, A., Ghoshal, U. C., Rajender, S., & Ro, S. (2021). Gut microbial dysbiosis in the pathogenesis of gastrointestinal dysmotility and metabolic disorders. *Journal of Neurogastroenterology and Motility*, 27(1), 19–34. <https://doi.org/10.5056/JNM20149>
- Spitz, L., & Coran, A. G. (2013). *Operative Pediatric Surgery*. (L. Spitz & A. G. Coran, Eds.) (7th ed.). CRC Press.

- Takır, H. B., Karakurt, Z., Saltürk, C., Balçı, M., Kargın, F., Moçin, Ö. Y., ... Adıgüzel, N. (2015). Does total parenteral nutrition increase the mortality of patients with severe sepsis in the ICU? *Turk Toraks Dergisi*, 16(2), 53–58. <https://doi.org/10.5152/ttd.2015.4323>
- Vaishnavi, C. (2013). Translocation of gut flora and its role in sepsis. *Indian Journal of Medical Microbiology*, 31(4), 334–342. <https://doi.org/10.4103/0255-0857.118870>
- Westerdijk, K., Simons, K. S., Zegers, M., Wever, P. C., Pickkers, P., & de Jager, C. P. C. (2019). The value of the neutrophil-lymphocyte count ratio in the diagnosis of sepsis in patients admitted to the Intensive Care Unit: A retrospective cohort study. *PLoS ONE*, 14(2), 1–13. <https://doi.org/10.1371/journal.pone.0212861>
- Wilar, R. (2019). Diagnostic value of eosinopenia and neutrophil to lymphocyte ratio on early onset neonatal sepsis. *Korean Journal of Pediatrics*, 62(6), 217–223. <https://doi.org/10.3345/kjp.2018.06723>