



## DAFTAR PUSTAKA

- Acet, H., Ertaş, F., Bilik *et al* (2015). The relationship between neutrophil to lymphocyte ratio, platelet to lymphocyte ratio and thrombolysis in myocardial infarction risk score in patients with ST elevation acute myocardial infarction before primary coronary intervention. *Postepy w Kardiologii Interwencyjnej*, 11(2), 126–135.
- Adivitiya, & Khasa, Y. P. (2017). The evolution of recombinant thrombolytics: Current status and future directions. *Bioengineered*, 8 (4), 331-358.
- Ali, M. R., Salim Hossain, M., Islam, M. A., Saiful Islam Arman, M., Sarwar Raju, G., Dasgupta, P., & Noshin, T. F. (2014). Aspect of thrombolytic therapy: A review. *Scientific World Journal*, 67 (1), 455-467.
- Arbel, Y., Finkelstein, *et al* (2012). Neutrophil/lymphocyte ratio is related to the severity of coronary artery disease and clinical outcome in patients undergoing angiography. *Atherosclerosis*, 225(2), 456–460.
- Azab, B., Chainani, V., Shah, N., & McGinn, J. T. (2013). Neutrophil-lymphocyte ratio as a predictor of major adverse cardiac events among diabetic population: A 4-year follow-up study. *Angiology*, 64(6), 456–465.
- Azab, B., Zaher, M., Weiserbs, K. F., Torbey, E., Lacossiere, K., Gaddam, S., Lafferty, J. (2010). Usefulness of neutrophil to lymphocyte ratio in predicting short-and long-term mortality after NonST-elevation myocardial infarction. *American Journal of Cardiology*, 106(4), 470–476.
- Bhat, T., Bhat, H., Raza, M., Khoueiry, G., Meghani, M., & Akhtar, M. (2013). Neutrophil to lymphocyte ratio and cardiovascular diseases : a review. *Expert Review of Cardiovascular Therapy*, 11(1), 55–60.
- Blum, A., & Yeganeh, S. (2003). The role of T-lymphocyte subpopulations in acute myocardial infarction. *European Journal of Internal Medicine*, 14(7), 407–410.
- Bugan, B., & Onar, L. C. (2014). The neutrophil-to-lymphocyte ratio: Does it represent inflammation sufficiently without other markers in patients with cardiovascular diseases. *Clinical and Applied Thrombosis/Hemostasis*, 20(3), 341–342.



Čeponienė I., Žaliaduonytė-Pekšienė D., Gustienė O., Tamošiūnas A., Žaliūnas R.  
2014. Association of major cardiovascular risk factors with the development of acute coronary syndrome in Lithuania. *Eur Heart J Suppl* 16(Suppl A): A80–A83.

Chan, D., & Ng, L. L. (2010). Biomarkers in acute myocardial infarction. *BMC Medicine*, 8. 8-34

Chia, S., Nagurney, J., Brown, D., Raffel, O., Bamberg, F., Senatore, F., Wackers, F., & Jang, I. (2009). Association of Leukocyte and Neutrophil Counts With Infarct Size, Left Ventricular Function and Outcomes After Percutaneous Coronary Intervention for ST-Elevation Myocardial Infarction. *The American Journal of Cardiology*, 103(3), 333-337.

Darmawan. (2016). Peran Rasio Netrofil Limfosit sebagai prediktor *major adverse events* tujuh hari dalam perawatan pada pasien sindrom koroner akut. *Penerbit Fakultas Kedokteran Universitas Indonesia* , 8-49

Dwijanarko, W., Maharani, E., & Anggrahini, D. W. (2017). T Peak–T End Interval Alteration as Parameter of Successful Fibrinolysis in Patients with ST Segment Elevation Acute Myocardial Infarction. *ACI (Acta Cardiologia Indonesiana)*, 3(1), 14.

Germany, G. H., Germany, A. K., Lenzen, M. J., Denmark, E. P., & Vranckx, P. (2018). 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST segment elevation, *Web Addenda The Task Force for the management of acute myocardial infarction* 119–177.

Germany, G. H., Germany, A. K., Lenzen, M. J., Denmark, E. P., & Vranckx, P. (2017). 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST segment elevation, *Web Addenda The Task Force for the management of acute myocardial infarction*, 1–8.

Girdhar, R., Kothari, Y., Raj, R. A., & Shastry, P. (2018). Successful or Unsuccessful Thrombolysis with Streptokinase in Acute Myocardial Infarction : A Descriptive Study, *Journal of Medical Science And clinical Research* 06(03), 731–735.

Girish Ronad, Girish & Dixit, Rohit. 2015. Factors influencing the outcome of thrombolysis in acute myocardial infarction. *Indian Journal of Basic and Applied Medical Research*; June 2015: Vol.-4, Issue- 3, P. 545-555



- Guo, Z., Yu, S., Xiao, L., *et al.* (2016). Dynamic change of neutrophil to lymphocyte ratio and hemorrhagic transformation after thrombolysis in stroke. *Journal of Neuroinflammation*, 1–8.
- Hancock, J. E., Cooke, J. C., Chin, D. T., & Monaghan, M. J. (2002). Determination of successful reperfusion after thrombolysis for acute myocardial infarction: A noninvasive method using ultrasonic tissue characterization that can be applied clinically. *Journal of Circulation*, 105(2), 157–161.
- Hamm CW., Bassand JP, Agewall S, Bax J., Boersma E., Hector B., Hubert K., *et al.* (2011). ESC Guidelines for management for acute coronary syndromes in patients presenting without persistent ST-segment elevation. *European Heart Journal* 32: 3003
- Han, Y. C., Yang, T. H., Kim, D. Il, Jin, H. Y., Chung, S. R., Seo, J. S., ... Kim, D. S. (2013). Neutrophil to lymphocyte ratio predicts long-term clinical outcomes in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. *Korean Circulation Journal*, 43(2), 93–99.
- Huang, G., Zhong, X. N., Zhong, B., Chen, Y. Q., Liu, Z. Z., Su, L., ... Yin, Y. H. (2009). Significance of white blood cell count and its subtypes in patients with acute coronary syndrome. *European Journal of Clinical Investigation*, 39(5), 348–358.
- Ibanez, B., James, S., Agewall, S., Antunes, M. J., Bucciarelli-Ducci, C., Bueno, H., ... Gale, C. P. (2018). 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *European Heart Journal*, 39 (2), 119-177.
- Juzar D. 2012. Sindrom Koroner Akut. In: L.I.Rilantono,1st ed. Penyakit kardiovaskular (PKV). Jakarta: Badan Penerbit Fakultas Kedokteran Universitas Indonesia, hal. 138-160.
- Kapoor S., Opneja A., Lalitha N. 2018. The Role of Neutrophils in Thrombosis. *Thromb Res.* 170 : 87-96.
- Khalifa, R., I.Chamouri W. *et al.* 2020. Predictors of Successful Thrombolysis in an Acute Myocardial Infarction. *Archives of Cardiovascular Diseases Supplements*, 12(1), 18.
- Kiefe C.I., Williams O.D., Bild D.E., Lewis C.E., Hilner J.E., Oberman A. 1997. Regional disparities in the incidence of elevated blood pressure among young adults: the cardia study. *Circulation* 96:1082–1088.



- Kleinschmidt K.C. 2006. Epidemiology and patophysiology of acute coronary syndrome. *J. Johns Hopkins Advances Studies in Nursing* (4): 72–7.
- Kovar D., Canon C.P., Bentley J.H., Charlesworth A., Rogers WJ. 2004. Does Initial and Delayed Heart Rate Predict Mortality in Patients with Acute Coronary Syndrome. *Clinical Cardiology* 27:80-8
- Kumar A dan Cannon C.P. 2009. Acute Coronary Syndromes: Diagnosis and Management, Part 1. *Mayo Clinic Proc* 84(10):917
- Kumar A dan Cannon C.P. 2009. Acute coronary syndromes: diagnosis and management part 1. *Mayo Clinic Proc* 84(10): 917–38.
- Kucia, A. M., & Zeitz, C. J. (2002). Failed reperfusion after thrombolytic therapy: Recognition and management. *Heart and Lung: Journal of Acute and Critical Care*, 31(2), 113–121.
- Kurtul, A., Murat, S. N., Yarlioglu, M., Duran, M., Celik, I. E., Kilic, A., & Ocek, A. H. (2015). Increased neutrophil-to-lymphocyte ratio predicts persistent coronary no-flow after wire insertion in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention. *Clinics*, 70(1), 34–40
- Kounis, N. G., Soufras, G. D., Tsikas, G., & Hahalis, G. (2015). White blood cell counts, leukocyte ratios, and eosinophils as inflammatory markers in patients with coronary artery disease. *Clinical and Applied Thrombosis/Hemostasis*, 21(2), 139–143.
- Libby, P. (2012). History of Discovery : Inflammation in Atherosclerosis. *Arterioscler Thromb Vasc Biol.*, 32(9), 2045–2051.
- Libby, P., Simon, D.I. 2001. Inflammation and Thrombosis: The Clot Thickens. *Journal of Circulation* 103, 1718-1720.
- Libby, P. & Theroux, P. 2005. Pathophysiology of Coronary Artery Disease. *Circulation American Heart Association* vol 11, 3481-3488.
- Madjid, M., & Fatemi, O. (2013). Components of the complete blood count as risk predictors for coronary heart disease: In-depth review and update. *Texas Heart Institute Journal*, 40(1), 17–29.



- Mor, A., Luboshits, G., Planer, D., Keren, G., & George, J. (2006). Altered status of CD4+CD25+ regulatory T cells in patients with acute coronary syndromes. *European Heart Journal*, 27(21), 2530–2537.
- Munir, T. A., & Afzal, M. N. (2010). Assessment of differential leukocyte count in patients with acute coronary syndrome. *Journal of the Pakistan Medical Association*, 60(7), 548–551.
- Nunez, J., Minana, G., Bodi, V., Nunez, E., Sanchis, J., Husser, O. (2011). Low Lymphocyte Count and Cardiovascular Diseases. *Curent Medicinal Chemistry*, 18, 3226- 3233.
- Oncel, R. C., Ucar, M., Karakas, M. S., Akdemir, B., Yanikoglu, A., Gulcan, A. R., ... Demir, I. (2015). Relation of neutrophil-to-lymphocyte ratio with GRACE risk score to in-hospital cardiac events in patients with ST-segment elevated myocardial infarction. *Clinical and Applied Thrombosis/Hemostasis*, 21(4), 383–388.
- Palareti, G., Legnani, C., Cosmi, B., Antonucci, E., Erba, N., Poli, D., Testa, S., & Tosetto, A. (2016). Comparison between different D-Dimer cutoff values to assess the individual risk of recurrent venous thromboembolism: Analysis of results obtained in the DULCIS study. *International Journal of Laboratory Hematology*, 38(1), 42–49.
- Park, J. S., Seo, K. W., Choi, B. J., Choi, S. Y., Yoon, M. H., Hwang, G. S., Tahk, S. J., & Shin, J. H. (2018). Importance of prognostic value of neutrophil to lymphocyte ratio in patients with ST-elevation myocardial infarction. *Medicine (United States)*, 97(48).
- Park, J. J., Jang, H. J., Oh, I. Y., Yoon, C. H., Suh, J. W., Cho, Y. S., ... Choi, D. J. (2013). Prognostic value of neutrophil to lymphocyte ratio in patients presenting with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention. *American Journal of Cardiology*, 111(5), 636–642.
- Pasceri, V., Andreotti, F., & Maseri, A. (1996). Clinical markers of thrombolytic success. *European Heart Journal*, 17 :35–41.
- PERKI (Perhimpunan Dokter Spesialis Kardiovaskular Indonesia). 2015. *Pedoman Tatalaksana Sindrom Koroner Akut*. [http://www.inaheart.org/upload/file/Pedoman\\_tatalaksana\\_Sindrom\\_Korone\\_r\\_Akut\\_2015.pdf](http://www.inaheart.org/upload/file/Pedoman_tatalaksana_Sindrom_Korone_r_Akut_2015.pdf)



- Shah, N., Parikh, V., Patel, N., Patel, N., Badheka, A., Deshmukh, A., Lafferty, J. (2014). Neutrophil lymphocyte ratio significantly improves the Framingham risk score in prediction of coronary heart disease mortality: Insights from the National Health and Nutrition Examination Survey-III. *International Journal of Cardiology*, 171(3), 390–397.
- Soehnlein, O. (2012). Multiple roles for neutrophils in atherosclerosis. *Circulation Research*, 110(6), 875–888.
- Soumerai, S. B., McLaughlin, T. J., Ross-Degnan, D., Christiansen, C. L., & Gurwitz, J. H. (2002). Effectiveness of thrombolytic therapy for acute myocardial infarction in the elderly: Cause for concern in the old-old. *Archives of Internal Medicine*, 162(5), 561–568.
- Tamhane, U. U., Aneja, S., Montgomery, D., Rogers, E. K., Eagle, K. A., & Gurm, H. S. (2008). Association Between Admission Neutrophil to Lymphocyte Ratio and Outcomes in Patients With Acute Coronary Syndrome. *American Journal of Cardiology*, 102(6), 653–657.

