VARIASI GENETIK E23K GENA POTASSIUM INWARDLY RECTIFYING CHANNEL SUB FAMILY J MEMBER 11 (KCNJ11) DAN
VARIASI GENETIK Ala55Val GENA UNCOUPLING PROTEIN-2 (UCP-2) SEBAGAI FAKTOR RISIKO DIABETES MELITUS
TIPE 2 DENGAN OBESITAS PADA ETNIK PAPUA DI KOTA JAYAPURA
JEMS KIFEN ROGET M, Dr. Dra. Pramudji Hastuti, Apt., MS; dr. Ahmad Hamim Sadewa, Ph.D
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

DAFTAR PUSTAKA

Abd-Elsalam K. (2003). Bioinformatic tools and guideline for PCR primer design. *J. Bio.* 2(5): 91-95.

Agyropoulous G., Brown A.M., Peterson R., Likes CE., Watson DK., Garvey WT. (1998). Structure and organization of the human Uncoupling Protein 2 gene and identification of a common biallelic variant in Caucasian and African-American Subject. *Diabetes*. 47: 685-687.

American Diabetes Association. (2006). Diagnosis and classification of diabetes mellitus. *Diabetes Care.* 29(1 Suppl): 43S - 48S.

American Diabetes Association. (2008). Standarts of medical care in diabetes mellitus-2008, *Diabetes Care.* 31(1 Suppl): 12S - 35S.

Ali Rastegari., Mohammad R., Hamid Mirmohammad Sadeghi., Elham Faghih Imani1., Akbar Hasanzadeh., Fatemeh Moazen. (2014). Association of KCNJ11 (E23K) gene polymorphism with susceptibility to type 2 diabetes in Iranian patients. *Adv. Biomed Res 2015;4;1. Sciences, Isfahan, Iran*. DOI: 10.4103/2277-9175.148256

Aschroft FM dan Rorsman P. (2004). Type 2 diabetes mellitus: not quite exciting enough. *Hum. Mol. Genet.* 13: R21 - R31

Aschroft FM. (2005). ATP-sensitive potassium channelopathies: focus on insulin secretion. JCI. 115 : 2047-58.

Astrup A., Toubro S., Daargard LT., Urhammer SA., Sorensen TI., Pedersen O. (1999). Impact of the v/v 55 polymorphism of the uncoupling protein 2 gene on 24-h energy expenditure and substrate oxidation. *Int. J. Obes Relat.* Metab. Disord. 23: 1030-1034.

Badman MK., Flier JS. (2005). The gut and anergy balance: visceral allies in the obesity wars. *(A review of the role of the gut in energy homeostatis)*. *Science*. 307.

Ball GD., Huang TT., Gower BA., Crus ML., Shaibi GO., Weigensberg MJ *et al.* (2006). Longitudinal Changes in Insulin sensitivity, insulin secretion, and beta-cell function during puberty. *J Pediatr.* 148: 16-22

Baroso I., Luan J., Middelberg RPS., Harding AH., Franks PW., Clayton D., *et al*. (2003). candidate gene association study in type 2 diabetes mellitus Indicates a role for genes involved in β-cell function as well as insulin action. *PLoS Biol.* 1(3):445.

VARIASI GENETIK E23K GENA POTASSIUM INWARDLY RECTIFYING CHANNEL SUB FAMILY J MEMBER 11 (KCNJ11) DAN
VARIASI GENETIK Ala55Val GENA UNCOUPLING PROTEIN-2 (UCP-2) SEBAGAI FAKTOR RISIKO DIABETES MELITUS
TIPE 2 DENGAN OBESITAS PADA ETNIK PAPUA DI KOTA JAYAPURA
JEMS KIFEN ROGET M, Dr. Dra. Pramudji Hastuti, Apt., MS; dr. Ahmad Hamim Sadewa, Ph.D
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Bogardus C., Lillioja S., Ravussin E. (2003). Familial dependence of the resting metabolic rate. *N. Engl. J. Med.* 315: 96-100.

Bouchard C. (1991). Current Understanding of Etiology of Obesity Genetic and Non-Genetic Factors. *Am J.Clin Nutr*. 53: 1561-1565.

Britz B., Siegfried W., Ziegler A., Lamertz C., Herpertz-Dahlmann BM., Remschmidt H *et al.* (2000). Rates of psychiatric disorders in a clinical study groupof adolescents with extreme obesity and in obese adolescents ascertained via a population based study. *Int. J Obes Relat Metab Disord;* 24: 1707-1714.

Clement K., Garner C., Harger J., Philippi A., LeDuc C., Carcy A., (1996). Indication for Linkage of the Human Ob Gene Region With Extreme Obesity. *Diabetes* 45: 687-690.

Comuzzie., Anthony G. (2003). The genetics of obesity in Mexican Americans: The Evidence From Genome Scanning Efforts in the San Antonio Family Heart Study. *Human Biology* 75(5): 635-646.

Corbalan MS., Marti A., Forga L., Martinez-Gonzales MA., Martines JA. (2002). b2-adrenergic receptor mutation and abdominal obesity risk: effect modification by gender and HDL-cholesterol. *Eur. J. Nutr*, 41.114-118.

Daargard L., & Pedersen. (2001) Uncoupling protein functional characteristics and role in the phatogenesis of obesity and type 2 diabetes. *Diabetologia*, 44(8), 946-65.

Daniels S.R. (2009). Complications of Obesity in Children and adolescents. *International J. of Obesity* (2005), 33 Suppl 1. S60-S65. doi: 10.1038/ijo. 2009.20.

Decroli Eva. (2019). Diabetes Melitus Tipe 2. Pusat Penerbit Bagian Ilmu Penyakit Dalam, *Fakultas Kedokteran Universitas Andalas.* Edisi Pertama. Padang.

DeFronzo RA. (1997). Pathogenesis of type 2 diabetes metabolic and molecule implication for identifying diabetes genes. Diabetes Rev. 4: 177-269

Defronzo R. (2008). Banting lecture dipresentasikan di Annual Meeting of ADA San Fransisco.

de Meyts P. (2004). Insulin and its receptor structure, function and evolution. *Bio. Essays.* 26: 1351-1362

UNIVERSITAS GADJAH MADA
VARIASI GENETIK E23K GENA POTASSIUM INWARDLY RECTIFYING CHANNEL SUB FAMILY J MEMBER 11 (KCNJ11) DAN
VARIASI GENETIK Ala55Val GENA UNCOUPLING PROTEIN-2 (UCP-2) SEBAGAI FAKTOR RISIKO DIABETES MELITUS
TIPE 2 DENGAN OBESITAS PADA ETNIK PAPUA DI KOTA JAYAPURA
JEMS KIFEN ROGET M, Dr. Dra. Pramudji Hastuti, Apt., MS; dr. Ahmad Hamim Sadewa, Ph.D
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

de Simone G., . Devereur RB., Daniels SR., Koren MJ., Meyer RA., Laragh JH. (1995). Effect of Growth on Variability of left ventricular mass: assessment of allometric signals in adults and children and their capacity to predict cardiovascular risk. *J Am Coll Cardiol.* 25: 1056-1062.

de Souza BM., Assmann TS., Kliemann LM., Gross JL., Canani LH., Crispim D. (2011). the role of uncoupling protein 2 (UCP2) on the development of type 2 diabetes mellitus and its chronic complications. *Arg. Bras. Endocrinol. Metab*. 55(4): 239-480.

Deurenberg-Yap M., Chew SK., Lin VF., Tan BY., Van Staveren W A., Deurenberg P. (1999). Relationships between indices of obesity and its comorbidities in multi-ethnic Singapore. *Int J. Obes Relat Metab Disord*: 23: 1136-42

Departemen Kesehatan (2008). *Laporan Riset Dasar Kesehatan Yogyakarta Tahun 2007*. Departemen Kesehatan RI, Jakarta.

Donohoue, P.A. (2004). Obesity dalam Nelson Textbook of Pediatric 17th Edition. *Philadelphia: Saunders*.

Drewnowski A, Popkin BM. (1997). The nutrition transition: new trends in the global diet. *Nutr Rev* 55, 31–43.

Ello-Martin JA., Ledikwe JH., Rolls BJ. (2005). The influence of food portion size and energy density on energy intake: implications for weight management. *Am J Clin Nutr.* 82: 236S-241S.

Emy Huriyati1., Harry F Luglio1., Prima D Ratrikaningtyas2., Ahmad F A Tsani1., Ahmad H Sadewa., Mohammad Juffrie. (2016). Dyslipidemia, insulin resistance and dietary fat intake in obese and normal weight adolescents: the role of uncoupling protein 2 -866G/A gene polymorphism. *Int J Mol Epidemiol Genet* 2016;7(1):67-73

Ezenwaka, C., Kalloo,R., Uhlig, M., Schwenk, R., & Eckel, J. 2005. The E23K variant in the Kir6.2 subunit of the ATP-sensitive K channel does not augment impaired glucose tolerance in Caribbean subjects with a family history of type 2 diabetes. *J Endocrin*. 185:439-44

Florez JC. 2008. The genetic of type 2 diabetes : A realistic appraisal in 2008. *J Clin endocrinol Metab* 93 : 4633-4642

UNIVERSITAS
GADJAH MADA
**VARIASI GENETIK E23K GENA POTASSIUM INWARDLY RECTIFYING CHANNEL SUB FAMILY J MEMBER 11 (KCNJ11) DAN
VARIASI GENETIK Ala55Val GENA UNCOUPLING PROTEIN-2 (UCP-2) SEBAGAI FAKTOR RISIKO DIABETES MELITUS
TIPE 2 DENGAN OBESITAS PADA ETNIK PAPUA DI KOTA JAYAPURA**
JEMS KIFEN ROGET M, Dr. Dra. Pramudji Hastuti, Apt., MS; dr. Ahmad Hamim Sadewa, Ph.D
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Fischer, A., Fischer, E., Ohlig, M.M., Schulze, M., Hoffmann, K., Weickert, M.O., e*t al.*, (2008). KCNJ11 E23K affects diabetes risk and is associated with the disposition index. *Diabetes Care*. 31: 87-89.

Frizzel JP. (2001). Handbook of Pathophysiology, *Springhouse Corporation,* Philadelphia.

Froguel P and Boutin P. (2001). Genetic of Pathways Regulating Body Weight in The Development of Obesity in Humans. *Exp Biol Med.* 226(11):991996.

Gloyn, A.L., Hashim, Y., Aschroft, S.J.H., Ashfield, R.. Wiltshire, S., Turner, R.C. (2001). Association studies of variants in promoter & coding region of beta-cell ATP-sensitive K-channel genes SUR1 and Kir6.2 with tyoe 2 diabetes mellitus (UKPDS 53). *Diabet Met.* 1: 206-212

Guyton, C.A., Hall, J.E., 2007. *Buku Ajar Fisiologi Kedokteran*. EGC, Jakarta.

Grant, S.F.A., Hakonarson, H. & Schwartz, S., (2010). Can the genetics of type 1 and type 2 diabetes shed light on the genetics of latent autoimmune diabetes in adults? *Endo rev,* 31, pp.183–193.

Haghvirdizadeh P., Mohamed Z., Abdulla NA., Haghvirdizadeh P., Haerian MS., Haerian BS. (2014). Review Article; KCNJ11: Genetic Polymorphism and Risk of Diabetes mellitus. *J Diab Res.* 1 - 9.

Hill J.O., Kriketos A.D dan Peters. (2000). Disturbances of Energy Balance dalam: *Biochemical and Physiological Aspect of Human Nutrition.* Philadelphia: Saunders.

Hirsch J., Salans L.B dan Aronne L.J. (1995). Obesity dalam Principles and Practice of Endocrinology and metabolism Second Edition. J.B. Philadelphia Lippincott Company.

Indra Rasjad Muhammad. (2006). Dasar Genetik Obesitas Viseral. Kedokteran Brawijaya, Vol. XXII. No.1. Malang.

International Association For The Study Of Obesity (IASO)/International Obesity Task Force (IOTF) (2000). The Asia-Pacific Perspective: Redefinig Obesity and its Treatment. *Health Communications.* Australia

VARIASI GENETIK E23K GENA POTASSIUM INWARDLY RECTIFYING CHANNEL SUB FAMILY J MEMBER 11 (KCNJ11) DAN
VARIASI GENETIK Ala55Val GENA UNCOUPLING PROTEIN-2 (UCP-2) SEBAGAI FAKTOR RISIKO DIABETES MELITUS
TIPE 2 DENGAN OBESITAS PADA ETNIK PAPUA DI KOTA JAYAPURA
JEMS KIFEN ROGET M, Dr. Dra. Pramudji Hastuti, Apt., MS; dr. Ahmad Hamim Sadewa, Ph.D
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

International Diabetes Federation / IDF. (2011)

Istiantho. (2009). Hubungan Faktor Pejamua, Agen dan Lingkungan terhadap peningkatan prevalensi DM di Ternate. FKUI., Jakarta.

Jensen M.D. (2008). Role of Body Fat distribution and the metabolic complications of obesity. *J.clin endocrinol metab*. 93. S57-S63. doi: 10.1210/jc.2008-1585.

Kadowaki T, Hara K, Yamauchi T, Terauchi Y, Tobe K, Nagai R. 2003. Molecular mechanism of insulin resistance and obesity. *Exp Biol Med*.228:111-117.

Kim YB., Nikoulina SE., Ciaraldi TP., Henry RR dan Kahn BB. (1999). Normal Insulin-dependent activation of Akt/protein kinase B, with diminished activation of phosphoinositide 3-kinase, in muscle in type 2 diabetes. *J. Clin. Invest.* 104: 733-741

Koentjaningrat. (2005). Pengantar Antropologi, Jilid I dan II, cetakan ke tiga.

Koster JC., Permutt MA dan Nichols CG. (2005). Perspectives in Diabetes : Diabetes and Insulin Secretion The ATP-Sensitive K+ Channel (KATP) Connection. *Diabetes*. 54: 3065-72.

Laukkanen O., Pihlajamaki J., Lindstrom J., Eriksson J., Valle TT., Hamalainen H., llanne-Parikka P., Keinanen-Kiukaanniemi S., Tuomilehto J., Uusitupa M., Laakso M. (2004). Polymorphisms of the SUR1 (ABCC8) and Kir6.2 (KCNJ11) Genes Predict Conversion from Impaired Glucose Tolerance to Type 2 Diabetes. The Finnish Diabetes. *J Clin Endocrinol Metab* 2004 Dec; 89(12): 6286-90. doi: 10.1210/jc.2004-1204

Laakso M., Edelman SV., Brechtel G., Baron AD. (1990). Decreased effect of insulin to stimulate skeletal muscle blood flow in obese man. *J Clin Invest*. 85: 1844-1852.

Lean ME, Han TS, Morrison CE. (1995). Waist circumference as a measure for indicating need for weight management. *BMJ.* ; 311:158-161

Low S*, et al.* (2009). Review on epidemic of obesity. *Ann Acad Med Singapore;* 38(1): 57-9.

Limin Wei, Meng Wei , Lei Chen, Shanshan Liang, Fanfan Gao, Xin Cheng, Hongli Jiang (2021). Low-density lipoprotein cholesterol : highdensity lipoprotein cholesterol ratio is associated with incident diabetes in Chinese adults: A retrospective cohort study. *J Diabetes Investig* 2021; 12: 91–98.

VARIASI GENETIK E23K GENA POTASSIUM INWARDLY RECTIFYING CHANNEL SUB FAMILY J MEMBER 11 (KCNJ11) DAN
VARIASI GENETIK Ala55Val GENA UNCOUPLING PROTEIN-2 (UCP-2) SEBAGAI FAKTOR RISIKO DIABETES MELITUS
TIPE 2 DENGAN OBESITAS PADA ETNIK PAPUA DI KOTA JAYAPURA
JEMS KIFEN ROGET M, Dr. Dra. Pramudji Hastuti, Apt., MS; dr. Ahmad Hamim Sadewa, Ph.D
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Madiyono B, Moeslihan S, Sastroasmoro, Budiman I, Purwanto SH. 2008. Dalam : Sastroasmoro, S, Ismael, S. (ed). Dasar-dasar Metodologi Penelitian Klinis, *3nd Ed. CV Sagung Seto*, Jakarta : 302-331

Mangmool S., Denkaew T., Parichatikanond W dan Kurose H. (2017). b-adrenergic Receptor and Insulin Resistance in the heart. *Biomol. Ther.* 25: 44-56

Marentek E. (2006). Resistensi Insulin pada Diabetes Melitus tipe 2. Poliklinik Endkrin Metabolik. *Bagian Penyakit Dalam Rumah Sakit Umum Gowa*. Makassar.

Maulana M. (2009). *Mengenal Diabetes Mellitus, Panduan Praktis mengenai penyakit kencing manis*. Ed 2. Katahati, Yogyakarta.

McCarthy M. 2010. Genomics, type 2 diabetes and obesity. *N Eng J Med*.363 : 2339-2350.

McGarry JD.2002. Dysregulation of fatty acid metabolism in the etiology of type 2 diabetes. *Diabetes.* 51:7-15.

Murray, R. K., Granner, D. K., & Rodwell, V. W. Biokimia harper (30 ed.). Jakarta: Buku Kedokteran EGC; 2020

Mozo J., Emre Y., Bouillaud F., Ricquier D & Criscuolo F. (2005). Thermoregulation: What role for UCPs in mammals and birds? *Biosci Rep* 25, 227-249.

Oberkofler H., Liu YM., Esterbauer H., Hell E., Krempler F., Patsch W. (1998). Uncoupling protein-2 gene: reduced mRNA expression in intraperitoneal adipose tissue of obese humans. *Diabetologia.* 41: 940-946

Oktavianthi S., Trimarsanto H., Febinia CA., Suastika K., Saraswati MR., Dwipayana P., Arindrarto W., Sudoyo H., Malik SG. (2012). Uncoupling Protein 2 gene Polymorphisms are associated with obesity. *Cardiovasc Diabetol* (CVD). 2012.11.41.

Oppert JM., Vohl MC., Chagnon M., Dionne FT., Cassad-Doulcier AM., Ricquier D., Perusse L., Bouchard C. (1994). DNA Polymorphism in the uncoupling protein (UCP) Gene and Human Body Fat. *Int J. Obes Relat Metab Disord*. 18: 526-531.

O'Rahilly S. (2009). Human geneticts illuminates the paths to metabolic disease. *Nature.* 462: 307 - 314.

Pecqueur C., Cassard-Doulcier AM., Raimbault S., Miroux B., Fleury C., Gelly C., Bouillaud F., Ricquier D. (1999). Functional Organization of the human

UNIVERSITAS GADJAH MADA
VARIASI GENETIK E23K GENA POTASSIUM INWARDLY RECTIFYING CHANNEL SUB FAMILY J MEMBER 11 (KCNJ11) DAN
VARIASI GENETIK Ala55Val GENA UNCOUPLING PROTEIN-2 (UCP-2) SEBAGAI FAKTOR RISIKO DIABETES MELITUS
TIPE 2 DENGAN OBESITAS PADA ETNIK PAPUA DI KOTA JAYAPURA
JEMS KIFEN ROGET M, Dr. Dra. Pramudji Hastuti, Apt., MS; dr. Ahmad Hamim Sadewa, Ph.D
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

uncoupling protein 2 gene, and juxtaposition to the uncoupling protein 3 gene. Biochem. *Biop. Res. Comm.* 255(1): 40-6.

PERKENI. (2019). Konsensus Pengelolaan dan Pencegahan Diabetes Mellitus tipe-2 di Indonesia, Pengurus Besar Perkumpulan Endokrinologi Indonesia.

Pine DS, Goldstein RB, Wolk S, Weissman MM. (2001). The association between childhood depression and adulthood body mass index. *Pediatrics*; 107: 1049–1056.

Popinat G, Cousse S, Goldfarb L, Becker S, Gardin I, Salaün M. (2019). Sub-cutaneous Fat Mass measured on multislice computed tomography of pretreatment PET/CT is a prognostic factor of stage IV non-small cell lung cancer treated by nivolumab. *J. OncoImmunology,* 8:5, e1580128, DOI: 10.1080/2162402X.2019.1580128

Pua YH., Ong PH. (2005). Anthropometric indices as screening tools for cardiovascular risk factors in Singaporean Women. *Asia Pac. J.Clin Nutr*. 14:74-9

Prabandari AS. (2009). *Polimorfieme p.E23K gene Potassium inwardly rectiflying channel sub family J member 11 (KCNJ11) sebagai faktor risiko diabetes melitus tipe 2 di yogyakarta (tesis),* Pascasarjana Universitas Gadjah Mada.

Radha KS., Vimaleswaran R., Deepa and Mohan V. (2003). The Genetic of diabetes mellitus. *Indian J Med Res.*117, pp 225-38.

Radha V and Mohan V.2007. Genetic predispotition to type 2 diabetes among Asian Indians. *Indian J Med Res*.125:259-60.

Rahilly S. (2001). Uncoupling protein 2: adiposity angel and diabetes devil?. *Nature Med.* 7,770 - 2.

Ricquier D., Bouillaud F. (2000). The Uncoupling Protein Homologues: UCP1, UCP2, UCP3, StUCP and AtUCP. *Biochem J.* 345(2): 161-79.

Riedel MJ., Boora P., Steckley D., de Vries G., dan Light PE. (2003). Kir6.2 Polymorphisms Sensitize β-Cell ATP Sensitive Pottassium Channels to Activation by Acyl CoAs : Possible Cellular Mechanism for increased Susceptibility to Type 2 Diabetes? *Diabetes 52(10). 2630 - 35*.

Riset Kesehatan Dasar (Riskesda). (2013). Badan Penelitian dan Pengembangan Kesehatan, Kementerian Kesehatan RI, Jakarta.

Rolls ET.(2005). Emotion Explained. Oxford University Press: Oxford

UNIVERSITAS GADJAH MADA
VARIASI GENETIK E23K GENA POTASSIUM INWARDLY RECTIFYING CHANNEL SUB FAMILY J MEMBER 11 (KCNJ11) DAN
VARIASI GENETIK Ala55Val GENA UNCOUPLING PROTEIN-2 (UCP-2) SEBAGAI FAKTOR RISIKO DIABETES MELITUS
TIPE 2 DENGAN OBESITAS PADA ETNIK PAPUA DI KOTA JAYAPURA
JEMS KIFEN ROGET M, Dr. Dra. Pramudji Hastuti, Apt., MS; dr. Ahmad Hamim Sadewa, Ph.D
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Rolls ET.(2007). Understanding the mechanisms of food intake and obesity. *Obesity reviews 8 (Suppl. 1)*. 67-72.

Rosen ED., Hsu CH., Wang X., Sakai S., Freeman M.W., Gonzales FJ., *et al*. (2002). 'C/EBP Induces Adipogenesis Through PPAR: A Unified Pathway'. *Genes Dev*., 16: 22-26

Rosen ED & Spiegelman BM. (2000). 'Molecular Regulation of Adipogenesis' Annu. Rev. *Cell Dev. Biol*. 16: 145-171.

Rousset S., Alves-Guerra MC., Mozo J., Miroux B., Cassard AM., Bouillaud F., Ricquier D. (2004). The biology of mitochondrial uncoupling protein. *Diabetes*. 53. Suppl 1: S130-5.

Salopura T. (2009). Variation in the UCP2 and UCP3 genes associates with abdominal obesity and serum lipids: The Finnish Diabetes prevention study. *BMC Med. Gen*. 10:94.

Sami F. Khalil, Mas S. Mohktar, and Fatimah Ibrahim., (2014). The Theory and Fundamentals of Bioimpedance Analysis in Clinical Status Monitoring and Diagnosis of Diseases. *J. **Sensors***. mdpi. *14*, 10895-10928;

Sandika, J. (2020). Rasio Triglyceride / High Density Lipoprotein-Cholesterole dan Resistensi Insulin sebagai Faktor Risiko Diabetes Melitus Tipe 2. Fakultas Kedokteran, Universitas Lampung. Majority | Volume 9 | Nomor 1 | 2020|

Schwanstecher C., Meyer U., dan Schwanstecher M. (2002). Kir6.2 Polymorphisme predisposes to Type 2 Diabetes by Inducing Overactivity of Pancreatic β-Cell ATP-Sensitive K+ Channels. *Diabetes* 51(3): 875-79.

Schwanstecher C dan Schwanstecher M. (2002). Nucleotida Sensitivity of Pancreatic ATP-Sensitive of Pottasium Channels and Type 2 Diabetes. *Diabetes* 51. Suppl 3: S358-S362.

Schwimmer JB., Deutsch R., Rauch JB., Behling C., Newbury R., Lavine JE. (2003). Obesity, insulin resistance, and other clinicopathological correlates of pediantric nonalcoholic fatty liver disease. J Pediatr: 143: 500-505.

Schwimmer JS., Burwinkle TM., Varni JW. (2003). Health-related quality of life of severely obese children and adolescents. *JAMA*; 289: 1813-1819.

Shashikiran U., Sudha V., Jayaprakash B. (2004). What is Obesity? *Med J Malaysia*. 59: 130-3

UNIVERSITAS GADJAH MADA
VARIASI GENETIK E23K GENA POTASSIUM INWARDLY RECTIFYING CHANNEL SUB FAMILY J MEMBER 11 (KCNJ11) DAN
VARIASI GENETIK Ala55Val GENA UNCOUPLING PROTEIN-2 (UCP-2) SEBAGAI FAKTOR RISIKO DIABETES MELITUS
TIPE 2 DENGAN OBESITAS PADA ETNIK PAPUA DI KOTA JAYAPURA
JEMS KIFEN ROGET M, Dr. Dra. Pramudji Hastuti, Apt., MS; dr. Ahmad Hamim Sadewa, Ph.D
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Sorof J & Daniels S. (2002). Obesity Hypertension in Children: a Problem of epidemic proportions. *Hypertension.* 40: 441-4417

Soeharto, I., 2002., Kolesterol dan lemak jahat, kolesterol dan lemak baik dan proses terjadinya serangan jantung dan stroke, Gramedia Pustaka Utama, Jakarta, pp.48, 72-74, 120-124.

Steinberg HO., Brechtel G., Johnson A., Fineberg N., Baron AD. (1994). Insulin-Mediated skeletal muscle vasodilation is nitric oxide release. *J Clin Invest.* 94: 1172-1179.

Steinberg HO., Paradisi G., Hook G., Crowder K., Cronin J., Baron AD. (2000). Free fatty Acid elevation impairs insulin-mediated vasodilation and nitic oxide production. *Diabetes.* 49: 1231-1238.

Sumiarsih T. (2006). Peran biologi Uncoupling Protein 2 (UCP2) pada obesitas dan diabetes melitus tipe 2 (DMT2). *J. Kedokteran Yarsi;* 14(3): 243-249.

Sunita R, Sadewa A H., Fatmawati A . (2015). Lower HOMA-β Values are detected among individuals with variant of E23K polymorphism of potassium inwardly-rectifying channel, subfamily J, member 11 *(KCNJ11)* gene. *The egyptian J. of Medical Hman Genetics.* 16, 227-231.

Suyono S., Wapadji S., Soegondo S., Soewondo P., Subekti I. (2009). *Penatalaksanaan Diabetes Melitus Terpadu, Panduan Penetalaksanaan Diabetes Melitus Bagi Dokter dan Edukator*, FKUI. Jakarta.

Tabara Y., Osawa H., Kawamoto R., Onuma H., Shimizu I., Makino H., Kohara K., Miki T. (2011). Genotipe risk score of common susceptible variants for prediction of type 2 diabetes mellitus in Japanese: the Shimanami Health Promoting Progam (J-SHIPP study). Development of type 2 diabetes mellitus and genotype risk score. *Metabolism*. 60 (11): 1634-40.

Thevenod F. (2008). Pathophysiology of diabetes mellitus type 2: Role of obesity, Insulin Resistance and β-Cell dysfunction. *F. Diab. Basel*. 19 ppl-18

Tischler. (2004). Neoruendocrinology: Obesity. *Lecture.* 52: 1-12

Wahjuni S, (2013). Biokimia metabolic, Udayana University Press, cetakan pertama,

Wang H., Chu WS., Lu T., Hasstedt SJ., Kern PA., Elbein SC. (2004). Uncoupling protein-2 polymorphismes in type 2 diabetes, obesity, and insulin secretion. Am. J. Physiol. Endocrinol. Metab. 286 (1): E1-E7.

UNIVERSITAS
GADJAH MADA
**VARIASI GENETIK E23K GENA POTASSIUM INWARDLY RECTIFYING CHANNEL SUB FAMILY J MEMBER 11 (KCNJ11) DAN
VARIASI GENETIK Ala55Val GENA UNCOUPLING PROTEIN-2 (UCP-2) SEBAGAI FAKTOR RISIKO DIABETES MELITUS
TIPE 2 DENGAN OBESITAS PADA ETNIK PAPUA DI KOTA JAYAPURA**
JEMS KIFEN ROGET M, Dr. Dra. Pramudji Hastuti, Apt., MS; dr. Ahmad Hamim Sadewa, Ph.D
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

White MF., Kahn CR. (1994). The Insulin signaling system. *J. Biologic. Chem*. 264: 1-4

World Heald Organization. (2016). *Obesity: preventing and managing the global epidemic. (WHO Tehnical Report Series no.894)*. Geneva, Switzerland

World Heald Organization, 2011, Diabetes, *Fact sheet*, No. 312

Xu K., Zhang M., Cui D., Fu Y., Qian L., Gu R., Wang M., Shen C., Yu R., Yang T. (2011). UCP2 -866G/A and Ala55Val, and UCP3 -55C/T Polymorphisms in association with type 2 diabetes susceptibility: a meta-analysis study. *Diabetologia*. 54 (9): 3215-24

Zhou D., Zhang D., Liu Y., Zhao T., Chen Z., Liu Z., Yu L *et al*. (2009). The E23K Variation in the KCNJ11 gene is associated with Type 2 Diabetes in Chinese and East Asian Population. *J Hum Genet.* 54:433-35

Zhang M., Wang M., Zhao Z-T. (2014). Uncoupling protein 2 gene polymorphism in association with overweight and obesity susceptibility : A meta analysis. *Meta Gene 2. 143-159.j.mgene.* 2013.10.009.