

DAFTAR PUSTAKA

- Alexopoulos EC, 2010. Introduction to Multivariate Regression Analysis. *Hippokratia* 14: 23–28.
- Asali, F., & Phillips, C., 2012. The management of urogynaecological problems in pregnancy and the early postpartum period. *Obstet. Gynaecol.* 14: 153–158. doi:10.1111/j.1744-4667.2012.00111.x
- Aydin, S., & Aydin, C.A., 2015. Evaluation of labor-related pelvic floor changes 3 months after delivery : a 3D transperineal ultrasound study. *Int. Urogynecol. J.* doi:10.1007/s00192-015-2774-x
- Barber, M.D., & Maher, C., 2013. Epidemiology and outcome assessment of pelvic organ prolapse. *Int. Urogynecol. J.* 24: 1783–1790. doi:10.1007/s00192-013-2169-9
- Barber, M.D., Walters, M.D., & Bump, R.C., 2005. Short forms of two condition-specific quality-of-life questionnaires for women with pelvic floor disorders (PFDI-20 and PFIQ-7). *Am. J. Obstet. Gynecol.* 193: 103–113. doi:10.1016/j.ajog.2004.12.025
- Blomquist, J.L., Muñoz, A., Carroll, M., & Handa, V.L., 2018. Association of Delivery Mode With Pelvic Floor Disorders After Childbirth. *JAMA* 21204. doi:10.1001/jama.2018.18315
- Boyle, R., Hay-smith, E.J.C., Cody, J.D., & Mo, S., 2014. Pelvic Floor Muscle Training for Prevention and Treatment of Urinary and Fecal Incontinence in Antenatal and Postnatal Women : A Short Version Cochrane Review. *Neurourol. Urodynamics, Wiley Online Libr.* 276: 269–276. doi:10.1002/nau
- Bozkurt, M., Yumru, A.E., & ahin, L., 2014. Pelvic floor dysfunction, and effects of pregnancy and mode of delivery on pelvic floor. *Taiwan. J. Obstet. Gynecol.* 53: 452–458. doi:10.1016/j.tjog.2014.08.001
- Brubaker, L., Bump, R., Fynes, M., Jaquetin, B., Karram, M., Kreder, K., et al., 2008. Surgery for Pelvic Organ Prolapse. *Int. Cont. Soc.* 1371–1402.
- Choi, K.H., & Hong, J.Y., 2014. Management of Pelvic Organ Prolapse 693–702.
- Corton, M.M., 2009. Anatomy of Pelvic Floor Dysfunction. *Obstet. Gynecol. Clin. North Am.* 36: 401–419. doi:10.1016/j.ogc.2009.09.002
- Cunningham, F.G., Leveno, K.J., & Bloom, S.L., 2013. Williams Obstetrics 23rd Edition-idw.
- Dave, A., Joshi, R., Goel, V., & Dave, A., 2017. Original Research Article MR imaging evaluation for the assessment of pelvic organ prolapse : a newer technique. *Int. J. Reprod. Contraception, Obstet. Gynecol.* 6: 3400–3404. doi:http://dx.doi.org/10.18203/2320-1770.ijrcog20173451

- Dhama, V., Chaudhary, R., Singh, S., & Singh, M., 2017. Evaluation of pelvic organ prolapse by standardized POP Q system for vaginal hysterectomy. *Int. J. Reprod. Contraception, Obstet. Gynecol.* 6: 2584. doi:10.18203/2320-1770.ijrcog20172356
- Dietz, H.P., 2004. Ultrasound imaging of the pelvic floor . Part I : two-dimensional aspects. *Ultrasound Obs. Gynecol* 23: 80–92. doi:10.1002/uog.939
- Dietz, H.P., Haylen, B.T., & Broome, J., 2001. Ultrasound in the quantification of female pelvic organ prolapse. *Ultrasound Obs. Gynecol* 2001; 18: 511–514 Blackwell.
- Dietz, H.P., & Shek, K.L., 2009. Levator defects can be detected by 2D translabial ultrasound. *Int. Urogynecol. J.* 20: 807–811. doi:10.1007/s00192-009-0839-4
- Dumolin, C., Hay-Smith, E., & Mac Habée-Séguin, G., 2019. Pelvic floor muscle training versus no treatment, or inactive control treatments, for urinary incontinence in women. *Res. Nurs. Heal.* 42: 234–235. doi:10.1002/nur.21946
- Gachon, B., Fritel, X., Pierre, F., & Nordez, A., 2021. Transperineal ultrasound shear - wave elastography is a reliable tool for assessment of the elastic properties of the levator ani muscle in women. *Sci. Rep.* 1–9. doi:10.1038/s41598-021-95012-8
- Gao, Y., Zhao, Z., Yang, Y., Zhang, M., Wu, J., & Miao, Y., 2019. Diagnostic value of pelvic floor ultrasonography for diagnosis of pelvic organ prolapse : a systematic review Diagnostic value of pelvic floor ultrasonography for diagnosis of pelvic organ prolapse : a systematic review. *Int. Urogynecol. J.* doi:10.1007/s00192-019-04066-w
- Gyhagen, M., Bullarbo, M., Nielsen, T.F., & Milsom, I., 2013. The prevalence of urinary incontinence 20 years after childbirth: A national cohort study in singleton primiparae after vaginal or caesarean delivery. *BJOG An Int. J. Obstet. Gynaecol.* 120: 144–151. doi:10.1111/j.1471-0528.2012.03301.x
- Hagen, S., Glazener, C., McClurg, D., Macarthur, C., Elders, A., Herbison, P., et al., 2017. Pelvic floor muscle training for secondary prevention of pelvic organ prolapse (PREVPROL): a multicentre randomised controlled trial. *Lancet* 389: 393–402. doi:10.1016/S0140-6736(16)32109-2
- Halle, T.K., Gunvor, J.S., Kari, H., Engh, M.E., & Siafarikas, F., 2020. Change in prevalence of major levator ani muscle defects from 6 weeks to 1 year postpartum , and maternal and obstetric risk factors : A longitudinal ultrasound study. *Acta Obstet. Gynecol. Scand.* 1403–1410. doi:10.1111/aogs.13878
- Jain, P., & Parsons, M., 2011. The effects of obesity on the pelvic floor. *Obstet. Gynaecol.* 13: 133–142. doi:10.1576/toag.13.3.133.27664

- Jelovsek, J.E., Ms, K.C., Gyhagen, M., Hagen, S., Wilson, D., Kattan, M.W., et al., 2018. Predicting risk of pelvic floor disorders 12 and 20 years after delivery. *Am. J. Obstet. Gynecol.* 218: 222.e1-222.e19. doi:10.1016/j.ajog.2017.10.014
- Jelovsek, J.E., Piccorelli, A., Barber, P.M.D., Tunitsky-bitton, E., & Kattan, M.W., 2013. Prediction Models for Postpartum Urinary and Fecal Incontinence in Primiparous Women. *female Pelvic Med. Reconstr. Surg.* 19: 110–118. doi:10.1097/SPV
- Jennewein, L., Kielland-kaisen, U., Paul, B., Mo, C.J., Klemm, S., Schulze, S., et al., 2018. Maternal and neonatal outcome after vaginal breech delivery at term of children weighing more or less than 3 . 8 kg : A FRABAT prospective cohort study 1–14.
- Jeon, J., & Na, S., 2017. Vacuum extraction vaginal delivery: Current trend and safety. *Obstet. Gynecol. Sci.* 60: 499–505. doi:10.5468/ogs.2017.60.6.499
- JNPK-KR, POGI, IDAI, IBI, & PPNI, 2008. Buku Acuan Pelatihan Klinik Asuhan Persalinan Normal, Asuhan Esensial, Pencegahan dan Penanggulangan Segera Komplikasi Persalinan dan Bayi Baru Lahir, Revisi 5. *JNPK-KR Depkes RI Jakarta.*
- Jundt, K., Peschers, U., & Kentenich, H., 2015. MEDIZIN: CME: Diagnostik und Therapie der Weiblichen Beckenbodendysfunktion. *Dtsch. Arztebl. Int.* 112. doi:10.3238/arztebl.2015.0564
- Karjalainen, P.K., Mattsson, N.K., Jalkanen, J.T., Nieminen, K., & Tolppanen, A.-M., 2020. Minimal important difference and patient acceptable symptom state for PFDI-20 and POPDI-6 in POP surgery. *Int. Urogynecol. J.* doi:https://doi.org/10.1007/s00192-020-04513-z
- Kementerian Kesehatan, 2018. Data dan Informasi, Profil Kesehatan Indonesia 2017. *Kementeri. Kesehat. Republik Indones. Jakarta.*
- Kementerian Kesehatan, 2016a. Ringkasan Eksekutif Data dan Informasi Kesehatan Provinsi DI Yogyakarta. *Kementeri. Kesehat. Republik Indones. Jakarta.*
- Kementerian Kesehatan, 2016b. Indonesian Case Based Groups (INA-CBG's) dan non Indonesian Case Based. *Peratur. MENTERI Kesehat. REPUBLIK Indones.* 52.
- Kementerian Kesehatan, WHO, IDI, POGI, & IBI, 2013. Buku Saku Pelayanan Kesehatan Ibu di Fasilitas Kesehatan Dsar dan Rujukan, Pedoman bagi Tenaga KesehatanN, 1st ed. *Kementreian Kesehat. Republik Indones. Jakarta.*
- Kochenour, N., Peterson, C., Adashi, C., & Burrows, P., 1997. Human Reproduction: Clinical, Pathologic and Pharmacologic Correlations. *Knowl. Weav. Spencer S. Eccles Heal. Sci. Libr. Univ. Utah.*

- Leeman, L., Rogers, R., Borders, N., Teaf, D., & Qualls, C., 2016. The Effect of Perineal Lacerations on Pelvic Floor Function and Anatomy at 6 Months Postpartum in a Prospective Cohort of Nulliparous Women. *Birth* 43: 293–302. doi:10.1111/birt.12258
- Mandrekar, J.N., 2010. Receiver operating characteristic curve in diagnostic test assessment. *J. Thorac. Oncol.* 5: 1315–1316. doi:10.1097/JTO.0b013e3181ec173d
- Masenga, G.G., Shayo, B.C., & Rasch, V., 2018. Prevalence and risk factors for pelvic organ prolapse in Kilimanjaro, Tanzania: A population based study in Tanzanian rural community. *PLoS One* 13: 1–13. doi:<https://doi.org/10.1371/journal.pone.0195910>
- MaBlo, K., Möllers, M., Murcia, K.O. de, Klockenbusch, W., & Ralf Schmitz, 2018. New Method for Assessment of Levator Avulsion Injury. *J Ultrasound Med* 00: 1–7. doi:10.1002/jum.14810
- Memon, H.U., & Handa, V.L., 2013. Vaginal childbirth and pelvic floor disorders. *Women's Heal.* 9: 265–277. doi:10.2217/whe.13.17
- Meston, C.M., 2003. Validation of the female sexual function index (Fsfi) in women with female orgasmic disorder and in women with hypoactive sexual desire disorder. *J. Sex Marital Ther.* 29: 39–46. doi:10.1080/713847100
- Pangastuti, N., 2018. Instrumen prediktor disfungsi dasar panggul pasca persalinan vaginal.
- Pangastuti, N., Sari, D.C.R., Santoso, B.I., Agustiniingsih, D., & Emilia, O., 2018. Gambaran Faktor Risiko Prolaps Organ Panggul Pasca Persalinan Vaginal di Daerah Istimewa Yogyakarta Risk Factors for Pelvic Organ Prolapse in Women with History of Vaginal Delivery in Yogyakarta. *Maj. Kedokt. Bandung* 50: 102–108.
- Persu, C., Chapple, C.R., Cauni, V., Gutue, S., & Geavlete, P., 2011. Pelvic Organ Prolapse Quantification System (POP-Q) - a new era in pelvic prolapse staging. *J. Med. Life* 4: 75–81.
- Pregazzi, R., Sartore, A., Bortoli, P., Grimaldi, E., Ricci, G., & Guaschino, S., 2002. Immediate postpartum perineal examination as a predictor of puerperal pelvic floor dysfunction. *Obstet. Gynecol.* 99: 581–584. doi:10.1016/S0029-7844(01)01763-X
- Schwertner-Tiepelmann, N., Thakar, R., Sultan, A.H., & Tunn, R., 2012. Obstetric levator ani muscle injuries: current status. *Ultrasound Obs. Gynecol* 39: 372–383. doi:10.1002/uog.11080
- Seo, J.T., & Kim, J.M., 2006. Voiding Dysfunction. *J. Urol.* 175: 1769–1772. doi:10.1016/S0022-5347(05)00993-6
- Sigrist, R.M.S., Liau, J., Kaffas, A. El, Chammas, M.C., & Willmann, J.K., 2017.

T h e r a n o s t i c s Ultrasound Elastography : Review of Techniques and Clinical Applications. *Theranostics* 7: 1303–1328. doi:10.7150/thno.18650

Utomo, E., Blok, B.F., Steensma, A.B., & Korfage, I.J., 2014. Validation of the pelvic floor distress inventory (PFDI-20) and pelvic floor impact questionnaire (PFIQ-7) in a Dutch population. *Int. Urogynecol. J.* 25: 531–544. doi:10.1007/s00192-013-2263-z

WHO, 2008. WHO POLICY ON NON-RECRUITMENT OF SMOKERS OR OTHER TOBACCO USERS. *World Health Organ.* 6–7.

Wilson, D., Dornan, J., Milsom, I., & Freeman, R., 2014. UR-CHOICE: can we provide mothers-to-be with information about the risk of future pelvic floor dysfunction? *Int. Urogynecol. J.* 25: 1449–1452. doi:10.1007/s00192-014-2376-z

Zhuang, R.R., Song, Y.F., Chen, Z.Q., Ma, M., Huang, H.J., Chen, J.H., et al., 2011. Levator avulsion using a tomographic ultrasound and magnetic resonance – based model. *Am. J. Obstet. Gynecol.* doi:10.1016/j.ajog.2011.03.052