

REFERENCES

- Abdul Gh, M.A., Defronzo, R.A., 2010. Pathogenesis of insulin resistance in skeletal muscle. *J. Biomed. Biotechnol.* 2010:47-79
- Algaidi, S.A., Eldomiaty, M.A., Elbastwisy, Y.M., Almasry, S.M., Desouky, M.K., Elnaggar, A.M., 2019. Effect of voluntary running on expression of myokines in brains of rats with depression. *Int. J. Immunopathol. Pharmacol.* 3:20-33
- Allen DL, Hittel DS, McPherron AC. 2011. Expression and function of myostatin in obesity, diabetes, and exercise adaptation. *Med Sci Sports Exerc.* 43(10). 1828-1835
- American Diabetes Association, 2005. Diagnosis and Classification of Diabetes Mellitus. Retrieved from https://care.diabetesjournals.org/content/37/Supplement_1/S81.full-text.pdf
- American Diabetes Association.,2010. Diagnosis and classification of diabetes mellitus [published correction appears in Diabetes Care. *Diabetes Care.* 33 (1)
- Amirican diabetes association, 2020. Classification and Diagnosis of Diabetes: Standards of Medical Care in Diabetes—2020. Retrieved from https://care.diabetesjournals.org/content/43/Supplement_1/S14
- Artasensi A, Pedretti A, Vistoli G, Fumagalli L, 2020. Type 2 Diabetes Mellitus: A Review of Multi-Target Drugs. *Molecules.* 25(8):1987
- Asiimwe, D., Mauti, G.O., Kiconco, R., 2020. Prevalence and Risk Factors Associated with Type 2 Diabetes in Elderly Patients Aged 45-80 Years at Kanungu District.

J. Diabetes Res. 2020:146-152

Azad, M., Khaledi, N., Hedayati, M., 2016. Effect of acute and chronic eccentric exercise on FOXO1 mRNA expression as fiber type transition factor in rat skeletal muscles. *Gene*. 584(2):180-184

Universitate de Barcelona, 2018. New tool to improve studies on exercise physiology - Universitat de Barcelona. Retrieved from https://www.ub.edu/web/ub/en/menu_eines/noticies/2019/07/033.html

Bassi, D., Bueno, P. de G., Nonaka, K.O., Selistre-Araujo, H.S., Leal, A.M. de O., 2015. Exercise alters myostatin protein expression in sedentary and exercised streptozotocin-diabetic rats. *Arch. Endocrinol. Metab.* 59(2): 148–153

Barrientos, G. *et al.* (2015) ‘Cholesterol removal from adult skeletal muscle impairs excitation-contraction coupling and aging reduces caveolin-3 and alters the expression of other triadic proteins’, *Frontiers in Physiology*, 6(MAR), pp. 1111–1115

Biglari, S., Afousi, A.G., Mafi, F., Shabkhiz, F., 2020. High-intensity interval training-induced hypertrophy in gastrocnemius muscle via improved IGF-I/Akt/FoxO and myostatin/Smad signaling pathways in rats. *Physiol. Int*

Bogdanovich, S., Perkins, K.J., Krag, T.O.B., Whittemore, L., Khurana, T.S., 2005. Myostatin propeptide-mediated amelioration of dystrophic pathophysiology. *FASEB J.* 19(6):543-549

Bois, P.R.J., Grosveld, G.C., 2003. FKHR (FOXO1a) is required for myotube fusion

of primary mouse myoblasts. *EMBO J.* 22(5):47-57

Bonaldo, P., Sandri, M., 2013. Cellular and molecular mechanisms of muscle atrophy.

DMM Dis. Model. Mech. 6(1): 25–39.

Bordoni B, V.M., 2021. Anatomy, Bony Pelvis and Lower Limb, Gastrocnemius

Muscle. Retrieved from <https://www.ncbi.nlm.nih.gov/books/NBK532946/>

Busquets, S., Toledo, M., Marmonti, E., Orpí, M., Capdevila, E., Betancourt, A.,

López-Soriano, F.J., Argilés, J.M., 2012. Formoterol treatment downregulates

the myostatin system in skeletal muscle of cachectic tumour-bearing rats. *Oncol.*

Lett. 3(1): 185–189.

Capozza, F., Combs, T.P., Cohen, A.W., Cho, Y., Park, S., Schubert, W., Williams,

T.M., Brasaemle, D.L., Jelicks, L.A., Scherer, P.E., Kim, J.K., Lisanti, M.P.,

Combs, T.P., Cohen, A.W., Park, S., Schubert, W., Williams, T.M., Brasaemle,

D.L., Jelicks, L.A., Scherer, P.E., Kim, J.K., Caveolin-, M.P.L., 2021. Caveolin-

3 knockout mice show increased adiposity and whole body insulin resistance ,

with ligand-induced insulin receptor instability in skeletal muscle. *Am J Physiol*

Cell Physiol. 288(6): 1317–1331

CARL J. CASPERSEN, KENNETH E. POWELL, GREGORY M. CHRISTENSON,

1985. Physical Activity, Exercise, and Physical Fitness: Definitions and

Distinctions for Health-Related Research Synopsis. *Public Health Rep.* 100(2):

126–131.

Carnac, G., Vernus, B., Bonnieu, A., 2007. Myostatin in the Pathophysiology of

Skeletal Muscle. *Curr Genomics.* 8(7):415-422

Chavanelle, V. *et al.* (2017) 'Effects of high-intensity interval training and moderate-intensity continuous training on glycaemic control and skeletal muscle mitochondrial function in db/db mice', *Scientific Reports*, 7(1), pp. 1–10

CDC, 2021. Diabetes Risk Factors. Retrieved from <https://www.cdc.gov/diabetes/basics/risk-factors.html>

Chavanelle, V., Boisseau, N., Otero, Y.F., Combaret, L., 2017. Effects of high-intensity interval training and moderate-intensity continuous training on glycaemic control and skeletal muscle mitochondrial function in db / db mice. *Sci Rep.* 7(1): 204

Cho, J.E., Fournier, M., Da, X., Lewis, M.I., 2010. Time course expression of Foxo transcription factors in skeletal muscle following corticosteroid administration. *J. Appl. Physiol.* 108(1): 137–145

Cohen, A.W., Combs, T.P., Scherer, P.E., Lisanti, M.P., 2003. Role of caveolin and caveolae in insulin signaling and diabetes. *Am. J. Physiol. - Endocrinol. Metab.* 285(6):1151-1160

Colberg, S.R., Sigal, R.J., Fernhall, B., Regensteiner, J.G., Blissmer, B.J., Rubin, R.R., Chasan-Taber, L., Albright, A.L., Braun, B., 2010. Exercise and type 2 diabetes: The American College of Sports Medicine and the American Diabetes Association: Joint position statement. *Diabetes Care.* 33(12):147-167

Colberg, S.R., Sigal, R.J., Yardley, J.E., Riddell, M.C., Dunstan, D.W., Dempsey, P.C., Horton, E.S., Castorino, K., Tate, D.F., 2016. Physical activity/exercise and diabetes: A position statement of the American Diabetes Association. *Diabetes Care* 39(11): 2065–2079

- Corrotte, M., Almeida, P.E., Tam, C., Castro-Gomes, T., Fernandes, M.C., Millis, B.A., Cortez, M., Miller, H., Song, W., Mangel, T.K., Andrews, N.W., 2013. Caveolae internalization repairs wounded cells and muscle fibers. *Elife* 2: 1–30
- Dalmau-Pastor, M., Fargues-Polo, B.J., Casanova-Martínez, D.J., Vega, J., Golanó, P., 2014. Anatomy of the triceps surae: a pictorial essay. *Foot Ankle Clin.* 19(4): 603–635.
- Diabetes UK, 2019. The risk factors of Type 2 diabetes. Retrived from <https://www.diabetes.org.uk/preventing-type-2-diabetes/diabetes-risk-factors>
- Dumitru, A. and Radu, M. (2018) ‘Section II Basic aspects of muscle atrophy 4 . Muscle changes during atrophy’, (February 2019)
- Draznin, B., 2006. Molecular mechanisms of insulin resistance: Serine phosphorylation of insulin receptor substrate-1 and increased expression of p85 α : The two sides of a coin. *Diabetes* 55(8): 2392–2397
- Edgerton, V.R., Smith, J.L., Simpson, D.R., 1975. Muscle fibre type populations of human leg muscles. *Histochem. J.* 7(3): 259–266
- El-Kawy Tayel, S. G. A. *et al.* (2021) ‘Histological structure of skeletal muscle in disuse muscle atrophy and role of exercise’, *Egyptian Journal of Histology*, 44(1), pp. 1–7
- Escribano, O., Beneit, N., Rubio-Longás, C., López-Pastor, A.R., Gómez-Hernández, A., 2017. The Role of Insulin Receptor Isoforms in Diabetes and Its Metabolic and Vascular Complications. *J. Diabetes Res.* 2017
- Fargion, S., Dongiovanni, P., Guzzo, A., Colombo, S., Valenti, L., Fracanzani, A.L.,

2005. Iron and insulin resistance. *Aliment. Pharmacol. Ther. Suppl.* 2: 61–63
- Fathi Hamouda, A., 2018. Ethical to using Rats in the Scientific Researches. *Pharm. Pharmacol. Int. J.* 6(1): 23–25
- Fiala GJ, Minguet S. Caveolin-1, 2018. The Unnoticed Player in TCR and BCR Signaling. *Adv Immunol.* 137:83-133
- Food and drugs administration, 2019. Retrived from <https://www.fda.gov/consumers/consumer-updates/how-safely-use-glucose-meters-and-test-strips-diabetes>
- Frontera, W.R., Ochala, J., 2015. Skeletal Muscle: A Brief Review of Structure and Function. *Behav. Genet.* 96(3): 183–195
- Frush, T.J., Noyes, F.R., 2015. Baker's Cyst: Diagnostic and Surgical Considerations. *Sports Health* 7(4): 359–365.
- Galbiati, F., Volonté, D., Chu, J.B., Li, M., Fine, S.W., Fu, M., Bermudez, J., Pedemonte, M., Weidenheim, K.M., Pestell, R.G., Minetti, C., Lisanti, M.P., 2000. Transgenic overexpression of caveolin-3 in skeletal muscle fibers induces a Duchenne-like muscular dystrophy phenotype. *Proc. Natl. Acad. Sci. U. S. A.* 97(17): 9689–9694
- Gazzerro, E., Sotgia, F., Bruno, C., Lisanti, M.P., Minetti, C., 2010. Caveolinopathies: From the biology of caveolin-3 to human diseases. *Eur. J. Hum. Genet.* 18: 137–145
- González Coraspe, J.A., Weis, J., Anderson, M.E., Münchberg, U., Lorenz, K., Buchkremer, S., Carr, S., Zahedi, R.P., Brauers, E., Michels, H., Sunada, Y.,

Lochmüller, H., Campbell, K.P., Freier, E., Hathazi, D., Roos, A., 2018.

Biochemical and pathological changes result from mutated Caveolin-3 in muscle.

Skelet. Muscle 8(1): 1–19

Groussard, C. *et al.* (2020) ‘Tissue-specific oxidative stress modulation by exercise: A comparison between MICT and HIIT in an obese rat model’, *Oxidative Medicine and Cellular Longevity*, 2019

Gross, D.N., Van Den Heuvel, A.P.J., Birnbaum, M.J., 2008. The role of FoxO in the regulation of metabolism. *Oncogene* 9(3):208-214.

Gu, X., Reagan, A.M., McClellan, M.E., Elliott, M.H., 2017. Caveolins and caveolae in ocular physiology and pathophysiology. *Prog. Retin. Eye Res.* 56: 84–106.

Greaves P, Chouinard L, Ernst H, Mecklenburg L, Pruimboom-Brees IM, Rinke M, Rittinghausen S, Thibault S, von Erichsen J, Yoshida T. 2013. Proliferative and non-proliferative lesions of the rat and mouse soft tissue, skeletal muscle, and mesothelium. *J Toxicol Pathol* 26(3 suppl):1S-26S. Abstract: <http://www.ncbi.nlm.nih.gov/pubmed/25035576>

Hagiwara, Y., Sasaoka, T., Araishi, K., Imamura, M., Yorifuji, H., Nonaka, I., Ozawa, E., Kikuchi, T., 2000. Caveolin-3 deficiency causes muscle degeneration in mice. *Hum. Mol. Genet.* 9: 3047–3054.

Harvard Medical School, 2018. Type 2 Diabetes Mellitus. Retrived from https://www.health.harvard.edu/a_to_z/type-2-diabetes-mellitus-a-to-z

Hernández-Deviez, D.J., Howes, M.T., Laval, S.H., Bushby, K., Hancock, J.F., Parton,

- R.G., 2008. Caveolin regulates endocytosis of the muscle repair protein, dysferlin. *J. Biol. Chem.* 283(10): 6476–6488
- Hittel, D. S. *et al.* (2009) 'Increased secretion and expression of myostatin in skeletal muscle from extremely obese women', *Diabetes*, 58(1), pp. 30–38
- Huang Y, Deng Y, Shang L, Yang L, Huang J, Ma J, Liao X, Zhou H, Xian J, Liang G, Huang Q, 2019. Effect of type 2 diabetes mellitus caveolin-3 K15N mutation on glycometabolism. *Exp Ther Med.* 18(4):2531-2539
- Jørgensen, S.B., Wojtaszewski, J.F.P., Viollet, B., Andreelli, F., Birk, J.B., Hellsten, Y., Schjerling, P., Vaulont, S., Neufer, P.D., Richter, E.A., Pilegaard, H., 2005. Effects of alpha-AMPK knockout on exercise-induced activation in mouse skeletal muscle. *FASEB J.* 19(9):1146-1148
- Kamei, Y., Hattori, M., Hatazawa, Y., Kasahara, T., Kanou, M., Kanai, S., Yuan, X., Suganami, T., Lamers, W.H., Kitamura, T., Ogawa, Y., 2014. FOXO1 activates glutamine synthetase gene in mouse skeletal muscles through a region downstream of 3'-UTR: Possible contribution to ammonia detoxification. *Am. J. Physiol. - Endocrinol. Metab.* 307(6): 485–493
- Kamei, Y., Miura, S., Suzuki, M., Kai, Y., Mizukami, J., Taniguchi, T., Mochida, K., Hata, T., Matsuda, J., Aburatani, H., Nishino, I., Ezaki, O., 2004. Skeletal muscle FOXO1 (FKHR) transgenic mice have less skeletal muscle mass, down-regulated type I (slow twitch/red muscle) fiber genes, and impaired glycemic control. *J. Biol. Chem.* 279(39): 41114–41123
- Kasuga, M., 1984. Structure and function of the insulin receptor. *Tanpakushitsu*

Kakusan Koso.29(7):533-550

Kazi, A.A., Blonde, L., 2001. Classification of diabetes mellitus. *Clin. Lab. Med.* 21(1):1-13

Kharroubi AT, Darwish HM., 2015. Diabetes mellitus: The epidemic of the century. *World J Diabetes*. 6(6):850-867

Konopka, A. R. *et al.* (2010) 'Molecular adaptations to aerobic exercise training in skeletal muscle of older women', *Journals of Gerontology - Series A Biological Sciences and Medical Sciences*, 65 A(11), pp. 1201–1207

Lenk, K., Schur, R., Linke, A., Erbs, S., Matsumoto, Y., Adams, V., Schuler, G., 2009. Impact of exercise training on myostatin expression in the myocardium and skeletal muscle in a chronic heart failure model. *Eur. J. Heart Fail.* 11(4): 342–348

Little, J.P., Gillen, J.B., Percival, M.E., Safdar, A., Tarnopolsky, M.A., Punthakee, Z., Jung, M.E., Gibala, M.J., 2011. Low-volume high-intensity interval training reduces hyperglycemia and increases muscle mitochondrial capacity in patients with type 2 diabetes. *J Appl Physiol* 111: 1554–1560

Louis, E. *et al.* (2007) 'Time course of proteolytic, cytokine, and myostatin gene expression after acute exercise in human skeletal muscle', *Journal of Applied Physiology*, 103(5), pp. 1744–1751

Machrina, Y., Damanik, H.A.-R., Purba, A., Lindarto, D., 2018. Effect various type of exercise to Insr gene expression, skeletal muscle insulin receptor and insulin

Resistance on Diabetes Mellitus Type-2 model Rats. *Int. J. Heal. Sci.* 6: 50–56

Machrina, Y., Purba, A., Lindarto, D., Maskoen, A.M., 2019. Exercise intensity alter insulin receptor gene expression in diabetic type-2 rat model. *Open Access Maced. J. Med. Sci.* 7(20): 3370–3375

MacInnis, M.J., Gibala, M.J., 2017. Physiological adaptations to interval training and the role of exercise intensity. *J. Physiol.* 595(9): 2915–2930

Mangmool, S., Denkaew, T., Parichatikanond, W., Kurose, H., 2017. β -Adrenergic receptor and insulin resistance in the heart. *Biomol. Ther.* 25(1): 44–56

Mann, S., Beedie, C. and Jimenez, A. (2014) 'Differential effects of aerobic exercise, resistance training and combined exercise modalities on cholesterol and the lipid profile: review, synthesis and recommendations', *Sports Medicine*, 44(2), pp. 211–221

Matsakas, A. *et al.* (2006) 'Effect of swimming on myostatin expression in white and red gastrocnemius muscle and in cardiac muscle of rats', *Experimental Physiology*, 91(6), pp. 983–994

Mayo Clinic, 2021. Type 2 diabetes - Symptoms and causes. Retrived from [https://www.mayoclinic.org/diseases-conditions/type-2 diabetes/symptoms-causes/syc-20351193](https://www.mayoclinic.org/diseases-conditions/type-2-diabetes/symptoms-causes/syc-20351193)

Mayo Clinic, 2019. Diabetes prevention: 5 tips for taking control. Retrived from <https://www.mayoclinic.org/diseases-conditions/type-2-diabetes/in-depth/diabetes-prevention/art-20047639>

- Mayo Clinic, 2011. Type 2 diabetes - Diagnosis and treatment. Retrived from <https://www.mayoclinic.org/diseases-conditions/type-2-diabetes/diagnosis-treatment/drc-20351199>
- McKinnell, I.W., Rudnicki, M.A., 2004. Molecular mechanisms of muscle atrophy. *Cell*. 119(7): 907-910
- McNally, E.M., De Sá Moreira, E., Duggan, D.J., Bönnemann, C.G., Lisanti, M.P., Lidov, H.G.W., Vainzof, M., Passos-Bueno, M.R., Hoffman, E.P., Zatz, M., Kunkel, L.M., 1998. Caveolin-3 in muscular dystrophy. *Hum. Mol. Genet.* 7(5): 871–877
- Melissa Dewulf., 2018. Role of caveolin-3 and caveolae mechanics in muscle pathophysiology. Cellular Biology. *Université Paris-Saclay English*. NNT: 2018SACLS067
- Mengarelli, A., Gentili, A., Strazza, A., Burattini, L., Fioretti, S., Di Nardo, F., 2018. Co-activation patterns of gastrocnemius and quadriceps femoris in controlling the knee joint during walking. *J. Electromyogr.* 42:117-122
- Milan, G., Romanello, V., Pescatore, F., Armani, A., Paik, J.H., Frasson, L., Seydel, A., Zhao, J., Abraham, R., Goldberg, A.L., Blaauw, B., DePinho, R.A., Sandri, M., 2015. Regulation of autophagy and the ubiquitin-proteasome system by the FoxO transcriptional network during muscle atrophy. *Nat. Commun*
- Mirzaeyan, E., Roozbani, M., Shirvani, H., 2020. The Effect of High Intensity Interval Training on the Expression of Enolase3, Caveolin 3 Genes and Plasma Levels of SGOT Enzyme in the Cardiac Tissue of Male Rats with Fatty Liver (Steatosis). *Yafteh* 22(3): 108–117

- Moreno-Villanueva, M. *et al.* (2019) 'Influence of acute exercise on dna repair and parp activity before and after irradiation in lymphocytes from trained and untrained individuals', *International Journal of Molecular Sciences*, 20(12)
- Motiani, K. K. *et al.* (2017) 'Two weeks of moderate-intensity continuous training, but not high-intensity interval training, increases insulin-stimulated intestinal glucose uptake', *Journal of Applied Physiology*, 122(5), pp. 1188–1197
- Muñoz, P., Mora, S., Sevilla, L., Kaliman, P., Tomàs, E., Gumà, A., Testar, X., Palacín, M., Zorzano, A., 1996. Expression and insulin-regulated distribution of caveolin in skeletal muscle: Caveolin does not colocalize with glut4 in intracellular membranes. *J. Biol. Chem.* 271(14): 8133–8139
- Myers, J., Kokkinos, P., Nyelin, E., 2019. Physical activity, cardiorespiratory fitness, and the metabolic syndrome. *Nutrients* 11(7): 1–13
- Nair, A. K. *et al.* (2012) 'Association analysis of common variants in FOXO3 with type 2 diabetes in a South Indian Dravidian population', *Gene*, 491(2), pp. 182–188
- Nayak, N.K., Khedkar, G.D., Khedkar, C.C., Khedkar, C.D., 2015. *Skeletal Muscle*, 1st ed, Encyclopedia of Food and Health. Elsevier Ltd
- Nijhuis, T.H.J., de Boer, S.A.S., Wahegaonkar, A.L., Bishop, A.T., Shin, A.Y., Hovius, S.E.R., Selles, R.W., 2013. A New Approach to Assess the Gastrocnemius Muscle Volume in Rodents Using Ultrasound; Comparison with the Gastrocnemius Muscle Index. *PLoS One*. 10(7):133-144

- Ohsawa, Y., Hagiwara, H., Nakatani, M., Yasue, A., Moriyama, K., Murakami, T., Tsuchida, K., Noji, S., Sunada, Y., 2006. Muscular atrophy of caveolin-3-deficient mice is rescued by myostatin inhibition. *J. Clin. Invest.* 116(14): 2924–2934
- Oshikawa, J., Otsu, K., Toya, Y., Tsunematsu, T., Hankins, R., Kawabe, J., Minamisawa, S., Umemura, S., Hagiwara, Y., Ishikawa, Y., 2004. Insulin resistance in skeletal muscles of caveolin-3-null mice. *Proc Natl Acad Sci U S A.* 101(34):12670-12675
- Park, S., Hong, Yunkyung, Lee, Y., Won, J., Chang, K.-T., Hong, Yonggeun, 2012. Differential expression of caveolins and myosin heavy chains in response to forced exercise in rats. *Lab. Anim. Res.* 28(1):1-9
- Pearson, S.J., 2015. High Intensity Interval Training Vs Moderate Intensity Continuous Training in the Management of Metabolic Type Disease. *MOJ Anat. Physiol.* 1(5):00027
- Peng, S. *et al.* (2015) ‘HDAC2 selectively regulates FOXO3a-mediated gene transcription during oxidative stress-induced neuronal cell death’, *Journal of Neuroscience*, 35(3), pp. 1250–1259
- Perry, B.D., Caldow, M.K., Brennan-Speranza, T.C., Sbaraglia, M., Jerums, G., Garnham, A., Wong, C., Levinger, P., Haq, M.A. ul, Hare, D.L., Price, S.R., Levinger, I., 2016. Muscle atrophy in patients with T2DM role of inflammatory pathways, physical activity and exercise. *Exerc Immunol Rev* 22: 94–109

- Pradhan BS., Prószyński TJ., 2020. A Role for Caveolin-3 in the Pathogenesis of Muscular Dystrophies. *Int J Mol Sci.* 21(22):8736
- Prescott, E., 2018. 18 - Lifestyle Interventions, in: de Lemos, J.A., Omland, T.B.T.-C.C.A.D. (Eds.). Elsevier, pp. 250–269. doi:<https://doi.org/10.1016/B978-0-323-42880-4.00018-2>
- Renna, L.V., Bosè, F., Brigonzi, E., Fossati, B., Meola, G., Cardani, R., 2019. Aberrant insulin receptor expression is associated with insulin resistance and skeletal muscle atrophy in myotonic dystrophies. *PLoS One* 14(3): 1–18
- Roostaei, mehdi, pirani, hossein, rashidlamir, amir, 2020. High intensity interval training induces the expression of Myostatin and Follistatin isoforms in rat muscle: differential effects on fast and slow twitch skeletal muscles. *Med. Lab. J.* 14: 48–53
- Russell, S.J., Schneider, M.F., 2020. Alternative signaling pathways from IGF1 or insulin to AKT activation and FOXO1 nuclear efflux in adult skeletal muscle fibers. *J. Biol. Chem.* 295: 15292–15306.
- Saeedi, P., Petersohn, I., Salpea, P., Malanda, B., Karuranga, S., Unwin, N., Colagiuri, S., Guariguata, L., Motala, A.A., Ogurtsova, K., Shaw, J.E., Bright, D., Williams, R., 2019. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9th edition. *Diabetes Res. Clin. Pract.* 157: 107843
- Sanchez, A.M.J., Candau, R.B., Bernardi, H., 2014. FoxO transcription factors: Their roles in the maintenance of skeletal muscle homeostasis. *Cell. Mol. Life Sci.* 71(9):

1657–1671

Sandri, M. *et al.* (2004) 'Foxo transcription factors induce the atrophy-related ubiquitin ligase atrogin-1 and cause skeletal muscle atrophy', *Cell*, 117(3), pp. 399–412

Schwencke, C., Braun-Dullaeus, R.C., Wunderlich, C., Strasser, R.H., 2006. Caveolae and caveolin in transmembrane signaling: Implications for human disease. *Cardiovasc. Res.* 70(1): 42–49.

Senf, S. M., Dodd, S. L. and Judge, A. R. (2010) 'FOXO signaling is required for disuse muscle atrophy and is directly regulated by Hsp70', *American Journal of Physiology - Cell Physiology*, 298(1), pp. 38–45

Shah, Dinesh S, Nisr, R.B., Stretton, C., Krasteva-christ, G., Hundal, H.S., 2020. Caveolin- 3 de fi ciencia associated with the dystrophy P 104 L mutation impairs skeletal muscle mitochondrial form and function. *J Cachexia Sarcopenia Muscle*. 11(3): 838–858

Shanazari, Z., Faramarzi, M., Banitalebi, E., Hemmati, R., 2019. Effect of moderate and high-intensity endurance and resistance training on serum concentrations of MSTN and IGF-1 in old male Wistar rats. *Horm. Mol. Biol. Clin. Investig.* 38

Shang, L., Chen, T., Deng, Y., Huang, Yiyuan, Huang, Yuanheng, Xian, J., Lu, W., Yang, L., Huang, Q., 2017. Caveolin-3 promotes glycometabolism, growth and proliferation in muscle cells. *PLoS One* 12(12): 1–12

Slopack, D., Roudier, E., Liu, S.T.K., Nwadozi, E., Birot, O., Haas, T.L., 2014. Forkhead BoxO transcription factors restrain exercise-induced angiogenesis. *J. Physiol.* 592(18): 4069–4082

- Srinivasan, K., Viswanad, B., Asrat, L., Kaul, C.L., Ramarao, P., 2005. Combination of high-fat diet-fed and low-dose streptozotocin-treated rat: A model for type 2 diabetes and pharmacological screening. *Pharmacol. Res.* 52(4): 313–320
- Stefanetti, R. J. *et al.* (2018) ‘Recent advances in understanding the role of FOXO3 [version 1; referees: 4 approved]’, *F1000Research*, 7(0), pp. 1–8
- Talukder, M.A.H., Preda, M., Ryzhova, L., Prudovsky, I., Pinz, I.M., 2016. Heterozygous caveolin-3 mice show increased susceptibility to palmitate-induced insulin resistance. *Physiol Rep.* 4(6): 1–14
- Tennant, J.N., Amendola, A., Phisitkul, P., 2014. Endoscopic gastrocnemius release. *Foot Ankle Clin.* 19(4): 787–793.
- Thomas, C.M., Smart, E.J., 2008. Caveolae structure and function: Caveolae Review Series. *J. Cell. Mol. Med.* 12(3): 796–809
- Tosun, B., Selek, O., Gok, U., Tosun, O., 2017. Medial gastrocnemius muscle flap for the reconstruction of unhealed amputation stumps. *J. Wound Care* 26(8): 504–507
- Vermeulen, C., Grade, C., Mantovani, C.S., Alvares, L.E., 2019. Myostatin gene promoter: structure, conservation and importance as a target for muscle modulation. *Anim Sci Biotechnol.* 5: 1–19
- Wang, R., Tian, H., Guo, D., Tian, Q., Yao, T., Kong, X., 2020. Impacts of exercise intervention on various diseases in rats. *J. Sport Heal. Sci.* 9(3): 211–227
- WHO Global Report on Diabetes, 2016. Global Report on Diabetes. *Isbn* 978: 6–86.
- Wu, Y., Ding, Y., Tanaka, Y., Zhang, W., 2014. Risk factors contributing to type 2

diabetes and recent advances in the treatment and prevention. *Int. J. Med. Sci.*

11(11): 1185–1200

Xavier, S., Sadanandan, J., George, N., Paulose, C.S., 2012. β 2-Adrenoceptor and insulin receptor expression in the skeletal muscle of streptozotocin induced diabetic rats: Antagonism by vitamin D 3 and curcumin. *Eur. J. Pharmacol.* 687(1-3): 14–20

Yanling Wu, Yanping Ding, Yoshimasa Tanaka, Wen Zhang, 2014. Risk Factors Contributing to Type 2 Diabetes and Recent Advances in the Treatment and Prevention. *Int. J. Med. Sci.* 11(11):1185-1200

Yasuko Hagiwara, Toshikuni Sasaoka, Kenji Araishi, Michihiro Imamura, Hiroshi Yorifuji, Ikuya Nonaka, Eijiro Ozawa, Tateki Kikuchi, Caveolin-3 deficiency causes muscle degeneration in mice, *Human Molecular Genetics*. 9 (20): 3047–3054

Yoon, S.O., Hyo, J.K., Sung, J.R., Cho, K.A., Young, S.P., Park, H., Kim, M.J., Chang, K.K., Sang, C.P., 2007. Exercise type and muscle fiber specific induction of caveolin-1 expression for insulin sensitivity of skeletal muscle. *Exp. Mol. Med.* 39(3): 395–401

Zheng, L., Rao, Z., Guo, Y., Chen, P., Xiao, W., 2020. High-Intensity Interval Training Restores Glycolipid Metabolism and Mitochondrial Function in Skeletal Muscle of Mice with Type 2 Diabetes. *Front. Endocrinol. (Lausanne)*. 11: 1–14