

## REFERENCE

- A. Ferreira, J., Lennartsson, P. and Taherzadeh, M. (2015) 'Production of ethanol and biomass from thin stillage by *Neurospora intermedia*: A pilot study for process diversification', *Engineering in Life Sciences*, 15. doi: 10.1002/elsc.201400213.
- Adedeji, J. *et al.* (1992) 'Characterization of glycosidically bound aroma compounds in the African mango (*Mangifera indica* L.)', *Journal of Agricultural and Food Chemistry*, 40(4), pp. 659–661. doi: 10.1021/jf00016a028.
- Alberti, F., Foster, G. and Bailey, A. (2017) 'Natural products from filamentous fungi and production by heterologous expression', *Applied Microbiology and Biotechnology*, 101, pp. 493–500. doi: 10.1007/s00253-016-8034-2.
- Almora, K. *et al.* (2004) 'Evaluation of volatiles from ripening papaya (*Carica papaya* L., var. Maradol roja)', *Food Chemistry*, 86(1), pp. 127–130. doi: <https://doi.org/10.1016/j.foodchem.2003.09.039>.
- Argenta, L. C., Fan, X. and Mattheis, J. P. (2003) 'Influence of 1-methylcyclopropene on Ripening, Storage Life, and Volatile Production by d'Anjou cv. Pear Fruit', *Journal of Agricultural and Food Chemistry*, 51(13), pp. 3858–3864. doi: 10.1021/jf034028g.
- Aubert, C. and Milhet, C. (2007) 'Distribution of the volatile compounds in the different parts of a white-fleshed peach (*Prunus persica* L. Batsch)', *Food Chemistry*, 102(1), pp. 375–384. doi: <https://doi.org/10.1016/j.foodchem.2006.05.030>.
- Baardseth, P. *et al.* (1995) 'Evaluation of carrot varieties for production of deep fried carrot chips — II. Sensory aspects', *Food Research International*, 28(6), pp. 513–519. doi: [https://doi.org/10.1016/0963-9969\(95\)00036-4](https://doi.org/10.1016/0963-9969(95)00036-4).
- Bakshi, M., Wadhwa, M. and Makkar, H. (2016) 'Waste to worth: vegetable wastes as animal feed station bulletins; Technical bulletins; Annual reports of

- Ministry of Agriculture of different countries; Annual reports of Ministry of Animal Husbandry of different countries; Annual reports of Ministry of ', *CAB Reviews Perspectives in Agriculture Veterinary Science Nutrition and Natural Resources*, 11, pp. 1–26.
- Bátori, V. *et al.* (2015) 'Ethanol and Protein from Ethanol Plant By-Products Using Edible Fungi *Neurospora intermedia* and *Aspergillus oryzae*.', *BioMed research international*, 2015, p. 176371. doi: 10.1155/2015/176371.
- Berger, R. G. (2007) *Flavours and Fragrances: Chemistry, Bioprocessing and Sustainability*. doi: 10.1007/978-3-540-49339-6.
- Branduardi, P. *et al.* (2007) 'Biosynthesis of vitamin C by yeast leads to increased stress resistance.', *PloS one*, 2(10), p. e1092. doi: 10.1371/journal.pone.0001092.
- Cai, R. *et al.* (2019) 'Antifungal activity and mechanism of citral, limonene and eugenol against *Zygosaccharomyces rouxii*', *LWT*, 106, pp. 50–56. doi: <https://doi.org/10.1016/j.lwt.2019.02.059>.
- Chang, P.-K. *et al.* (2014) 'ASPERGILLUS | Introduction', in Batt, C. A. and Tortorello, M. Lou (eds) *Encyclopedia of Food Microbiology (Second Edition)*. Second Edi. Oxford: Academic Press, pp. 77–82. doi: <https://doi.org/10.1016/B978-0-12-384730-0.00010-0>.
- Chao, L. K. *et al.* (2005) 'Study on the Antiinflammatory Activity of Essential Oil from Leaves of *Cinnamomum osmophloeum*', *Journal of Agricultural and Food Chemistry*, 53(18), pp. 7274–7278. doi: 10.1021/jf0511151u.
- Cheema, A. *et al.* (2018) 'Postharvest hexanal vapor treatment delays ripening and enhances shelf life of greenhouse grown sweet bell pepper (*Capsicum annum* L.)', *Postharvest Biology and Technology*, 136. doi: 10.1016/j.postharvbio.2017.10.006.
- Chen, R. E. and Thorner, J. (2007) 'Function and regulation in MAPK signaling pathways: lessons learned from the yeast *Saccharomyces cerevisiae*.', *Biochimica et biophysica acta*, 1773(8), pp. 1311–1340. doi:

10.1016/j.bbamcr.2007.05.003.

Chin, H.-W. and Lindsay, R. C. (1994) 'Mechanisms of formation of volatile sulfur compounds following the action of cysteine sulfoxide lyases', *Journal of Agricultural and Food Chemistry*, 42(7), pp. 1529–1536. doi: 10.1021/jf00043a026.

Côté, J. *et al.* (2011) 'Antimicrobial effect of cranberry juice and extracts', *Food Control*, 22(8), pp. 1413–1418. doi: <https://doi.org/10.1016/j.foodcont.2011.02.024>.

Dixon, J. and Hewett, E. W. (2000) 'Factors affecting apple aroma/flavour volatile concentration: A Review', *New Zealand Journal of Crop and Horticultural Science*, 28(3), pp. 155–173. doi: 10.1080/01140671.2000.9514136.

Drewnowski, A. (1997) 'Taste preferences and food intake.', *Annual review of nutrition*, 17, pp. 237–253. doi: 10.1146/annurev.nutr.17.1.237.

Drewnowski, A. and Gomez-Carneros, C. (2000) 'Bitter taste, phytonutrients, and the consumer: a review', *The American Journal of Clinical Nutrition*, 72(6), pp. 1424–1435. doi: 10.1093/ajcn/72.6.1424.

Ernst, R., Ejsing, C. S. and Antonny, B. (2016) 'Homeoviscous Adaptation and the Regulation of Membrane Lipids.', *Journal of molecular biology*, 428(24 Pt A), pp. 4776–4791. doi: 10.1016/j.jmb.2016.08.013.

Esparza, I. *et al.* (2020) 'Fruit and vegetable waste management: Conventional and emerging approaches', *Journal of Environmental Management*, 265, p. 110510. doi: <https://doi.org/10.1016/j.jenvman.2020.110510>.

Fan, L. *et al.* (2006) 'Effect of Hexanal Vapor on Spore Viability of *Penicillium expansum*, Lesion Development on Whole Apples and Fruit Volatile Biosynthesis', *Journal of Food Science*, 71, pp. M105–M109. doi: 10.1111/j.1365-2621.2006.tb15632.x.

FAO (2020) *No Title, The International Year of Fruits and Vegetables, 2021, background paper*. doi: <https://doi.org/10.4060/cb2395en>.

- Farewell, A. and Neidhardt, F. C. (1998) 'Effect of temperature on in vivo protein synthetic capacity in *Escherichia coli*.', *Journal of bacteriology*, 180(17), pp. 4704–4710. doi: 10.1128/JB.180.17.4704-4710.1998.
- Ferreira, J. A. *et al.* (2016) 'Waste biorefineries using filamentous ascomycetes fungi: Present status and future prospects.', *Bioresource technology*, 215, pp. 334–345. doi: 10.1016/j.biortech.2016.03.018.
- Forney, C. F. *et al.* (2015) 'Relationships between fruit composition and storage life in air or controlled atmosphere of red raspberry', *Postharvest Biology and Technology*, 110, pp. 121–130. doi: <https://doi.org/10.1016/j.postharvbio.2015.07.017>.
- Forney, C. F., Kalt, W. and Jordan, M. A. (2000) 'The Composition of Strawberry Aroma Is Influenced by Cultivar, Maturity, and Storage', *HortScience HortSci*, 35(6), pp. 1022–1026. doi: 10.21273/HORTSCI.35.6.1022.
- Forney, C. F. and Song, J. (2017) 'Flavors and Aromas: Chemistry and Biological Functions', in *Fruit and Vegetable Phytochemicals*. John Wiley & Sons, Ltd, pp. 515–540. doi: <https://doi.org/10.1002/9781119158042.ch24>.
- Franco, M. R. B., Rodriguez-Amaya, D. and Lancas, F. (2004) 'Volatile composition of three cultivars of mango (*Mangifera indica* L.)', *Food Science and Technology (Campinas)*, 24, pp. 165–169. doi: 10.1590/S0101-20612004000200002.
- Galanakis, C. M. (2012) 'Recovery of high added-value components from food wastes: Conventional, emerging technologies and commercialized applications', *Trends in Food Science & Technology*, 26(2), pp. 68–87. doi: <https://doi.org/10.1016/j.tifs.2012.03.003>.
- Gerday, C. *et al.* (1997) 'Psychrophilic enzymes: a thermodynamic challenge.', *Biochimica et biophysica acta*, 1342(2), pp. 119–131. doi: 10.1016/s0167-4838(97)00093-9.
- Gibbs, B. F. *et al.* (2004) 'Production and characterization of bioactive peptides from soy hydrolysate and soy-fermented food', *Food Research*

*International*, 37(2), pp. 123–131. doi:  
<https://doi.org/10.1016/j.foodres.2003.09.010>.

Gmoser, R. *et al.* (2018) ‘Pigment Production by the Edible Filamentous Fungus *Neurospora Intermedia*’, *Fermentation*, 4. doi:  
[10.3390/fermentation4010011](https://doi.org/10.3390/fermentation4010011).

Gómez, P. and Artés, F. (2004) ‘Improved keeping quality of minimally fresh processed celery sticks by modified atmosphere packaging’, *LWT-Food Science and Technology*, 38, pp. 323–329.

Gomi, K. (2014) ‘ASPERGILLUS | *Aspergillus oryzae*’, in Batt, C. A. and Tortorello, M. Lou (eds) *Encyclopedia of Food Microbiology (Second Edition)*. Second Edi. Oxford: Academic Press, pp. 92–96. doi:  
<https://doi.org/10.1016/B978-0-12-384730-0.00011-2>.

Gouka, R. J., Punt, P. J. and van den Hondel, C. A. (1997) ‘Efficient production of secreted proteins by *Aspergillus*: progress, limitations and prospects.’, *Applied microbiology and biotechnology*, 47(1), pp. 1–11. doi:  
[10.1007/s002530050880](https://doi.org/10.1007/s002530050880).

Griffin, S., Wyllie, S. G. and Markham, J. (1999) ‘Determination of octanol–water partition coefficient for terpenoids using reversed-phase high-performance liquid chromatography’, *Journal of Chromatography A*, 864(2), pp. 221–228. doi: [https://doi.org/10.1016/S0021-9673\(99\)01009-2](https://doi.org/10.1016/S0021-9673(99)01009-2).

Grimm, L. H. *et al.* (2005) ‘Morphology and productivity of filamentous fungi.’, *Applied microbiology and biotechnology*, 69(4), pp. 375–384. doi:  
[10.1007/s00253-005-0213-5](https://doi.org/10.1007/s00253-005-0213-5).

El Hadi, M. A. M. *et al.* (2013) ‘Advances in fruit aroma volatile research’, *Molecules (Basel, Switzerland)*, 18(7), pp. 8200–8229. doi:  
[10.3390/molecules18078200](https://doi.org/10.3390/molecules18078200).

Hakala, M. A., Lapveteläinen, A. T. and Kallio, H. P. (2002) ‘Volatile Compounds of Selected Strawberry Varieties Analyzed by Purge-and-Trap Headspace GC-MS’, *Journal of Agricultural and Food Chemistry*, 50(5), pp. 1133–

1142. doi: 10.1021/jf0111256.

Häkkinen, M. *et al.* (2015) 'The effects of extracellular pH and of the transcriptional regulator PACI on the transcriptome of *Trichoderma reesei*', *Microbial Cell Factories*, 14(1), p. 63. doi: 10.1186/s12934-015-0247-z.

Hammer, K. A., Carson, C. F. and Riley, T. V (2003) 'Antifungal activity of the components of *Melaleuca alternifolia* (tea tree) oil', *Journal of Applied Microbiology*, 95(4), pp. 853–860. doi: <https://doi.org/10.1046/j.1365-2672.2003.02059.x>.

Ho, C. C. (1986) 'Identity and characteristics of *Neurospora intermedia* responsible for oncom fermentation in Indonesia', *Food Microbiology*, 3(2), pp. 115–132. doi: [https://doi.org/10.1016/S0740-0020\(86\)80035-1](https://doi.org/10.1016/S0740-0020(86)80035-1).

Jayanty, S. *et al.* (no date) 'Temporal Relationship between Ester Biosynthesis and Ripening Events in Bananas', *Journal of the American Society for Horticultural Science jashs*, 127(6), pp. 998–1005. doi: 10.21273/JASHS.127.6.998.

Jones, P. G. and Inouye, M. (1996) 'RbfA, a 30S ribosomal binding factor, is a cold-shock protein whose absence triggers the cold-shock response.', *Molecular microbiology*, 21(6), pp. 1207–1218. doi: 10.1111/j.1365-2958.1996.tb02582.x.

Kahle, K. *et al.* (2005) 'On-line gas chromatography combustion/pyrolysis isotope ratio mass spectrometry (HRGC-C/P-IRMS) of major volatiles from pear fruit (*Pyrus communis*) and pear products', *Food Chemistry*, 91(3), pp. 449–455. doi: <https://doi.org/10.1016/j.foodchem.2004.06.026>.

Kansci, G., Koubala, B. B. and Mbome, I. L. (2008) 'Biochemical and Physicochemical Properties of Four Mango Varieties and Some Quality Characteristics of Their Jams', *Journal of Food Processing and Preservation*, 32(4), pp. 644–655. doi: <https://doi.org/10.1111/j.1745-4549.2008.00204.x>.

Klein, D. A. and Paschke, M. W. (2004) 'Filamentous fungi: the indeterminate

lifestyle and microbial ecology.’, *Microbial ecology*, 47(3), pp. 224–235.  
doi: 10.1007/s00248-003-1037-4.

Maarse, H. (2017) *Volatile compounds in foods and beverages*. Routledge.

Machida, M., Yamada, O. and Gomi, K. (2008) ‘Genomics of *Aspergillus oryzae*: learning from the history of Koji mold and exploration of its future’, *DNA research : an international journal for rapid publication of reports on genes and genomes*, 15(4), pp. 173–183. doi: 10.1093/dnares/dsn020.

Makris, D. P. and Rossiter, J. T. (2001) ‘Domestic processing of onion bulbs (*Allium cepa*) and asparagus spears (*Asparagus officinalis*): effect on flavonol content and antioxidant status.’, *Journal of agricultural and food chemistry*, 49(7), pp. 3216–3222. doi: 10.1021/jf001497z.

Mandras, N. *et al.* (2016) ‘Liquid and vapour-phase antifungal activities of essential oils against *Candida albicans* and non-*albicans Candida*’, *BMC Complementary and Alternative Medicine*, 16. doi: 10.1186/s12906-016-1316-5.

Masanetz, C. and Grosch, W. (1998) ‘Key odorants of parsley leaves (*Petroselinum crispum* [Mill.] Nym. ssp. *crispum*) by Odour–activity values’, *Flavour and Fragrance Journal*, 13(2), pp. 115–124. doi: [https://doi.org/10.1002/\(SICI\)1099-1026\(199803/04\)13:2<115::AID-FFJ706>3.0.CO;2-6](https://doi.org/10.1002/(SICI)1099-1026(199803/04)13:2<115::AID-FFJ706>3.0.CO;2-6).

Meyer, V. *et al.* (2016) ‘Current challenges of research on filamentous fungi in relation to human welfare and a sustainable bio-economy: a white paper.’, *Fungal biology and biotechnology*, p. 6. doi: 10.1186/s40694-016-0024-8.

Miermont, A. *et al.* (2013) ‘Severe osmotic compression triggers a slowdown of intracellular signaling, which can be explained by molecular crowding’, *Proceedings of the National Academy of Sciences*. doi: 10.1073/pnas.1215367110.

Misran, A. *et al.* (2014) ‘Composition of phenolics and volatiles in strawberry cultivars and influence of preharvest hexanal treatment on their profiles’,

*Canadian Journal of Plant Science*, 95, p. 140903050330008. doi: 10.4141/CJPS-2014-245.

Multari, S., Stewart, D. and Russell, W. R. (2015) 'Potential of Fava Bean as Future Protein Supply to Partially Replace Meat Intake in the Human Diet', *Comprehensive Reviews in Food Science and Food Safety*, 14(5), pp. 511–522. doi: <https://doi.org/10.1111/1541-4337.12146>.

Muntean, M.-V. *et al.* (2016) 'High Pressure Processing in Food Industry – Characteristics and Applications', *Agriculture and Agricultural Science Procedia*, 10, pp. 377–383. doi: <https://doi.org/10.1016/j.aaspro.2016.09.077>.

Muszewska, A., Pawłowska, J. and Krzyściak, P. (2014) 'Biology, systematics, and clinical manifestations of Zygomycota infections', *European Journal of Clinical Microbiology & Infectious Diseases*, 33(8), pp. 1273–1287. doi: 10.1007/s10096-014-2076-0.

Oshoma, C. *et al.* (2017) 'Fruit Wastes as Substrate For the Production of Amylase by *Aspergillus niger*', *Tropical Journal of Natural Product Research*, 1, pp. 182–185. doi: 10.26538/tjnpr/v1i4.9.

Panesar, R., Kaur, S. and Panesar, P. S. (2015) 'Production of microbial pigments utilizing agro-industrial waste: a review', *Current Opinion in Food Science*, 1, pp. 70–76. doi: <https://doi.org/10.1016/j.cofs.2014.12.002>.

Persson, E. and Halle, B. (2008) 'Cell water dynamics on multiple time scales', *Proceedings of the National Academy of Sciences*, 105(17), pp. 6266–6271. doi: 10.1073/pnas.0709585105.

Pino, J. A. and Mesa, J. (2006) 'Contribution of volatile compounds to mango (*Mangifera indica* L.) aroma', *Flavour and Fragrance Journal*, 21(2), pp. 207–213. doi: <https://doi.org/10.1002/ffj.1703>.

Plazzotta, S., Manzocco, L. and Nicoli, M. C. (2017) 'Fruit and vegetable waste management and the challenge of fresh-cut salad', *Trends in Food Science & Technology*, 63, pp. 51–59. doi:

<https://doi.org/10.1016/j.tifs.2017.02.013>.

- Puupponen-Pimiä, R. *et al.* (2005) 'Bioactive berry compounds-novel tools against human pathogens.', *Applied microbiology and biotechnology*, 67(1), pp. 8–18. doi: 10.1007/s00253-004-1817-x.
- Russell, A. D. (2003) 'Similarities and differences in the responses of microorganisms to biocides', *Journal of Antimicrobial Chemotherapy*, 52(5), pp. 750–763. doi: 10.1093/jac/dkg422.
- Russell, N. (2008) 'Membrane Components and Cold Sensing', in *Psychrophiles: From Biodiversity to Biotechnology*, pp. 177–190. doi: 10.1007/978-3-540-74335-4\_11.
- Sagar, N. A. *et al.* (2018) 'Fruit and Vegetable Waste: Bioactive Compounds, Their Extraction, and Possible Utilization', *Comprehensive Reviews in Food Science and Food Safety*, 17(3), pp. 512–531. doi: <https://doi.org/10.1111/1541-4337.12330>.
- Salles, C. (2006) 'Odour–taste interactions in flavour perception', in *Flavour in Food*, pp. 345–368. doi: 10.1533/9781845691400.3.345.
- Shaw, J. A. *et al.* (1991) 'The function of chitin synthases 2 and 3 in the *Saccharomyces cerevisiae* cell cycle.', *The Journal of cell biology*, 114(1), pp. 111–123. doi: 10.1083/jcb.114.1.111.
- Shu, H. *et al.* (2019) 'Antimicrobial Activity and Proposed Action Mechanism of 3-Carene against *Brochothrix thermosphacta* and *Pseudomonas fluorescens*', *Molecules (Basel, Switzerland)*, 24(18), p. 3246. doi: 10.3390/molecules24183246.
- Silva, C. *et al.* (2013) 'Production of enzymes from *Lichtheimia ramosa* using Brazilian savannah fruit wastes as substrate on solid state bioprocesses', *Electronic Journal of Biotechnology*, 16, p. 9. doi: 10.2225/vol16-issue5-fulltext-7.
- Skoneczny, M. and Skoneczna, A. (2018) 'Stress Response Mechanisms in Fungi'.

- Sues, A. *et al.* (2005) 'Ethanol production from hexoses, pentoses, and dilute-acid hydrolyzate by *Mucor indicus*', *FEMS Yeast Research*, 5(6–7), pp. 669–676. doi: 10.1016/j.femsyr.2004.10.013.
- Tilburn, J. *et al.* (1995) 'The *Aspergillus* PacC zinc finger transcription factor mediates regulation of both acid- and alkaline-expressed genes by ambient pH.', *The EMBO journal*, 14(4), pp. 779–790.
- Tokitomo, Y. *et al.* (2005) 'Odor-active constituents in fresh pineapple (*Ananas comosus* [L.] Merr.) by quantitative and sensory evaluation.', *Bioscience, biotechnology, and biochemistry*, 69(7), pp. 1323–1330. doi: 10.1271/bbb.69.1323.
- Tsanova-Savova, S., Ribarova, F. and Gerova, M. (2005) '(+)-Catechin and (–)-epicatechin in Bulgarian fruits', *Journal of Food Composition and Analysis*, 18(7), pp. 691–698.
- Tsushida, T. and Suzuki, M. (1996) 'Content of Flavonol Glucosides and Some Properties of Enzymes Metabolizing the Glucosides in Onion', *NIPPON SHOKUHIN KAGAKU KOGAKU KAISHI*, 43(5), pp. 642–649. doi: 10.3136/nskkk.43.642.
- USDA (2021) *No Title, Food Availability (Per Capita) Data System*. Available at: <https://www.ers.usda.gov/data-products/food-availability-per-capita-data-system/>.
- Valente, A. *et al.* (2011) 'Ascorbic acid content in exotic fruits: A contribution to produce quality data for food composition databases', *Food Research International*, 44(7), pp. 2237–2242. doi: <https://doi.org/10.1016/j.foodres.2011.02.012>.
- Virgilio, S. *et al.* (2017) 'Regulation of the reserve carbohydrate metabolism by alkaline pH and calcium in *Neurospora crassa* reveals a possible cross-regulation of both signaling pathways', *BMC Genomics*, 18(1), p. 457. doi: 10.1186/s12864-017-3832-1.
- Wakai, S. *et al.* (2017) 'Future insights in fungal metabolic engineering',

- Bioresource Technology*, 245, pp. 1314–1326. doi:  
<https://doi.org/10.1016/j.biortech.2017.04.095>.
- Walker, G. M. and Basso, T. O. (2020) ‘Mitigating stress in industrial yeasts’,  
*Fungal Biology*, 124(5), pp. 387–397. doi:  
<https://doi.org/10.1016/j.funbio.2019.10.010>.
- Wang, C.-Y., Chen, Y.-W. and Hou, C.-Y. (2019) ‘Antioxidant and antibacterial activity of seven predominant terpenoids’, *International Journal of Food Properties*, 22, pp. 229–237. doi: 10.1080/10942912.2019.1582541.
- Werkhoff, P. *et al.* (1998) ‘Vacuum Headspace Method in Aroma Research: Flavor Chemistry of Yellow Passion Fruits’, *Journal of Agricultural and Food Chemistry*, 46(3), pp. 1076–1093. doi: 10.1021/jf970655s.
- Wiesel, I., Rehm, H. J. and Bisping, B. (1997) ‘Improvement of tempe fermentations by application of mixed cultures consisting of *Rhizopus* sp. and bacterial strains’, *Applied Microbiology and Biotechnology*, 47(3), pp. 218–225. doi: 10.1007/s002530050916.
- Wikandari, R. *et al.* (2012) ‘Isolation and characterization of zygomycetes fungi from tempe for ethanol production and biomass applications.’, *Applied biochemistry and biotechnology*, 167(6), pp. 1501–1512. doi: 10.1007/s12010-012-9587-x.
- Yanai, K. *et al.* (1992) ‘Purification of two chitinases from *Rhizopus oligosporus* and isolation and sequencing of the encoding genes’, *Journal of bacteriology*, 174(22), pp. 7398–7406. doi: 10.1128/jb.174.22.7398-7406.1992.
- Yousufi, M. K. (2012) ‘TO DETERMINE PROTEIN CONTENT OF SINGLE CELL PROTEIN PRODUCED BY USING VARIOUS COMBINATIONS OF FRUIT WASTES AND TWO STANDARD FOOD FUNGI’, in.
- Zhang, Jianguo and Zhang, Jining (2016) ‘The filamentous fungal pellet and forces driving its formation.’, *Critical reviews in biotechnology*, 36(6), pp. 1066–1077. doi: 10.3109/07388551.2015.1084262.

Zhang, S.-B. *et al.* (2020) 'Effects of hexanal fumigation on fungal spoilage and grain quality of stored wheat', *Grain & Oil Science and Technology*, 4. doi: 10.1016/j.gaost.2020.12.002.