



UNIVERSITAS
GADJAH MADA

**HPLC CHARACTERIZATION OF AMINO ACIDS EXTRACTION FROM GOLDEN APPLE SNAIL
(Pomacea canaliculata) FLOUR
BY USING KERATIN HYDROLYSATE FROM CHICKEN FEATHER**
RAFI FARHANDY, Dr. Agus Kuncaka, DEA.; Dr.rer.nat. Adhitasari Suratman, S.Si., M.Si.

Universitas Gadjah Mada, 2021 | Diunduh dari <http://etd.repository.ugm.ac.id/>

REFERENCES

- Adelina, 1999, Pengaruh pakan dengan kadar protein dan rasio energy protein berbeda terhadap pertumbuhan bennih ikan bawal air tawar (*cholosomma macropomum*), *Tesis*, Institut Pertanian Bogor, Bogor.
- Agrahari, S., 2013, Production of Extracellular Keratinase Enzymes From *Bacillus Pumilis Sn3* Isolated From Soil Sample Of Ghazipur Poultry Waste Site, *Int. J. Sustain. Dev.*, 2, 2315-4721.
- Alashwal, B. Y., Gupta, A., and Husain, M. S., 2019, Characterization of dehydrated keratin protein extracted from chicken feather, *IOP Conf. Ser.: Matter. Sci. Eng.*, 702, 012033.
- Aldheeb, M. A., Asrar, W., Sulaeman, E., and Omar, A. A., 2016, A review on aerodynamics of Non-Flapping bird wings, *J. Aerosp. Technol. Manag.*, 8(1), 7-17.
- Ali, T. H., Nadia, H. A., and Latifa, A. M., Production, purification, and some properties of extracellular keratinase from feathers degradation by *Aspergillus oryzae* NRRL-447, *J. Appl. Sci. Environ. Sanit.*, 6, 123-136.
- Bishop, C. D., Angus, R. A., and Watts, S. A., 1995, The use of feather meal as a replacement for fish meal in the diet of *Oreochromis niloticus* fry, *Bioresour. Technol.*, 54(3), 291-295.
- Cooper, G., 2000, *The cell*, Washington, ASM Press.
- Edwards, P., and Allan, G. L., 2004, *Feeds and feeding for inland aquaculture in Mekong Region countries*, ACT: Australian Centre for International Agricultural Research, Canberra.
- El-Sayed, A. M., 1999, Alternative dietary protein sources for farmed tilapia *Oreochromis* spp., *Aquaculture*, 179(1-4), 149-168.
- Fauzi, A., 2007, Peningkatan kualitas pupuk organik cair keluaran instalasi biogas fermentasi lanjut dengan penambahan tepung bulu ayam dan tepung silase kepala ikan patin, *Tesis*, Institut Pertanian Bogor, Bogor.
- Funke, A., and Ziegler, F., 2010, Hydrothermal carbonization of biomass: A summary and discussion of chemical mechanisms for process engineering, *Biofuel. Bioprod. Biorefin.*, 4(2), 160–177.



UNIVERSITAS
GADJAH MADA

**HPLC CHARACTERIZATION OF AMINO ACIDS EXTRACTION FROM GOLDEN APPLE SNAIL
(Pomacea canaliculata) FLOUR
BY USING KERATIN HYDROLYSATE FROM CHICKEN FEATHER**
RAFI FARHANDY, Dr. Agus Kuncaka, DEA.; Dr.rer.nat. Adhitasari Suratman, S.Si., M.Si.

Universitas Gadjah Mada, 2021 | Diunduh dari <http://etd.repository.ugm.ac.id/>

- Fowler, L, 1991, Poultry by-product meal as a dietary protein source in fall chinook salmon diets, *Aquac.*, 99(3-4), 309-321.
- Gao, Y., He, J., He, Z., Li, Z., Zhao, B., Mu, Y., Lee, J., and Chu, Z., 2017, Effects of fulvic acid on growth performance and intestinal health of juvenile loach *Paramisgurnus dabryanus* (Sauvage), *Fish Shellfish Immunol.*, 62, 47-56.
- Garrido, T., Lecet, I., Caba, K. D., and Guerrero, P., 2018, Chicken feathers as a natural source of sulphur to develop sustainable protein films with enhanced properties, *Int. J. Biol. Macromol.*, 106, 521-531.
- Ghufran, M. H., and Kordi, K., 2012, *Jurus jitu pengelolaan tambak budi daya perikanan ekonomis*, Lily Publisher, Yogyakarta.
- Ghosh, S., Jung, C., and Meyer-Rochow, V. B., 2017, Snail as mini-livestock: Nutritional potential of farmed *Pomacea Canaliculata* (Ampullariidae), *Agric. Nat. Resour.*, 51(6), 504-511.
- Grazziotin, A., Pimentel F. A., Eyde, J., and Brandelli, A., Nutritional improvement of feather protein by treatment with microbial keratinase, *Anim. Feed. Sci. Technol.*, 126(1-2), 135-144.
- Halwart, M., Litsinger, J., Barrion, A., Viray, M., and Kaule, G., 2012, Efficacy of common carp and nile tilapia as biocontrol agents of rice insect pests in the Philippines, *Int. J. Pest Manag.*, 58(4), 330-346.
- Harrow, B. and Mazur, A., 1954, *Textbook of Biochemistry 6th Ed.*, W.B. Ders. Company, Philadelphia.
- Heidari, M., Dutta, A., Acharya, B., and Mahmud, S., 2018, A review of the current knowledge and challenges of hydrothermal carbonization for biomass conversion, *J. Energy Inst.*, 92(6), 1779-1799.
- Henriksson, P. J. G., Banks, L. K., Suri, S. K., Pratiwi, T. Y., Fatan, N. A., and Troell, M., 2019, Indonesian aquaculture futures-identifying interventions for reducing environmental impacts. *Environ. Res. Lett.*, 14(12), 124062.
- Hickel, E. R., Scheuermann, K. K., and Eberhardt, D. S., 2012, Manejo de caramujos em lavouras de arroz irrigado, em sistema de cultivo pre-germinado, *Agro. Catarinense*, 25(1), 54-57.



UNIVERSITAS
GADJAH MADA

**HPLC CHARACTERIZATION OF AMINO ACIDS EXTRACTION FROM GOLDEN APPLE SNAIL
(*Pomacea canaliculata*) FLOUR
BY USING KERATIN HYDROLYSATE FROM CHICKEN FEATHER**
RAFI FARHANDY, Dr. Agus Kuncaka, DEA.; Dr.rer.nat. Adhitasari Suratman, S.Si., M.Si.

Universitas Gadjah Mada, 2021 | Diunduh dari <http://etd.repository.ugm.ac.id/>

- Huang, Y., Wen, X., Li, S., Xuan, X., and Zhu, D., 2016, Effects of protein levels on growth, feed utilization, body composition, amino acid composition and physiology indices of Juvenile chu's croaker, *Nibea coibo*, *Aquac. Nutr.*, 23(3), 594-602.
- Jawa, G., Suwitari, N. K., and Sanjaya, I. G., 2020, The appearance of broiler that given golden apple snail meat (*Pomacea canaliculata*) flour as replacement of fish flour, *SEAS (Sustainable Environment Agricultural Science)*, 4(1), 60-64.
- Kaushik, S. J., and Seiliez, I., 2010, Protein and amino acid nutrition and metabolism in fish: Current knowledge and future needs, *Aquac. Res.*, 41(3), 322-332.
- Ketaren, N., 2008, Pemanfaatan limbah bulu ayam sebagai sumber protein ayam pedaging dalam pengelolaan lingkungan hidup, *Tesis*, Universitas Sumatera Utara, Medan.
- Kuncaka, A., Arvianto, R. I., Latifa, A. S. R. B., Rambe, M. R., Suratman, A., and Triono, S., 2021, Analysis and characterization of solid and liquid organic fertilizer from hydrothermal carbonization (HTC) of chicken feather and blood waste, *Indones. J. Chem.*, 21(3), 651.
- Mazotto, A. M., Coelho, R. R. R., Cedrola, S. M. L., de Lima, M. F., Couri, S., de Souza, E. P., and Vemelho, A. B., 2011, Keratinase production by three *Bacillus* spp. using feather meal and whole feather as substrate in a submerged fermentation, *Enzyme Res.*, 2011, 1-7.
- Merkel, J.A., 1981, *Managing livestock wastes*, AVI Publishing Company Inc., Westport.
- Mousavi, S., Salouti, M., Shapoury, R., and Heidari, Z., Research article: Optimization of keratinase production for feather degradation by *Bacillus subtilis*, *Jundishapur J. Microbiol.*, 6, 7160.
- Mulia, D. S., Yuliningsih, R. T., Maryanto, H., dan Purbomartono, C., 2016, Pemanfaatan limbah bulu ayam menjadi bahan pakan ikan dengan fermentasi *Bacillus subtilis*, *J. People Environ.*, 23(1), 49-57.
- Nursinatrio and Nugroho, R.A., 2019, Hydrolyzed chicken feather meal as protein source for red tilapia (*Oreochromis* sp.) aquafeeds, *Pak. J. Zool.*, 51(4), 1489-1496.



UNIVERSITAS
GADJAH MADA

**HPLC CHARACTERIZATION OF AMINO ACIDS EXTRACTION FROM GOLDEN APPLE SNAIL
(Pomacea canaliculata) FLOUR
BY USING KERATIN HYDROLYSATE FROM CHICKEN FEATHER**
RAFI FARHANDY, Dr. Agus Kuncaka, DEA.; Dr.rer.nat. Adhitasari Suratman, S.Si., M.Si.

Universitas Gadjah Mada, 2021 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Peddu, J. L., Chitturi, C. M. K., and Lakshmi, M. M., 2013, Efficient degradation of feather by keratinase producing *Bacillus* sp., *Int. J. Microbiol.*, 2, 608321.

Pertiwi, M. P., and Saputri, D. D., 2020, Golden apple snail (*Pomacea canaliculata*) as an alternative protein source in Pasupati catfish (*Pangasius* sp.) fish feed, *Nusantara Biosci.*, 2(12), 162-167.

Puastuti, W., 2007, Teknologi pemrosesan bulu ayam dan pemanfaatannya sebagai sumber protein pakan ruminansia, *Wartazoa*, 17(2), 53-60.

Purmaningsih, A., 2010, Pengaruh penambahan tepung keong mas (*Pomacea canaliculata*) dalam ransum terhadap kualitas telur itik, *Tesis*, Universitas Sebelas Maret, Surakarta.

Puspitasari, D., 2010, Pengaruh penambahan tepung keong mas (*Pomacea canaliculata Lamarck*) dalam ransum terhadap performan produksi itik petelur, *Tesis*, Universitas Sebelas Maret, Surakarta.

Said, M.I., 2014, *By product ternak terknologi dan aplikasinya*, *Tesis* Institut Pertanian Bogor, Bogor.

Sandjojo, H., Hasan, H., dan Dewantoro, E., 2014, Pemanfaatan tepung keong mas (*Pomacea canaliculata*) sebagai bahan substitusi tepung ikan dalam pakan terhadap keragaan pertumbuhan ikan nila gift (*Oreochromis niloticus*), *Tesis*, Universitas Muhammadiyah Pontianak, Pontianak.

Setalaphruk, C., and Price, L.L., 2007, Children's traditional knowledge of wild food resources: a case study in rural village in Northeast Thailand, *J. Ethnobiol. Ethnomed.*, 3, 33.

Sidik, M., 2016, Karakteristik kimiawi tepung bulu limbah pengolahan kerupuk kulit sapi menggunakan NaOH dan lama perendaman berbeda, *Tesis*, Universitas Hasanuddin, Makassar.

Simamora, A., 2015, Asam mino, Peptida, dan Protein, *Tesis*, Universitas Kristen Krida Wacana, Jakarta.

Tacon, A.G.J., 1993. *Feed ingredients for warmwater fish. Fish meal and other processed feedstuffs*, FAO: Food and Agriculture Organization, Rome.



UNIVERSITAS
GADJAH MADA

**HPLC CHARACTERIZATION OF AMINO ACIDS EXTRACTION FROM GOLDEN APPLE SNAIL
(Pomacea canaliculata) FLOUR
BY USING KERATIN HYDROLYSATE FROM CHICKEN FEATHER**
RAFI FARHANDY, Dr. Agus Kuncaka, DEA.; Dr.rer.nat. Adhitasari Suratman, S.Si., M.Si.

Universitas Gadjah Mada, 2021 | Diunduh dari <http://etd.repository.ugm.ac.id/>

- Tarmizi, A., 2001, Evaluasi nilai nutrisi tepung bulu yang difermentasi dengan menggunakan bacillus licheniformis pada ayam broiler, *Tesis*, Institut Pertanian Bogor, Bogor.
- Wan, M. Y., Dong, G., Yang, B. Q., and Feng, H., Identification and characterization of a novel antioxidant peptide from feather keratin hydrolysate, *Biotechnol. Lett.*, 38(4), 643-9.
- Wang, B., Yang, W., Mckittrick, J., and Meyers, M. A., 2016, Keratin: structure, mechanical properties, occurrence in biological organisms, and efforts at bioinspiration, *Prog. Mater Sci.*, 76, 229-318.
- Wu, R. B., Wu, C. L., Liu, D., Yang, X. H., Huang, J. F., Zhang, J., Liao, B., He, H. L., and Li, H., Overview of antioxidant peptides derived from marine resources: the sources, characteristic, purification, and evaluation methods, *Appl. Biochem. Biotechnol.*, 176(7), 1815-33.
- Wijayanti, R., Wibowo, L., and Solikhin, 2016, Pengaruh varietas padi (*Oriza sativa L.*) dan jenis kelamin keong emas (*Pomacea sp.*) terhadap daya rusak keong emas pada tanaman padi, *J. Agrotek Tropika*, 4(2), 141-145.
- Yahav Spitzer, R., Mau, V., and Gross, A., 2018, Using hydrothermal carbonization for sustainable treatment and reuse of human excreta, *J. Clean. Prod.*, 205, 955-963.
- Zhao, P., Ge, S., Ma, D., Areeprasert, C., and Yoshikawa, K., 2014, Effect of hydrothermal pretreatment on convective drying characteristics of paper sludge, *ACS Sustain. Chem. Eng.*, 2(4), 665-671.