

DAFTAR PUSTAKA

- Adham, M., Kurniawan, A.N., Muhtadi, A.I., Roezin, A., Hermani, B., Gondhowiardjo, S., Tan, I.B., Middeldorp, J.M. (2012) 'Nasopharyngeal carcinoma in indonesia: Epidemiology, incidence, signs, and symptoms at presentation', *Chinese Journal of Cancer*, 31(4), pp. 185–196. doi: 10.5732/cjc.011.10328.
- Armstrong, R. W., Yu, M. C., and Henderson, B. E. (1983) 'Salted Fish and Inhalants as Risk Factors for Nasopharyngeal Carcinoma in Malaysian Chinese', *Cancer Research*, 43(6), pp. 2967–2970.
- Arnold, M., Wildeman, M.A., Visser, O., Karim-Kos, K.E., Middeldrop, J.M., Fles, R., Tan, I.B., Coebergh, J.W. (2013) 'Lower mortality from nasopharyngeal cancer in the Netherlands since 1970 with differential incidence trends in histopathology', *Oral Oncology*, 49(3), pp. 237–243. doi: 10.1016/j.oraloncology.2012.09.016.
- Autexier, C. and Lue, N. F. (2006) 'The Structure and Function of Telomerase Reverse Transcriptase', *Annual Review of Biochemistry*, 75(1), p. 493–517. doi: 10.1146/annurev.biochem.75.103004.142412.
- Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan Republik Indonesia. (2013) 'Riset Kesehatan Dasar (RISKESDAS) 2013', *Laporan Nasional 2013*, pp. 1–384. doi: 1 Desember 2013.
- Barnes, L., Eveson, J.W., Reichart, P., Sidransky, D. (2003) 'CHAPTER 2 Tumours of the Nasopharynx', *WHO classification of tumours. Pathology & Genetics: head and Neck Tumors*, pp. 82–106.
- Bei, J. X., Jia, W. H., and Zeng, Y. X. (2012) 'Familial and large-scale case-control studies identify genes associated with nasopharyngeal carcinoma', *Seminars in Cancer Biology*. Elsevier Ltd, 22(2), pp. 96–106. doi: 10.1016/j.semcancer.2012.01.012.
- Bei, J. X., Su, W., Ng, C., Yu, K., Chin, Y., Lou, P., Hsu, W.L., McKay, J.D., Chen, C.J., Chang, Y.S., Chen, L.Z., Chen, M.Y., Cui, Q., Feng, F.T., Feng, Q.S., Guo, Y.M., Jia, W.H., Khoo, A.S., Liu, W.S., Mo, H.Y., Pua, K.C., Teo, S.H., Tse, K.P., Xia, Y.F., Zhang, H., Zhou, G.Q., Liu, J.J., Zeng, Y.X., Hildesheim, A. (2015) 'A GWAS meta-analysis and replication study identifies a novel locus within CLPTM1L/TERT associated with nasopharyngeal carcinoma in individuals of Chinese ancestry', *Cancer Epidemiology Biomarkers and Prevention*, 25(1), pp. 188–192. doi: 10.1158/1055-9965.EPI-15-0144.

- Bray, F., Haugen, M., Moger, T.A., Tretli, S., Aalen, O.O., Grotmol, T. (2008) 'Age-incidence curves of nasopharyngeal carcinoma worldwide: Bimodality in low-risk populations and aetiologic implications', *Cancer Epidemiology Biomarkers and Prevention*, 17(9), pp. 2356–2365. doi: 10.1158/1055-9965.EPI-08-0461.
- Brennan B. (2006) 'Nasopharyngeal carcinoma', *Orphanet Journal of Rare Diseases*, (1), p. 23. doi:10.1186/1750-1172-1-23.
- Cao, Y., Miao, X.P., Huang, M.Y., Deng, L., Hu, L.F., Ernberg, I., Zeng, Y.X., Lin, D.X., Shao, J.Y. (2006) 'Polymorphisms of XRCC1 genes and risk of nasopharyngeal carcinoma in the Cantonese population.', *BMC cancer*, 6, p. 167. doi: 10.1186/1471-2407-6-167.
- Cao, S. M., Liu, Z., Jia, W.H., Huang, Q.H., Liu, Q., Guo, X., Huang, T.B., Ye, W., Hong, M.H. (2011) 'Fluctuations of epstein-barr virus serological antibodies and risk for nasopharyngeal carcinoma: A prospective screening study with a 20-year follow-up', *PLoS ONE*, 6(4), pp. 1–10. doi: 10.1371/journal.pone.0019100.
- Chang, E. T. and Adami, H. O. (2006) 'The enigmatic epidemiology of nasopharyngeal carcinoma', *Cancer Epidemiology Biomarkers and Prevention*, 15(10), pp. 1765–1777. doi: 10.1158/1055-9965.EPI-06-0353.
- Charles, I. A., Ogbolosingha, A. J., and Afia, I. U. (2017) 'Health risk assessment of instant noodles commonly consumed in Port Harcourt, Nigeria', *Environmental Science and Pollution Research*. doi: 10.1007/s11356-017-0583-0.
- Chen, Z., Pan, S., Sun, Q., Yao, K. (1984) 'Renewal of nasopharyngeal squamous epithelium of rats treated with N,N'-Dinitrosopiperazine', *Bulletin of Hunan Medical College*, 9(4), pp.317-323.
- Cho, E., Hildesheim, A., Chen, C., Hsu, M., Chen, I., Mittl, B.F., Levine, P.H., Liu, M.Y., Chen, J.Y., Brinton, L.A., Cheng, Y.J., Yang, C.S. (2003) 'Nasopharyngeal Carcinoma and Genetic Polymorphisms of DNA Repair Enzymes XRCC1 and hOGG1 Nasopharyngeal Carcinoma and Genetic Polymorphisms of DNA Repair', *Prevention*, 12(October), pp. 1100–1104.
- Di Maso, M., Talamini, R., Bosetti, C., Montella, M., Zucchetto, A., Libra, M., Negri, E., Levi, F., La Vecchia, C., Franceschi, S., Serraino, D., Polesel, J. (2013) 'Red meat and cancer risk in a network of case-control studies focusing on cooking practices', *Annals of Oncology*, 24(12), pp. 3107–3112. doi: 10.1093/annonc/mdt392.
- Diggs, Deacquita L., Huderson, Ashley C., Harris, Kelly L., Myers, Jeremy N., Banks, Leah D., Rekhadevi, Perumalla V., Niaz, Mohammad S., Ramesh,

- Aramandla. (2011) 'Polycyclic Aromatic Hydrocarbons and digestive tract cancer – a perspective,' (1), pp. 1–31. doi: 10.1080/10590501.2011.629974.Polycyclic.
- Fachiroh, J., Sangrajang, S., Johansson, M., Renard, H., Gaborieau, V., Chabrier, A., Chindavijak, S., Brennan, P., McKay, J.D. (2012) 'Tobacco consumption and genetic susceptibility to nasopharyngeal carcinoma (NPC) in Thailand', *Cancer Causes and Control*, 23(12), pp. 1995–2002. doi: 10.1007/s10552-012-0077-9.
- Fang, M. Z. (2003) 'Tea polyphenol (-)-epigallocatechin-3-gallate inhibits DNA methyltransferase and reactivates methylation-silenced genes in cancer cell lines', *Cancer Res.*, 63, pp. 7563–7570.
- Fareed, M. and Afzal, M. (2013) 'Single nucleotide polymorphism in genome-wide association of human population: A tool for broad spectrum service', *Egyptian Journal of Medical Human Genetics*, 14(2), pp. 123–134. doi: 10.1016/j.ejmhg.2012.08.001.
- Food and Agriculture Organization of the United Nations. (2013) Current Worldwide Annual Meat Consumption per capita, Livestock and Fish Primary Equivalent.
- Guo, X., Johnson, R.C., Deng, H., Liao, J., Guan, L., Nelson, G.W., Tang, M., Zheng, Y., de The, G., O'Brien, S.J., Winkler, C.A., Zeng, Y. (2009) 'Evaluation of non-viral risk factors for nasopharyngeal carcinoma in a high-risk population of Southern China', *Int J Cancer*, 124(12), pp. 2942–4947. doi: 10.1038/nbt.3121.ChIP-nexus.
- Handoyo, D. and Rudiretna, A. (2001) 'Prinsip Umum dan Pelaksanaan Polymerase Chain Reaction (PCR) [General Principles and Implementation of Polymerase', 9(1), pp. 17–29.
- Hildesheim, A., Anderson, L.M., Chen, C.J., Cheng, Y.J., Brinton, L.A., Daly, A.K., Reed, C.D., Chen, I.H., Caporaso, N.E., Hsu, M.M., Chen, J.Y., Idle, J.R., Hoover, R.N., Yang, C.S., Chhabra, S.K. (1997) 'CYP2E1 genetic polymorphisms and risk of nasopharyngeal carcinoma in Taiwan', *JNCI: Journal of the National Cancer Institute*, 89(16), pp. 1207–1212, <https://doi.org/10.1093/jnci/89.16.1207>
- Hildesheim, A. and Wang, C. (2012) 'Genetic predisposition factors and nasopharyngeal carcinoma risk: A review of epidemiological association studies, 2000–2011: Rosetta Stone for NPC: Genetics, viral infection, and other environmental factors', *Semin. Cancer Biol*, Available at: <https://dx.doi.org/10.1016/j.semcancer.2012.01.007>
- <https://www.cancer.org/cancer/nasopharyngeal-cancer/detection-diagnosis->

staging/staging.html

- IARC Working Group on the Evaluation of Carcinogenic Risk to Humans. (2012) IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, No. 100E : Personal Habits and Indoor Combustions. Vol. 100E. International Agency for Research on Cancer. Lyon, France. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK304384/>
- Jackson, L. S. and Al-Taher, F. (2005) 'Effects of consumer food preparation on acrylamide formation', *Advances in Experimental Medicine and Biology*, 561, pp. 447–465. doi: 10.1007/0-387-24980-X_34.
- James, M. A., Wen, W., Wang, Y., Byers, L.A, Heymach, J.V., Coombeset, K.R., Girard, L., Minna, J., You, M. (2012) 'Functional characterization of CLPTM1L as a lung cancer risk candidate gene in the 5p15.33 locus', *PLoS ONE*, 7(6), pp. 1–9. doi: 10.1371/journal.pone.0036116.
- Jemal, A., Bray, F., Center, M.M., Ferlay, J., Ward, E., Forman, D. (2011) 'Global Cancer Statistics: 2011', *CA: A Cancer Journal for Clinicians*, 61(2), pp. 69–90. doi: 10.3322/caac.20107. Available.
- Jia, W.H., Luo, X.Y., Feng, B.J., Ruan, H.L., Bei, J.X., Liu, W.S., Qin, H.D., Feng, Q.S., Chen, L.Z., Yao, S.Y., Zeng, Y.X. (2010) 'Traditional Cantonese Diet and Nasopharyngeal Carcinoma Risk: A Large-Scale Case-Control Study in Guangdong, China', *BMC cancer*, 10, p. 446. doi: 10.1186/1471-2407-10-446.
- Jia, W.-H. and Qin, H.-D. (2012) 'Non-viral environmental risk factors for nasopharyngeal carcinoma: A systematic review', *Seminars in Cancer Biology*. Elsevier Ltd, 22(2), pp. 117–126. doi: 10.1016/j.semcancer.2012.01.009.
- Kelsey, J.L., Whittemore, A.S., Evans, A.S., Thompson, W.D. (1996) *Methods in Observational Epidemiology*. Oxford University Press.
- Koonanuwachaidet, P. (1984) 'N-Nitroso Compounds: Environmental Carcinogens', *J. Sci. Soc. Thailand*, pp. 207–220. doi: 10.2306/scienceasia1513-1874.1984.10.207.
- Luka, J., Kallin, B., and Klein, G. (1979) 'Induction of the Epstein-Barr virus (EBV) cycle in latently infected cells by n-butyrate', *Virology*, 94(1), pp. 228–231. doi: 10.1016/0042-6822(79)90455-0.
- Luo, X., Lamsal, L.P., Xu, W., Lu, J., Lu, Y., Shen, Y., Guan, Q. (2014) 'Genetic variant in CLPTM1L confers reduced risk of lung cancer: a replication study in Chinese and a meta-analysis.', *Asian Pacific journal of cancer prevention : APJCP*, 15(21), pp. 9241–7. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25422207>.

Mafra, I., Amaral, J.S., and Oliveira M.B. (2010) *Polycyclic aromatic hydrocarbons (PAH) in olive oils and other vegetable oils: potential for carcinogenesis*. Victor R. Preedy and Ronald Ross Watson, editors. *Olives and Olive Oil in Health and Disease Prevention*, pp. 489–498.

McKay, J.D., Hung, R.J., Gaborieau, V., Boffetta, P., Chabrier, A., Byrnes, G., Zaridze, D., Mukeria, A., Szeszenia-Dabrowska, N., Lissowska, J., Rudnai, P., Fabianova, E., Mates, D., Bencko, V., Foretova, L., Janout, V., McLaughlin, J., Shepherd, F., Montpetit, A., Narod, S., Krokhan, H.E., Skorpene, F., Elvestad, M.B., Vatten, L., Njølstad, I., Axelsson, T., Chen, C., Goodman, G., Barnett, M., Loomis, M.M., Lubinski, J., Matyjasik, J., Lener, M., Oszutowska, D., Field, J., Liloglou, T., Xinarianos, G., Cassidy, A., EPIC Study., Vineis, P., Clavel-Chapelon, F., Palli, D., Tumino, R., Krogh, V., Panico, S., González, CA., Ramón Quirós, J., Martínez, C., Navarro, C., Ardanaz, E., Larrañaga, N., Kham, K.T., Key, T., Bueno-de-Mesquita, H.B., Peeters, P.H., Trichopoulou, A., Linseisen, J., Boeing, H., Hallmans, G., Overvad, K., Tjønneland, A., Kumle, M., Riboli, E., Zelenika, D., Boland, A., Delepine, M., Foglio, M., Lechner, D., Matsuda, F., Blanche, H., Gut, I., Heath, S., Lathrop, M., Brennan, P. (2008) ‘Lung cancer susceptibility locus at 5p15.33’, *Lung Cancer*, 40(12), pp. 1404–1406. doi: 10.1038/ng.254.Lung.

Mocellin, S., Verdi, D., Pooley, K.A., Landi, M.T., Egan, K.M., Baird, D.M., Prescott, J., De Vivo, I., Nitti, D. (2012) ‘Telomerase reverse transcriptase locus polymorphisms and cancer risk: A field synopsis and meta-analysis’, *Journal of the National Cancer Institute*, 104(11), pp. 840–854. doi: 10.1093/jnci/djs222.

Nazar-Stewart, V., Vaughan, T.L., Burt, R.D., Chen, C., Berwick, M., Swanson, G.M. (1999) ‘Glutathione S-transferase M1 and susceptibility to nasopharyngeal carcinoma’, *Cancer Epidemiol Biomarkers Prev*, 8(6), pp. 547–551.

Parada, H., Steck, S.E., Bradshaw, P.T., Engel, L.S., Conway, K., Teitelbaum, S.L., Neugut, A.I., Santella, R.M., Gammon M.D. (2017) ‘Grilled, Barbecued, and Smoked Meat Intake and Survival Following Breast Cancer’, *Journal of the National Cancer Institute*, 109(6), pp. 1–8. doi: 10.1093/jnci/djw299.

Pfeifer, G. P., Denissenko, M.F., Olivier, M., Tretyakova, N., Hecht, S.S., Hainaut, P. (2002) ‘Tobacco smoke carcinogens, DNA damage and p53 mutations in smoking-associated cancers’, *Oncogene*, 21(48), pp. 7435–7451. doi: 10.1038/sj.onc.1205803.

Rostkowska, K., Zwierz, K., Rózański, A., Moniuszko-Jakoniuk, J., Roszczenko

- A. (1998) 'Formation and metabolism of N-nitrosamines', *Polish Journal of Environmental Studies*, 7(6), pp. 321–325.
- Soeripto. (1998) 'Epidemiology of nasopharyngeal carcinoma', *Berita Majalah Kedokteran Masyarakat*, XIII, pp. 207-211.
- Sugimura, T. (2000) 'Nutrition and dietary carcinogens', *Carcinogenesis*, 21, pp. 387-395.
- Suhda, S. (2015) Polimorfisme Gen *Cytochrome* P450 Family 2 Subfamili EPolipeptida 1 (CYPE21) dan Kebiasaan Merokok sebagai Faktor Risiko Karsinoma Nasofaring di Yogyakarta. Thesis. Fakultas Kedokteran Universitas Gadjah Mada.
- Sun, X., Tong, L., Wang, Y., Wu, Y., Sheng, H., Lu, L., Wang, W. (2011) 'Can global variation of nasopharynx cancer be retrieved from the combined analyses of IARC cancer information (CIN) databases?', *PLoS ONE*, 6(7), pp. 1–9. doi: 10.1371/journal.pone.0022039.
- Tang, M., Zeng Y., Poisson, A., Marti, D., Guan, L., Zheng, Y., Deng, H., Liao, J., Guo, X., Sun, S., Nelson, G., de Thé, G., Winkler, CA., O'Brien, SJ., Carrington, M., Gao, X. (2010) 'Haplotype-dependent HLA susceptibility to nasopharyngeal carcinoma in a Southern Chinese population', *Genes and immunity*, 11(4), pp 334-342. doi:10.1038/gene.2009.109.
- Tsao, S. W., Yip, Y.L., Tsang, C.M., Pang, P.S., Lau, V.M.Y., Zhang, G., Lo, K.W. (2014) 'Etiological factors of nasopharyngeal carcinoma', *Oral Oncology*. Elsevier Ltd, 50(5), pp. 330–338. doi: 10.1016/j.oraloncology.2014.02.006.
- Van Hassel H.J., and Topping, J.W. (1977) 'Trotter's syndrome. A review'. *Oral Surg. Oral Med. Oral Pathol.* 44 (1): pp.125–127. PMID 267870
- World Cancer Research Fund, American Institute for Cancer Research. (2007) Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective. Cancer Research. p. 517.
- WHO-IARC. (2012) Globocan 2012 : estimated cancer incidence, mortality and prevalence worldwide in 2012. Available at : http://globocan.iarc.fr/Pages/fact_sheets_population.aspx
- Xie, S., Yu, I.T., Tse, L.A., Au, J.S.K., Wang, F., Lau, J.S., Zhang, B. (2014) 'Domestic Incense Burning and Nasopharyngeal Carcinoma : A Case-Control Study in HongKong Chinese', *Environmental and Molecular Mutagenesis*, 55, pp.751-756
- Xue, W. Q., Qin, H.D., Ruan, H.L., Shugart, Y.Y., Jia, W.H. (2013) 'Quantitative association of tobacco smoking with the risk of nasopharyngeal carcinoma: A comprehensive meta-analysis of studies conducted between 1979 and

- 2011', *American Journal of Epidemiology*, 178(3), pp. 325–338. doi: 10.1093/aje/kws479.
- Xun, X., Wang, H., Yang, H., Wang, H., Wang, B., Kang, L., Jin, T., Chen, C. (2014) 'CLPTM1L Genetic Polymorphisms and Interaction With Smoking and Alcohol Drinking in Lung Cancer Risk', *Medicine*, 93(28), p. e289. doi: 10.1097/MD.0000000000000289.
- Yee Ko, J. M., Dai, W., Wun Wong, E.H., Kwong, D., Tong Ng, W., Lee, A., Kai Cheong Ngan, R., Chung Yau, C., Tung, S., Li Lung, M. (2014) 'Multigene pathway-based analyses identify nasopharyngeal carcinoma risk associations for cumulative adverse effects of TERT-CLPTM1L and DNA double-strand breaks repair', *International Journal of Cancer*, 135(7), pp. 1634–1645. doi: 10.1002/ijc.28802.
- Ye, S., Dhillon, S., Ke, X., Collins, A.R., Day, I.N.M. (2001) 'An efficient procedure for genotyping single nucleotide polymorphisms', *Nucl. Acid Res*, 29(17), p. e88
- Yin, J., Wang L., Zheng L., Wang X., Shi Y., Shao A., Ding G., Liu C., Chen S., Tang W., Gu H. (2014) 'TERT-CLPTM1L Rs401681 C>T polymorphism was associated with a decreased risk of esophageal cancer in a chinese population', *PLoS ONE*, 9(7), pp. 3–9. doi: 10.1371/journal.pone.0100667.
- Young, L.S., Arrand, J.R., and Murray, P.G. (2007) EBV gene expression and regulation. In: Arvin A, Campadelli-Fiume G, Mocarski E, Moore P.S., Roizman, B., Whitley, R., Yamanishi, K., editors. *Human Herpesviruses: Biology, Therapy, and Immunoprophylaxis*. Cambridge: Cambridge University Press; Chapter 27. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK47431/>
- Yu, M.C., Chi-chun, Mo., Wei-xiong, Chong., Fu-sun, Yeh., and Brian, E Henderson. (1988) 'Preserved Foods and Nasopharyngeal Carcinoma: A Case-Control Study in Guangxi, China Preserved Foods and Nasopharyngeal Carcinoma: A Case-Control Study in', pp. 1954–1959.
- Yuan, J.M., Wang, X.L., Xiang, Y.B., Gao, Y.T., Ross, R.K., Yu, M.C. (2000) 'Preserved foods in relation to risk of nasopharyngeal carcinoma in Shanghai, China', *International Journal of Cancer*, 85(3), pp. 358–363. doi: 10.1002/(SICI)1097-0215(20000201)85:3<358::AID-IJC11>3.0.CO;2-E.
- Zhang, Y., Zhang, X., Zhang, H., Zhai, Y., Wang, Z., Li, P., Yu, L., Xia, X., Zhang, Y., He, F., Zhou, G. (2015) 'Common variations in TERT-CLPTM1L locus are reproducibly associated with the risk of nasopharyngeal carcinoma in Chinese populations', *Oncotarget*, 7(1), pp. 759-770

Zheng, X., Luo, Y., Christensson, B., Drettner, B. (1994) 'Induction of nasal and nasopharyngeal tumours in Sprague-Dawley rats fed with Chinese salted fish', *Acta Otolaryngol*, 114, pp.98–104

Zou, X.N., Lu, S.H., and Liu, B. (1994) 'Volatile N-nitrosamines and their precursors in Chinese salted fish—a possible etiological factor for NPC in china', *Int J Cancer*, 59, pp.155–158.