

DAFTAR PUSTAKA

- Abdel-Misih, S. R. Z. And Bloomston, M. (2010) 'Liver anatomy.', *The Surgical clinics of North America*, 90(4), pp. 643–653. Doi: 10.1016/j.suc.2010.04.017.
- Attwa, M. H. And El-Etreby, S. A. (2015) 'Guide for diagnosis and treatment of hepatocellular carcinoma', *World journal of hepatology*, 7(12), pp. 1632–1651. Doi: 10.4254/wjh.v7.i12.1632.
- Baues, C. *Et al.* (2019) 'Volumetric assessment of mediastinal lymphoma masses in Hodgkin lymphoma.', *Leukemia & lymphoma*, 60(13), pp. 3244–3250. Doi: 10.1080/10428194.2019.1623888.
- Beaumont, H. *Et al.* (2015) 'Changes of lung tumour volume on CT - prediction of the reliability of assessments', *Cancer imaging: the official publication of the International Cancer Imaging Society*, 15, p. 17. Doi: 10.1186/s40644-015-0052-2.
- Bonekamp, S. *Et al.* (2011) 'Hepatocellular Carcinoma: arsse to TACE Assessed with Semiautomated Volumetric and Functional Analysis of Diffusion-weighted and Contrast-enhanced MR Imaging Data', *Radiology*, 260(3), pp. 752–761. Doi: 10.1148/radiol.11102330.
- Bray, F. *Et al.* (2018) 'Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries.', *CA: a cancer journal for clinicians*, 68(6), pp. 394–424. Doi: 10.3322/caac.21492.
- Carlsson, G., Gullberg, B. And Hafström, L. (1983) 'Estimation of liver tumor volume using different formulas - an experimental study in rats.', *Journal of cancer research and clinical oncology*, 105(1), pp. 20–23. Doi: 10.1007/BF00391826.
- Cassinotto, C., Aubé, C. And Dohan, A. (2017) 'Diagnosis of hepatocellular carcinoma: An update on international guidelines', *Diagnostic and Interventional Imaging*, 98(5), pp. 379–391. Doi: <https://doi.org/10.1016/j.diii.2017.01.014>.
- Chalian, H. *Et al.* (2011) 'Radiologic assessment of response to therapy: comparison of RECIST Versions 1.1 and 1.0.', *Radiographics: a review publication of the Radiological Society of North America, Inc*, 31(7), pp. 2093–2105. Doi: 10.1148/rg.317115050.
- Chalian, H. *Et al.* (2012) 'Hepatic tumors: region-of-interest versus volumetric analysis for quantification of attenuation at CT.', *Radiology*, 262(3), pp. 853–861. Doi: 10.1148/radiol.11110106.
- Dachman, A. H. *Et al.* (2001) *Tumor Size on Computed Tomography Scans Is One Measurement Enough?*
- Dinkel, J. *Et al.* (2013) 'Inter-observer reproducibility of semi-automatic tumor diameter measurement and volumetric analysis in patients with lung cancer.', *Lung cancer (Amsterdam, Netherlands)*, 82(1), pp. 76–82. Doi: 10.1016/j.lungcan.2013.07.006.
- Eisenhauer EA, Therasse P, Bogaerts J, et al. (2009) 'New response evaluation criteria in solid tumours: re-vised RECIST guideline (version 1.1)', *Eur J Cancer*, 45(2), pp. 228–247.
- Erasmus, J. J. *Et al.* (2003) 'Interobserver and intraobserver variability in measurement of non-small-cell carcinoma lung lesions: implications for assessment of tumor response.', *Journal of clinical oncology: official journal of the American Society*

- Universitas Gadjah Mada, 2021 | Diunduh dari <http://etd.repository.ugm.ac.id/>
of *Clinical Oncology*, 21(13), pp. 2574–2582. Doi: 10.1200/JCO.2003.01.144.
- Fuentes, D. *Et al.* (2019) ‘Automated Volumetric Assessment of Hepatocellular Carcinoma Response to Sorafenib: A Pilot Study.’, *Journal of computer assisted tomography*, 43(3), pp. 499–506. Doi: 10.1097/RCT.0000000000000866.
- Ghouri, Y. A., Mian, I. And Rowe, J. H. (2017) ‘Review of hepatocellular carcinoma: Epidemiology, etiology, and carcinogenesis’, *Journal of carcinogenesis*, 16, p. 1. Doi: 10.4103/jcar.jcar_9_16.
- Gomaa, A. I. *Et al.* (2009) ‘Diagnosis of hepatocellular carcinoma’, *World journal of gastroenterology*, 15(11), pp. 1301–1314. Doi: 10.3748/wjg.15.1301.
- Gonzalez-Guindalini, F. D. *Et al.* (2013) ‘Assessment of liver tumor response to therapy: role of quantitative imaging.’, *Radiographics: a review publication of the Radiological Society of North America, Inc*, 33(6), pp. 1781–1800. Doi: 10.1148/rg.336135511.
- Gregory, J. *Et al.* (2020) ‘Evaluation of liver tumour response by imaging.’, *JHEP reports: innovation in hepatology*, 2(3), p. 100100. Doi: 10.1016/j.jhepr.2020.100100.
- Hayes, S. A. *Et al.* (2016) ‘Comparison of CT volumetric measurement with RECIST response in patients with lung cancer’, *European journal of radiology*. 2016/01/02, 85(3), pp. 524–533. Doi: 10.1016/j.ejrad.2015.12.019.
- Höink, A. J. *Et al.* (2017) ‘Response Evaluation of Malignant Liver Lesions After TACE/SIRT: Comparison of Manual and Semi-Automatic Measurement of Different Response Criteria in Multislice CT.’, *rofo: Fortschritte auf dem Gebiete der Röntgenstrahlen und der Nuklearmedizin*, 189(11), pp. 1067–1075. Doi: 10.1055/s-0043-116220.
- Imber, B. S. *Et al.* (2019) ‘Comparison of Radiographic Approaches to Assess Treatment Response in Pituitary Adenomas: Is RECIST or RANO Good Enough?’, *Journal of the Endocrine Society*, 3(9), pp. 1693–1706. Doi: 10.1210/js.2019-00130.
- Juza, R. M. And Pauli, E. M. (2014) ‘Clinical and surgical anatomy of the liver: a review for clinicians.’, *Clinical anatomy (New York, N.Y.)*, 27(5), pp. 764–769. Doi: 10.1002/ca.22350.
- Kanwal, F. *Et al.* (2011) ‘Increasing Prevalence of HCC and Cirrhosis in Patients With Chronic Hepatitis C Virus Infection’, *Gastroenterology*, 140(4), pp. 1182–1188.e1. Doi: <https://doi.org/10.1053/j.gastro.2010.12.032>.
- Keil, S. *Et al.* (2010) ‘Semiautomated versus manual evaluation of liver metastases treated by radiofrequency ablation.’, *Journal of vascular and interventional radiology: JVIR*, 21(2), pp. 245–251. Doi: 10.1016/j.jvir.2009.10.024.
- Kharbach, A. *Et al.* (2017) ‘Towards a Novel Approach for Tumor Volume Quantification’, *Journal of Imaging*, 3, p. 41. Doi: 10.3390/jimaging3040041.
- Kim, H. J. And Kim, W. (2012) ‘Method of tumor volume evaluation using magnetic resonance imaging for outcome prediction in cervical cancer treated with concurrent chemotherapy and radiotherapy’, *Radiation oncology journal*. 2012/06/30, 30(2), pp. 70–77. Doi: 10.3857/roj.2012.30.2.70.
- Kobayashi, S. *Et al.* (2020) ‘Intraarterial and intravenous contrast enhanced CT and MR imaging of multi-step hepatocarcinogenesis defining the early stage of hepatocellular carcinoma development’, *Hepatoma Research*, 6, p. 36. Doi: 10.20517/2394-5079.2020.24.
- Kudo, M. (2010) ‘Will Gd-EOB-MRI Change the Diagnostic Algorithm in Hepatocellular Carcinoma?’, *Oncology*, 78(suppl 1(Suppl. 1), pp. 87–93. Doi:

- Kudo, M. *Et al.* (2019) ‘Response Evaluation Criteria in Cancer of the Liver version 5 (RECICL 2019 revised version).’, *Hepatology research : the official journal of the Japan Society of Hepatology*, 49(9), pp. 981–989. Doi: 10.1111/hepr.13394.
- Levine, Z. H. *Et al.* (2010) ‘RECIST versus volume measurement in medical CT using ellipsoids of known size.’, *Optics express*, 18(8), pp. 8151–8159. Doi: 10.1364/OE.18.008151.
- Levine, Z. H. *Et al.* (2011) ‘Tumor volume measurement errors of RECIST studied with ellipsoids.’, *Medical physics*, 38(5), pp. 2552–2557. Doi: 10.1118/1.3577602.
- Loho, I. M. *Et al.* (2016) ‘Hepatocellular Carcinoma in a Tertiary Referral Hospital in Indonesia: Lack of Improvement of One-Year Survival Rates between 1998-1999 and 2013-2014.’, *Asian Pacific journal of cancer prevention : APJCP*, 17(4), pp. 2165–2170. Doi: 10.7314/apjcp.2016.17.4.2165.
- Ltd, P. H. P. (2021) *Predible*.
- M. Sopiudin Dahlan (2013) *Statistik Untuk Kedokteran dan Kesehatan: Deskriptif, Bivariat, dan Multivariat Dilengkapi Aplikasi Menggunakan SPSS*. Edisi 6 ce. Jakarta: Epidemiologi Indonesia.
- Mardian, Y. *Et al.* (2019) “‘Sarcopenia and intramuscular fat deposition are associated with poor survival in Indonesian patients with hepatocellular carcinoma: a retrospective study””, *BMC Gastroenterology*, 19. Doi: 10.1186/s12876-019-1152-4.
- Mcglynn, K. A., Petrick, J. L. And London, W. T. (2015) ‘Global epidemiology of hepatocellular carcinoma: an emphasis on demographic and regional variability.’, *Clinics in liver disease*, 19(2), pp. 223–238. Doi: 10.1016/j.cld.2015.01.001.
- MEVIS, F. (2021) *About MEVIS*.
- Mittal, S. And El-Serag, H. B. (2013) ‘Epidemiology of hepatocellular carcinoma: consider the population’, *Journal of clinical gastroenterology*, 47 Suppl(0), pp. S2–S6. Doi: 10.1097/MCG.0b013e3182872f29.
- Moawad, A. W. *Et al.* (2020) ‘Feasibility of Automated Volumetric Assessment of Large Hepatocellular Carcinomas’ Responses to Transarterial Chemoembolization.’, *Frontiers in oncology*, 10, p. 572. Doi: 10.3389/fonc.2020.00572.
- Mozley, P. D. *Et al.* (2012) ‘Measurement of tumor volumes improves RECIST-based response assessments in advanced lung cancer’, *Translational oncology*. 2012/02/01, 5(1), pp. 19–25. Doi: 10.1593/tlo.11232.
- Patella, F. *Et al.* (2019) ‘Assessment of the response of hepatocellular carcinoma to interventional radiology treatments.’, *Future oncology (London, England)*, 15(15), pp. 1791–1804. Doi: 10.2217/fon-2018-0747.
- Pereira, N. *Et al.* (2014) ‘Hemivaginal septum resection in a patient with a rare variant of Herlyn-Werner-Wunderlich syndrome.’, *Journal of minimally invasive gynecology*, 21(6), pp. 1113–1117. Doi: 10.1016/j.jmig.2014.05.001.
- Van Persijn van Meerten, E. L., Gelderblom, H. And Bloem, J. L. (2010) ‘RECIST revised: implications for the radiologist. A review article on the modified RECIST guideline.’, *European radiology*, 20(6), pp. 1456–1467. Doi: 10.1007/s00330-009-1685-y.
- Pupulim, L. F. *Et al.* (2018) ‘Volumetric measurement of hepatic tumors: Accuracy of manual contouring using CT with volumetric pathology as the reference method.’, *Diagnostic and interventional imaging*, 99(2), pp. 83–89. Doi: 10.1016/j.diii.2017.11.002.

- Universitas Gadjah Mada, 2021 | Diunduh dari <http://etd.repository.ugm.ac.id/>
- Ronot, M. *Et al.* (2014) 'Alternative Response Criteria (Choi, European association for the study of the liver, and modified Response Evaluation Criteria in Solid Tumors [RECIST]) Versus RECIST 1.1 in patients with advanced hepatocellular carcinoma treated with sorafenib.', *The oncologist*, 19(4), pp. 394–402. Doi: 10.1634/theoncologist.2013-0114.
- Rothe, J. H. *Et al.* (2013) 'Size determination and response assessment of liver metastases with computed tomography--comparison of RECIST and volumetric algorithms.', *European journal of radiology*, 82(11), pp. 1831–1839. Doi: 10.1016/j.ejrad.2012.05.018.
- Schiavon, G. *Et al.* (2012) 'Tumor Volume as an Alternative Response Measurement for Imatinib Treated GIST Patients', *plos ONE*, 7(11). Doi: 10.1371/journal.pone.0048372.
- Singal, A. G., Lampertico, P. And Nahon, P. (2020) 'Epidemiology and surveillance for hepatocellular carcinoma: New trends', *Journal of Hepatology*, 72(2), pp. 250–261. Doi: 10.1016/j.jhep.2019.08.025.
- Suzuki, C. *Et al.* (2008) 'Radiologic measurements of tumor response to treatment: practical approaches and limitations.', *Radiographics : a review publication of the Radiological Society of North America, Inc*, 28(2), pp. 329–344. Doi: 10.1148/rg.282075068.
- Suzuki, K. *Et al.* (2011) 'Quantitative radiology: automated CT liver volumetry compared with interactive volumetry and manual volumetry.', *AJR. American journal of roentgenology*, 197(4), pp. W706-12. Doi: 10.2214/AJR.10.5958.
- Tacher, V. *Et al.* (2013) 'Semiautomatic volumetric tumor segmentation for hepatocellular carcinoma: comparison between C-arm cone beam computed tomography and MRI.', *Academic radiology*, 20(4), pp. 446–452. Doi: 10.1016/j.acra.2012.11.009.
- Tacher, V. *Et al.* (2016) 'Comparison of Existing Response Criteria in Patients with Hepatocellular Carcinoma Treated with Transarterial Chemoembolization Using a 3D Quantitative Approach.', *Radiology*, 278(1), pp. 275–284. Doi: 10.1148/radiol.2015142951.
- Tanwar, S. *Et al.* (2020) 'Inflammation and fibrosis in chronic liver diseases including non-alcoholic fatty liver disease and hepatitis C.', *World journal of gastroenterology*, 26(2), pp. 109–133. Doi: 10.3748/wjg.v26.i2.109.
- Therasse, P. *Et al.* (2000) 'New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada.', *Journal of the National Cancer Institute*, 92(3), pp. 205–216. Doi: 10.1093/jnci/92.3.205.
- Tirkes, T. *Et al.* (2013) 'Response criteria in oncologic imaging: review of traditional and new criteria.', *Radiographics : a review publication of the Radiological Society of North America, Inc*, 33(5), pp. 1323–1341. Doi: 10.1148/rg.335125214.
- Tirumani, S. H. *Et al.* (2016) 'Accuracy and feasibility of estimated tumour volumetry in primary gastric gastrointestinal stromal tumours: validation using semiautomated technique in 127 patients', *European radiology*. 2015/05/20, 26(1), pp. 286–295. Doi: 10.1007/s00330-015-3829-6.
- Usermanualfujifilm (2019) 'Synapse 3D: User manual Software Version V5.4US'. West Avenue Stamford, CT 06902, U.S.A.: FUJIFILM MEDICAL SYSTEMS, U.S.A, pp. 99, 412.
- Vietti Violi, N. *Et al.* (2019) 'Radiological Diagnosis and Characterization of HCC.', in

- Waller, L. P., Deshpande, V. And Pysopoulos, N. (2015) ‘Hepatocellular carcinoma: A comprehensive review’, *World journal of hepatology*, 7(26), pp. 2648–2663. Doi: 10.4254/wjh.v7.i26.2648.
- Welsh, J. L. *Et al.* (2012) ‘Comparison of response evaluation criteria in solid tumors with volumetric measurements for estimation of tumor burden in pancreatic adenocarcinoma and hepatocellular carcinoma.’, *American journal of surgery*, 204(5), pp. 580–585. Doi: 10.1016/j.amjsurg.2012.07.007.
- Winter, K. S. *Et al.* (2018) ‘Towards volumetric thresholds in RECIST 1.1: Therapeutic response assessment in hepatic metastases.’, *European radiology*, 28(11), pp. 4839–4848. Doi: 10.1007/s00330-018-5424-0.
- Yamamichi, J. *Et al.* (2020) ‘Assessment of tumor volume and density as a measure of the response of advanced hepatocellular carcinoma to sorafenib: Application of automated measurements on computed tomography scans.’, *JGH open : an open access journal of gastroenterology and hepatology*, 4(2), pp. 145–152. Doi: 10.1002/jgh3.12230.
- Zhao, B. *Et al.* (2013) ‘Exploring intra- and inter-reader variability in uni-dimensional, bi-dimensional, and volumetric measurements of solid tumors on CT scans reconstructed at different slice intervals.’, *European journal of radiology*, 82(6), pp. 959–968. Doi: 10.1016/j.ejrad.2013.02.018.