

DAFTAR PUSTAKA

- Antara, I.K.Y., Wiargitha, I.K., Mahadewa, T.G.B. (2017). Validitas New Injury Severity Score (NISS) dalam mendeteksi terjadinya koagulopati pada pasien multiple trauma. *Jurnal Bedah Nasional*, 10 (1): 15-19.
- Azab, B., Camacho-Rivera, M., Taioli, E. (2014). Average Values and Racial Differences of Neutrophil Lymphocyte Ratio among a Nationally Representative Sample of United States Subjects. *Plos One*, 9 (11): 1-6.
- Baker, S.P., O'Neill, B., Haddon, W., Long, W.B. (1974). The Injury Severity Score: a method for describing patients with multiple injuries and evaluating emergency care. *J Trauma*; 14 (8):187-196.
- Betensley, A., Sharif, R., Karamichos, D. (2017). A Systematic Review of the Role of Dysfunctional Wound Healing in the Pathogenesis and Treatment of Idiopathic Pulmonary Fibrosis. *J. Clin. Med*, 6 (2): 1-19.
- Bolorunduro O.B., Villegas C., Oyetunji, Haut, E.R., Stevens, K.A., Chang, D.C., Cornwell, E.E., Efron, D.T., Haider, A.H. (2011). Validating the Injury Severity Score (ISS) in different populations: ISS predicts mortality better among Hispanics and females. *J Surg Res.*; 166(1): 40-44.
- Charalampidis, C., Youroukou, A., Lazaridis, G., Baka, S., Mpoukovinas, I., Karavasilis, V., Kioumis, I., Pitsion, G., Papaiwannou, A., Karavergou, A., et al. (2015). Pleura space anatomy. *J. Thorac.Dis.*, 7 (S1): S27-S32.
- Cho, H., Hur, H.W., Kim, S.W., Kim, S.H., Kim, J.H., Kim, Y.T., Lee, K. (2009). Pre-treatment neutrophil to lymphocyte ratio is elevated in epithelial ovarian cancer and predicts survival after treatment. *Cancer Immunol. Immunother.*, 58: 15-23.
- Darmawan. (2016). Peran rasio neutrofil limfosit sebagai prediktor major adverse events tujuh hari dalam perawatan pada pasien sindrom koroner akut. *Tesis*. Fakultas Kedokteran Universitas Indonesia. Jakarta.
- Di Bartolomeo, S., Sanson, G., Nardi, G., Scian, F., Michelutto, V., Lattuada, L. (2001). A population-based study on pneumothorax in severely traumatized patients. *J. Trauma*, 51(4): 677-682.
- Dixon, J.B., O'Brien, P.E. (2006). Obesity and the White Blood Cell Count: Changes with Sustained Weight Loss. *Obesity Surgery*, 16: 251-257.
- Duchesne, J.C., Tatum, D., Jones, G., Davis, B., Robledo, R., DeMoya, M., O'Keeffe, T., Ferrada, P., Jacome, T., Schroll, R., Wlodarczyk, J., Prakash, P., Smith, B., Inaba, K., Khor, D., Duke, M., Khan, M. (2017). Multi-institutional analysis of neutrophil-to-lymphocyte ratio (NLR) in patients with severe hemorrhage: A new mortality predictor value. *J Trauma Acute Care Surg*; 83 (5): 888-893.

- Fisher, A., Srikusalanuku, W., Fisher, L., Smith, P. (2016). The Neutrophil to Lymphocyte Ratio on Admission and Short-Term Outcomes in Orthogeriatric Patients. *International Journal of Medical Sciences*; 13(8): 588-602. doi: 10.7150/ijms.15445.
- Forget, P., Khalifa, C., Defour, J.P., Latinne, D., Pel, M.C.V., De Kock, M. (2017). What is the normal value of the neutrophil-to-lymphocyte ratio?. *BMC Res. Notes*, 10 (12): 1-4.
- Graziosi, L., Marino, E., De Angelis, V., Rebonato, A., Cavazzoni, E., Donini, A., (2015). Prognostic value of preoperative neutrophils to lymphocytes ratio in patients resected for gastric cancer. *The American Journal of Surgery*, 209 (2): 333-337.
- Gwak, M.S., Choi, S.J., Kim, J.A, Ko, J.S., Kim, T.H., Lee, S.M., Park, J., Kim., M.H. (2007). Effects of Gender on White Blood Cell Populations and Neutrophil-Lymphocyte Ratio Following Gastrectomy in Patients with Stomach Cancer. *J. Korean Med. Sci.*, 22 (Suppl): S104-S108.
- Hussain, M., Babar, M.Z.M., Akhtar, L., Hussain, M.S. (2017). Neutrophil lymphocyte ratio (NLR): A well assessment tool of glycemic control in Type-2 diabetic patients. *Pak. J. Med. Sci.*, 33 (6): 1366-1370
- Idrees, M.M., Ingleby, A.M., Wali, S.O. (2003). Evaluation and management of pneumothorax. *Saudi Med. J.*, 24 (5): 447-452.
- Jilma, B., Blann, A., Pernerstorfer, T., Stohlawetz, P., Eichler, H.G., Vondrovec, B., Amiral, J., Richter, V., Wagner, O.F. (1999). Regulation of Adhesion Molecules during Human Endotoxemia. No Acute Effects of Aspirin. *AM. J. Respi. Care Med.*, 159: 857-863.
- Katleya, F., Anam, M.S., Dadiyanto, D.W. (2015). Rasio Jumlah Neutrofil Limfosit pada Awal Masuk Rawat sebagai Faktor Risiko Luar Pnemonia Anak. *Sari Pediatri*, 17 (1): 47-51.
- Lavoie, A., Moore, L., LeSage, N., Liberman, M., Sampalis, J.S. (2005). The Injury Severity Score or the New Injury Severity Score for predicting intensive care unit admission and hospital length of stay?. *Injury, Int. J. Care Injured*; 36: 477—483.
- Linn, S., Linn, R., Sheps, S., Sarid, M., Michaelson, M., Geva, H., Wiener, M., Brandes, J.M., & Revach, M. (1993). Injury Severity Scoring and Length of Stay in Hospital of War Casualties—Demonstration of an Association and Possible Selection Bias. *International Journal of Epidemiology*; 22 (5): 905-910.
- McKnight, C.L., Burns, B. (2020). *Pneumothorax*. StatPearls Publishing. Download: <https://pubmed.ncbi.nlm.nih.gov/28722915/>.
- Missaoui, L., Hassine, H., Ayari, A., Dhahri, B., Baccar, M.A., Azzebi, S., Amar, J.B., Aouina, H. (2016). What factors influence the duration of pneumothorax drainage?. *European Respiratory Journal* 2016; 48: PA5048; DOI: 10.1183/13993003.congress-2016.PA5048

- Muttaqien, F., Bermansyah, Saleh, I. (2019). Pengaruh durasi pneumotorak terhadap tingkat stress oksidatif paru tikus wistar. *Qanun Medika*, 3 (1): 45-53.
- Noppen, M., De Keukeleire, T. (2008). Pneumothorax. *Respiration*, 76 (2): 121-127.
- Ozkaya, M., Bedel, C. (2018). Can neutrophil to lymphocyte ratio and platelet to lymphocyte ratio predict the severity of sternum fractures?. *Current Thoracic Surgery*, 3 (3): 116-122.
- Paramanathan, A., Saxena, A., Morris, D.L. (2014). A systematic review and meta-analysis on the impact of pre-operative neutrophil lymphocyte ratio on long term outcomes after curative intent resection of solid tumours. *Surgical Oncology*, 23: 31-39.
- Petridou, E.T., & Antonopoulos, C.N. (2017). *Injury Epidemiology. International Encyclopedia of Public Health*, 258–274. doi:10.1016/b978-0-12-803678-5.00233-2.
- Pusponegoro A.D., 2005. Luka. Dalam: Sjamsuhidajat R, De Jong W, penyunting. Buku Ajar Ilmu Bedah. Edisi ke-2. Jakarta: EGC: 66-88.
- Riandy, M.D., Purba, B.A., Puspita, Y., Zainal, R., Siswo, L. (2019). Hubungan Rasio Neutrofil Limfosit dengan Mortalitas ≤ 28 hari di General Intensive Care Unit RSUP Dr. Mohammad Hoesin Palembang. *Jurnal Anestesi Perioperatif*, 7 (3): 188-197.
- Rodriguez, J.A.O., Hipskind, J.E. (2020). Iatrogenic pneumothorax. *Statpearls*. Download dari: <https://www.ncbi.nlm.nih.gov/books/NBK526057/>
- Safitri, W.R. (2016). Analisis korelasi Pearson dalam menentukan hubungan antara kejadian demam berdarah dengue dengan kepadatan penduduk di kota surabaya pada tahun 2012-2014. *Jurnal Ilmu Keperawatan*, 2 (2).
- Sakurai, K., Chubachi, S., Irie, H., Tsutsumi, A., Kameyama, N., Kamatani, T., Koh, H., Terashima, T., Nakamura, H., Asano, K., Betsuyaku, T. (2018). Clinical utility of blood neutrophil lymphocyte ratio in Japanese COPD patients. *BMC Pulmonary Medicine*, 18 (65): 1-11.
- Saliccioli, J.D., Marshall, D.C., Pimentel, M.A.F., Santos, M.D., Pollard, T., Celi, L.A., Shalhoub, J. (2015). The association between the neutrophil to lymphocyte ratio and mortality in critical illness: an observational cohort study. *Critical Care*, 19 (13): 1-8.
- Salim, Caroline. (2015). Sistem Penilaian Trauma. *Cermin Dunia Kedokteran*-232, 42 (9): 702-209.
- Saricam, M., Ozkan, B., Turk, Y. (2018). Management of traumatic pneumothorax in isolated blunt chest trauma. *Eur. Res. J.*, 1-5.
- Sharma, A., Jindal, P. (2008). Principles of diagnosis and management of traumatic pneumothorax. *J. Emerg. Trauma Shock*, 1 (1): 34-41.
- Simandjuntak, U.R.D. (2018). Rasio Platelet Limfosit (PLR) dan Rasio Neutrofil Limfosit (NLR) Pre Treatment Sebagai Faktor Prognostik Pasien Kanker Rektum Stadium II dan IV di RSUP H. Adam Malik Medan Tahun 2011-2013. Download dari Repositori Institusi USU, Universitas Sumatera Utara. <http://repositori.usu.ac.id/handle/123456789/8208>.

- Slobodan, M., Marko, S., Bojan, M. (2015). Pneumothorax - diagnosis and treatment. *Sanamed*, 10 (3): 221-228.
- Soehnlein, O. (2012). Multiple Roles for Neutrophils in Atherosclerosis. *Circulation Research*, 875-888.
- Soulaiman, S.E., Dopa, D., Raad, A.T., Hasan, W., Ibrahim, N., Hasan, A., Sulaiman, H.A., Darwich, M. (2020). Cohort retrospective study: the neutrophil to lymphocyte ratio as an independent predictor of outcomes at the presentation of the multi-trauma patient. *International Journal of Emergency Medicine*, 13 (5): 1-5.
- Sudiarta, I.P.G., Wiargitha, I.K., Mahadewa, T.G.B. (2020). Perbedaan nilai Neutrophil Lymphocyte Ratio (NLR) terhadap pemeriksaan kultur darah dalam mendiagnosis sepsis pada pasien peritonitis di RSUP Sanglah, Bali, Indonesia. *Intisari Sains Medis*, 11 (1): 165-171.
- Toptas, M., Samanci, N.S., Akkoc, E., Yucetas, E., Cebeci, E., Sen, O., Can, M.M., and Ozturk, S. (2018). Factors Affecting the Length of Stay in the Intensive Care Unit: Our Clinical Experience. *BioMed Research International Volume 2018*. Article ID 9438046: 1-4. <https://doi.org/10.1155/2018/9438046>
- Webster, J., Scuffham, P., Sherriff, K.L., Stankiewicz, M., Chaboyer, W.P. (2012). Negative pressure wound therapy for skin grafts and surgical wounds healing by primary intention. *Cochrane Database of Systematic Reviews*; 4:1-45.
- Widjaya, D.P., Amin, Z., Suprayitno, Afifi, R., Shatri, H. (2014). Karakteristik dan Faktor-Faktor yang Mempengaruhi Kesintasan Pasien Pneumotoraks di Rumah Sakit Cipto Mangunkusumo, Jakarta. *Ina. J. Chest Crit. and Emerg. Med*; 1 (3): 113-119.
- Yadollahi, M., Ghiassee, A., Anvar, M., Ghaem, H., Farahmand, M. (2017). Analysis of Shahid Rajaei hospital administrative data on injuries resulting from car accidents in Shiraz, Iran: 2011-2014 data. *Chinese Journal of Traumatology*; 20: 27-33. <http://www.elsevier.com/locate/CJTEE>
- Yen, S.T. (2016). Correlation of Injury Severity Score and Average Length of Stay. *Health Care Current Reviews*; 4:2 (Suppl). <http://dx.doi.org/10.4172/2375-4273.C1.018>