

DAFTAR PUSTAKA

- Abdul-Ghani, M. A. and DeFronzo, R. A. (2010). Pathogenesis of insulin resistance in skeletal muscle. *Journal of Biomedicine and Biotechnology*, 2010 : 1–19.
- Abe, A. E., De Oliveira, C. E., Dalboni, T. M., Chagas-Paula, D. A., Rocha, B. A., De Oliveira, R. B., *et al.* (2015). Anti-inflammatory sesquiterpene lactones from *Tithonia diversifolia* trigger different effects on human neutrophils. *Revista Brasileira de Farmacognosia*, 25 : 111–116.
- ADA. (2020). Classification and diagnosis of diabetes: Standards of Medical Care in Diabetes-2020. *Diabetes Care*, 43 (Supplement 1) : S14–S31.
- ADA. (2021). 2. Classification and diagnosis of diabetes: Standards of medical care in diabetes-2021. *Diabetes Care*, 44 (Supplement 1) : S15–S33.
- Aditama, T. Y. (2014). *Jamu dan Kesehatan*. Jakarta: Lembaga Penerbit Balitbangkes (LPB).
- Ajao, A. A. and Moteetee, A. N. (2017). *Tithonia diversifolia* (Hemsl) A. Gray. (Asteraceae: Heliantheae), an invasive plant of significant ethnopharmacological importance: A review. *South African Journal of Botany*. Elsevier B.V., 396–403.
- Alberti, K. G. M. M. and Zimmet, P. Z. (1998). Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: Diagnosis and classification of diabetes mellitus. Provisional report of a WHO consultation. *Diabetic Medicine*, 15 (7).
- Antunes, L. C., Elkfury, J. L., Jornada, M. N., Foletto, K. C. and Bertoluci, M. C. (2016). Validation of HOMA-IR in a model of insulin-resistance induced by a high-fat diet in Wistar rats. *Archives of Endocrinology and Metabolism*, 60 (2) : 138–142.
- Anwar, K., Hariadi, R. E. P., Kamalia, N., Santoso, H. B. and Ngindra, A. P. L. (2016). Perbandingan Efek Ekstrak Etanol, Fraksi N-Butanol, dan Fraksi Petroleum Eter Daun Kembang Bulan (*Tithonia diversifolia* (Hemsley) A. Gray) Terhadap Penurunan Kadar Glukosa Darah Mencit Jantan Yang Diinduksi Aloksan. *Jurnal Pharmascience*, 03 (02) : 80–88.

- Armstrong, R. B. and Phelps, R. O. (1984). Muscle fiber type composition of the rat hindlimb. *American Journal of Anatomy*, 171 (3) : 259–272.
- Au, T. H., Skarbek, C., Pethe, S., Labruere, R., Baltaze, J.-P., Phuong, T., *et al.* (2021). Structural modification and biological activity studies of tagitinin C and its derivatives. *Tetrahedron*.
- Balitbangkes. (2019). *Laporan Nasional RISKESDAS 2018*. Lembaga Penerbit Badan Penelitian dan Pengembangan Kesehatan (LPB), Kementerian Kesehatan Republik Indonesia.
- Broca, C., Gross, R., Petit, P., Sauvaire, Y., Manteghetti, M., Tournier, M., *et al.* (1999). 4-hydroxyisoleucine: Experimental evidence of its insulinotropic and antidiabetic properties. *American Journal of Physiology - Endocrinology and Metabolism*, 277 (4 40-4).
- Cecilia, E. O., Gbadura, A. I., Olalekan, E. O. and Kundu, R. (2017). Triterpene from *Tithonia diversifolia* Potentiate Insulin Release: Invitro and in silico Approach. Orlando: Endocrine Society.
- Chagas-Paula, D. A., Oliveira, R. B., Rocha, B. A. and Da Costa, F. B. (2012). Ethnobotany, chemistry, and biological activities of the genus *Tithonia* (Asteraceae). *Chemistry and Biodiversity*.
- Chagas-Paula, D. A., Zhang, T., Da Costa, F. B. and Edrada-Ebel, R. (2015). A Metabolomic Approach to Target Compounds from the Asteraceae Family for Dual COX and LOX Inhibition. *Metabolites*, 5 : 404–430.
- Chunudom, L., Thongsom, M., Karim, N., Rahman, M. A., Rana, M. N. and Tangpong, J. (2020). *Tithonia diversifolia* aqueous fraction plays a protective role against alloxan-induced diabetic mice via modulating GLUT2 expression. *South African Journal of Botany*, 133 : 118–123.
- Darmawi, A. R., Saleh, C. and Kartika, R. (2015). Aktivitas antihiperlikemik dari ekstrak etanol dan n-heksana daun kembang bulan [*Tithonia diversifolia* A.Gray] pada tikus jantan. *Jurnal Kimia Mulawarman*, 12 (2) : 59–63.
- DeFronzo, R. A. (1999). Pharmacologic Therapy for Type 2 Diabetes Mellitus. *Annals of Internal Medicine*, 131 (4) : 281.
- DeFronzo, R. A., Ferrannini, E., Groop, L., Henry, R. R., Herman, W. H., Holst, J.

- J., *et al.* (2015). Type 2 diabetes mellitus. *Nature Reviews Disease Primers*, 1 (1) : 15019.
- DeFronzo, R. A. and Tripathy, D. (2009). Skeletal muscle insulin resistance is the primary defect in type 2 diabetes. *Diabetes care*, 32 (Suppl 2) : S157.
- DepKesRI. (2008). *Farmakope Herbal Indonesia Edisi 1*. Departemen Kesehatan Republik Indonesia.
- Van Dijk, T. H., Laskewitz, A. J., Grefhorst, A., Boer, T. S., Bloks, V. W., Kuipers, F., *et al.* (2013). A novel approach to monitor glucose metabolism using stable isotopically labelled glucose in longitudinal studies in mice. *Laboratory Animals*, 47 : 79–88.
- Dlamini, B. S., Chen, C.-R., Shyu, D. J. H. and Chang, C.-I. (2020). Flavonoids from *Tithonia diversifolia* and their Antioxidant and Antibacterial Activity. *Chemistry of Natural Compounds*, 56 (5) : 906–908.
- Ennequin, G., Capel, F., Caillaud, K., Chavanelle, V., Etienne, M., Teixeira, A., *et al.* (2017). Neuregulin 1 improves complex 2-mediated mitochondrial respiration in skeletal muscle of healthy and diabetic mice. *Scientific Reports*, 7 (1).
- Fasuyi, A. O., Dairo, F. A. S. and Ibitayo, F. J. (2010). Ensiling wild sunflower (*Tithonia diversifolia*) leaves with sugar cane molasses. *Livestock Research for Rural Development*, 22 (3).
- Fauziyah, Y., Sunarti, Hanoum, I. F. and Wahyuningsih, M. S. H. (2018). Ethanol extract of *Tithonia diversifolia* (Hemsley) A Gray standardized ameliorates hyperglycemia, polyphagia, and weight loss in diabetic rats. *Molekul*, 13 (1) : 72–79.
- Fryer, L. G. D., Parbu-Patel, A. and Carling, D. (2002). The anti-diabetic drugs rosiglitazone and metformin stimulate AMP-activated protein kinase through distinct signaling pathways. *Journal of Biological Chemistry*, 277 (28).
- Garnier, A., Fortin, D., Deloménie, C., Momken, I., Veksler, V. and Ventura-Clapier, R. (2003). Depressed mitochondrial transcription factors and oxidative capacity in rat failing cardiac and skeletal muscles. *Journal of*

Physiology, 551 (2) : 491–501.

Gayoso-Diz, P., Otero-González, A., Rodriguez-Alvarez, M. X., Gude, F., García, F., De Francisco, A., *et al.* (2013). Insulin resistance (HOMA-IR) cut-off values and the metabolic syndrome in a general adult population: effect of gender and age: EPIRCE cross-sectional study. *BMC Endocrine Disorders*, 13 (1) : 47.

Giacomo, C. Di, Vanella, L., Sorrenti, V., Santangelo, R., Barbagallo, I., Calabrese, G., *et al.* (2015). Effects of *Tithonia diversifolia* (Hemsl.) A. Gray extract on adipocyte differentiation of human mesenchymal stem cells. *PLoS ONE*, 10 (4) : 1–15.

Gilkerson, R., Bravo, L., Garcia, I., Gaytan, N., Herrera, A., Maldonado, A., *et al.* (2013). The mitochondrial nucleoid: Integrating mitochondrial DNA into cellular homeostasis. *Cold Spring Harbor Perspectives in Biology*, 5 : a011080.

Haupt, E., Knick, B., Koschinsky, T., Liebermeister, H., Schneider, J. and Hirche, H. (1991). Oral antidiabetic combination therapy with sulphonylureas and metformin. in *Diabete et Metabolisme*.

Heinrich, M., Robles, M., West, J. E., Ortiz De Montellano, B. R. and Rodriguez, E. (1998). Ethnopharmacology of Mexican asteraceae (compositae). *Annual Review of Pharmacology and Toxicology*.

Himsworth, H. P. (2013). Diabetes mellitus: its differentiation into insulin-sensitive and insulin-insensitive types. 1936. *International journal of epidemiology*, 42 (6).

Hu, F. B. (2011). Globalization of diabetes: The role of diet, lifestyle, and genes. in *Diabetes Care*.

Jackman, M. R. and Willis, W. T. (1996). Characteristics of mitochondria isolated from type I and type IIb skeletal muscle. *American Journal of Physiology-Cell Physiology*, 270 (2) : C673–C678.

John-Dewole, O. O. and Oni, S. O. (2013). Phytochemical and Antimicrobial Studies of Extracts from the Leaves of *Tithonia Diversifolia* for Pharmaceutical Importance. *IOSR Journal of Pharmacy and Biological*

Sciences, 6 (4) : 21–25.

- Kaikini, A. A., Kanchan, D. M., Nerurkar, U. N. and Sathaye, S. (2017). Targeting mitochondrial dysfunction for the treatment of diabetic complications: Pharmacological interventions through natural products. *Pharmacognosy Reviews*, 11 (22) : 128–135.
- Kelley, D. E., He, J., Menshikova, E. V and Ritov, V. B. (2002). Dysfunction of Mitochondria in Human Skeletal Muscle in Type 2 Diabetes. *Diabetes*, 51 : 2944–2950.
- Kooti, W., Farokhipour, M., Asadzadeh, Z., Ashtary-Larky, D. and Asadi-Samani, M. (2016). The role of medicinal plants in the treatment of diabetes: a systematic review. *Electronic Physician*, 8 (1) : 1832–1842.
- Kukidome, D., Nishikawa, T., Sonoda, K., Imoto, K., Fujisawa, K., Yano, M., *et al.* (2006). Activation of AMP-activated protein kinase reduces hyperglycemia-induced mitochondrial reactive oxygen species production and promotes mitochondrial biogenesis in human umbilical vein endothelial cells. *Diabetes*, 55 (1).
- Kuroda, M., Yokosuka, A., Kobayashi, R., Jitsuno, M., Kando, H., Nosaka, K., *et al.* (2007). Sesquiterpenoids and flavonoids from the aerial parts of *Tithonia diversifolia* and their cytotoxic activity. *Chemical and Pharmaceutical Bulletin*, 55 (8).
- Kuznetsov, A. V, Mayboroda, O., Kunz, D., Winkler, K., Schubert, W. and Kunz, W. S. (1998). Functional Imaging of Mitochondria in Saponin-permeabilized Mice Muscle Fibers. *The Journal of Cell Biology*, 140 (5) : 1091–1099.
- Ladeska, V., Dewanti, E. and Sari, D. I. (2019). Pharmacognostical Studies and Determination of Total Flavonoids of Paitan (*Tithonia diversifolia* (Hemsl.) A. Gray. *Pharmacognosy Journal*, 11 (6) : 1256–1261.
- Leabman, M. K., Huang, C. C., DeYoung, J., Carlson, E. J., Taylor, T. R., De la Cruz, M., *et al.* (2003). Natural variation in human membrane transporter genes reveals evolutionary and functional constraints. *Proceedings of the National Academy of Sciences of the United States of America*, 100 (10).

- Lee, C. H., Shih, A. Z. L., Woo, Y. C., Fong, C. H. Y., Leung, O. Y., Janus, E., *et al.* (2016). Optimal Cut-Offs of Homeostasis Model Assessment of Insulin Resistance (HOMA-IR) to Identify Dysglycemia and Type 2 Diabetes Mellitus: A 15-Year Prospective Study in Chinese. *PLOS ONE*. Edited by C. Hu, 11 (9) : e0163424.
- Lin, H. R. (2012). Sesquiterpene lactones from *Tithonia diversifolia* act as peroxisome proliferator-activated receptor agonists. *Bioorganic and Medicinal Chemistry Letters*, 22 (8) : 2954–2958.
- Masiello, P., Broca, C., Gross, R., Roye, M., Manteghetti, M., Hillaire-Buys, D., *et al.* (1998). Experimental NIDDM: Development of a new model in adult rats administered streptozotocin and nicotinamide. *Diabetes*, 47 (2) : 224–229.
- Mather, K. (2009). Surrogate measures of insulin resistance: of rats, mice, and men. *Am J Physiol Endocrinol Metab*, 296 : 398–399.
- Matthews, D. R., Hosker, J. P., Rudenski, A. S., Naylor, B. A., Treacher, D. F. and Turner, R. C. (1985). Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*, 28 (7).
- Medeiros, D. M. (2008). Assessing mitochondria biogenesis. *Methods*, 46 (4) : 288–294.
- MIMS. (2021). *Metformin: Indication, Dosage, Side Effect, Precaution | MIMS Indonesia*.
- Miranda, M. A. F. M., Varela, R. M., Torres, A., Molinillo, M. G., Gualtieri, S. C. J. and Macías, F. A. (2015). Phytotoxins from *Tithonia diversifolia*. *J. Nat. Prod*, 78 : 1083–1092.
- Miura, T., Nosaka, K., Ishii, H. and Ishida, T. (2005). Antidiabetic Effect of Nitobegiku, the Herb *Tithonia diversifolia*, in KK-Ay Diabetic Mice. *Notes Biol. Pharm. Bull*, 28 (11) : 2152–2154.
- Mogensen, M., Sahlin, K., Fernström, M., Glinborg, D., Vind, B. F., Beck-Nielsen, H., *et al.* (2007). Mitochondrial respiration is decreased in skeletal muscle of patients with type 2 diabetes. *Diabetes*, 56 (6).

- Mogensen, M. and Sahlin, K. (2005). Mitochondrial efficiency in rat skeletal muscle: influence of respiration rate, substrate and muscle type. *Acta Physiologica Scandinavica*, 185 (3) : 229–236.
- Montgomery, M. K. and Turner, N. (2015). Mitochondrial dysfunction and insulin resistance: an update. *Endocrine Connections*, 4 : R1–R15.
- Muin, R. (2019). Uji Efektivitas Antidiabetes Ekstrak Etanol Daun Kembang Bulan (*Tithonia diversifolia*). *Journal of Pharmaceutical Science and Herbal Technology*, 4 (1) : 1–3.
- Musi, N., Hirshman, M. F., Nygren, J., Svanfeldt, M., Bavenholm, P., Rooyackers, O., *et al.* (2002). Metformin increases AMP-activated protein-kinase activity in skeletal muscle of subjects with type 2 diabetes. *Diabetes*, 51 (7).
- Olayinka, B. U., Raiyemo, D. A. and Etejere, E. O. (2015). Phytochemical and Proximate Composition of *Tithonia diversifolia* (Hemsl.) A. Gray. *Annals. Food Science and Technology*, 16 (1) : 195–200.
- Olukunle, J. O., Okediran, B., Sogebi, E. and Jacobs, E. (2014). Hypoglycaemic and Hypolipidaemic Effects of the Aqueous Leaf Extracts of *Tithonia diversifolia*. *Annual Research & Review in Biology*, 4 (16) : 2655–2662.
- Omolola, T. O. (2019). Phytochemical, Proximate and Elemental Composition of *Tithonia diversifolia* (Hemsl.) A. Gray leaves. *International Annals of Science*, 8 (1).
- Otusanya, O. and Ilori, O. (2012). Phytochemical Screening and the Phytotoxic Effects of Aqueous Extracts of *Tithonia diversifolia* (Hemsl.) A. Gray. *International Journal of Biology*, 4 (3).
- Owoyele, V. B., Wuraola, C. O., Soladoye, A. O. and Olaleye, S. B. (2004). Studies on the anti-inflammatory and analgesic properties of *Tithonia diversifolia* leaf extract. *Journal of Ethnopharmacology*, 90 (2–3) : 317–321.
- Owoyele, V. B., Wuraola, C. O. and Soladoye, A. O. (2004). Studies on the anti-inflammatory and analgesic properties of *Tithonia diversifolia* leaf extract. *Journal of Ethnopharmacology*, 90 : 317–321.
- Patti, M. E. and Corvera, S. (2010). The role of mitochondria in the pathogenesis of type 2 diabetes. *Endocrine Reviews*.

- PERKENI. (2019). *Pedoman Pengelolaan dan Pencegahan Diabetes Mellitus Tipe 2 Dewasa di Indonesia*. Pengurus Besar Perkumpulan Endokrinologi Indonesia (PB PERKENI).
- Petersen, K. F., Befroy, D., Dufour, S., Dziura, J., Ariyan, C., Rothman, D. L., *et al.* (2003). Mitochondrial dysfunction in the elderly: Possible role in insulin resistance. *Science*, 300 (5622).
- Petersen, K. F., Dufour, S., Befroy, D., Garcia, R. and Shulman, G. I. (2004). Impaired Mitochondrial Activity in the Insulin-Resistant Offspring of Patients with Type 2 Diabetes. *New England Journal of Medicine*, 350 (7).
- Picard, M., Csukly, K., Robillard, M.-E., Godin, R., Ascah, A., Bourcier-Lucas, C., *et al.* (2008). Resistance to Ca²⁺-induced opening of the permeability transition pore differs in mitochondria from glycolytic and oxidative muscles. *Am J Physiol Regul Integr Comp Physiol*, 295 : 659–668.
- Picard, M., Hepple, R. T. and Burrelle, Y. (2012). Mitochondrial functional specialization in glycolytic and oxidative muscle fibers: tailoring the organelle for optimal function. *Am J Physiol Cell Physiol*, 302 : 629–641.
- Prasetyo, A., Denashurya, T. G., Putri, W. S. and Ilmiawan, M. I. (2016). Perbandingan Efek Hipoglikemik Infusa Daun Kembang Bulan (*Tithonia diversifolia* (Hemsley) A. Gray) dan Metformin pada Tikus yang Diinduksi Aloksan. *Cermin Dunia Kedokteran*, 43 (2) : 91–94.
- Qin, Z. Y., Zhang, M., Dai, Y. M., Wang, Y. M., Zhu, G. Z., Zhao, Y. P., *et al.* (2014). Metformin prevents LYRM1-induced insulin resistance in 3T3-L1 adipocytes via a mitochondrial-dependent mechanism. *Experimental Biology and Medicine*, 239 (12).
- Ranti, I., Wahyuningsih, M. S. H. and Wirohadidjojo, Y. W. (2018). The antifibrotic effect of isolate tagitinin C from *Tithonia diversifolia* (Hemsley) A. gray on keloid fibroblast cell. *Pan African Medical Journal*, 30.
- Di Re, M., Sembongi, H., He, J., Reyes, A., Yasukawa, T., Martinsson, P., *et al.* (2009). The accessory subunit of mitochondrial DNA polymerase γ determines the DNA content of mitochondrial nucleoids in human cultured cells. *Nucleic Acids Research*, 37 (17) : 5701–5713.

- Robinson, M. M. and Zhang, X. (2011). The World Medicines Situation 2011 Traditional Medicines : Global Situation , Issues and Challenges. *World Health Organization*.
- Rüngeler, P., Lyß, G., Castro, V., Mora, G., Pahl, H. and Merfort, I. (1998). Study of Three Sesquiterpene Lactones from *Tithonia diversifolia* on their Anti-Inflammatory Activity Using the Transcription Factor NF- κ B and Enzymes of the Arachidonic Acid Pathway as Targets. *Planta Medica*, 64 (07) : 588–593.
- Saeedi, Pouya, Petersohn, I., Salpea, P., Malanda, B., Karuranga, S., Unwin, N., *et al.* (2019). Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9th edition. *Diabetes Research and Clinical Practice*, 157 : 107843.
- Salpeter, S. R., Greyber, E., Pasternak, G. A. and Salpeter, E. E. (2003). Risk of Fatal and Nonfatal Lactic Acidosis with Metformin Use in Type 2 Diabetes Mellitus: Systematic Review and Meta-analysis. *Archives of Internal Medicine*.
- Samuel, V. T. and Shulman, G. I. (2012). Mechanisms for insulin resistance: Common threads and missing links. *Cell*.
- Sánchez-Duarte, S., Márquez-Gamiño, S., Montoya-Pérez, R., Villicaña-Gómez, E. A., Vera-Delgado, K. S., Caudillo-Cisneros, C., *et al.* (2021). Nicorandil decreases oxidative stress in slow- and fast-twitch muscle fibers of diabetic rats by improving the glutathione system functioning. *Journal of Diabetes Investigation*, 12 (7) : 1152–1161.
- Sasmita, F. W., Susetyarini, E., Husamah, H. and Pantiwati, Y. (2017). Efek Ekstrak Daun Kembang Bulan (*Tithonia diversifolia*) terhadap Kadar Glukosa Darah Tikus Wistar (*Rattus norvegicus*) yang Diinduksi Alloxan. *Biosfera*, 34 (1) : 22.
- Siedle, B., García-Piñ, A. J., Murillo, R., Rgen Schulte-Mönting, J., Castro, V., Rü, P., *et al.* (2004). Quantitative Structure-Activity Relationship of Sesquiterpene Lactones as Inhibitors of the Transcription Factor NF-KB. *J.*

Med. Chem., 47 : 6042–6054.

- Silva, A. M. R., Ferreira, N. L. O., Oliveira, A. E., Leonardo, L. B. and Conceição, E. C. (2017). Comparison of ultrasound-Assisted extraction and dynamic maceration over content of tagitinin C obtained from *Tithonia diversifolia* (Hemsl.) A. gray leaves using factorial design. *Pharmacognosy Magazine*, 13 (50) : 270–274.
- Skuratovskaia, D., Komar, A., Vulf, M. and Litvinova, L. (2020). Mitochondrial destiny in type 2 diabetes: The effects of oxidative stress on the dynamics and biogenesis of mitochondria. *PeerJ*, 8 : e9741.
- Suwa, M., Egashira, T., Nakano, H., Sasaki, H. and Kumagai, S. (2006). Metformin increases the PGC-1 α protein and oxidative enzyme activities possibly via AMPK phosphorylation in skeletal muscle in vivo. *J Appl Physiol*, 101 : 1685–1692.
- Teixeira, M. A., Chaguri, L. do C. A. G., Carissimi, A. S., Souza, N. L. de, Mori, C. M. C., Gomes, V. M. W., *et al.* (2000). Hematological and biochemical profiles of rats (*Rattus norvegicus*) kept under microenvironmental ventilation system. *Brazilian Journal of Veterinary Research and Animal Science*, 37 (5) : 341–347.
- Thomford, N. E., Senthebane, D. A., Rowe, A., Munro, D., Seele, P., Maroyi, A., *et al.* (2018). Natural Products for Drug Discovery in the 21st Century: Innovations for Novel Drug Discovery. *International Journal of Molecular Sciences*, 19 (6) : 1578.
- Thongsom, M., Chunglok, W., Kuanchuea, R. and Tangpong, J. (2013). Antioxidant and hypoglycemic effects of *Tithonia diversifolia* aqueous leaves extract in alloxan-induced diabetic mice. *Advances in Environmental Biology*, 7 (9) : 2116–2125.
- Timóteo, A. T., Miranda, F., Carmo, M. M. and Ferreira, R. C. (2014). Optimal Cut-Off Value for Homeostasis Model Assessment (HOMA) Index of Insulin-Resistance in a Population of Patients Admitted Electively in a Portuguese Cardiology Ward. *Acta Médica Portuguesa*, 27 (4) : 473–479.
- Wahyuningsih, M. S. H., Wijayanti, M. A., Budiyanto, A. and Hanafi, M. (2015).

- Isolation and Identification of Potential Cytotoxic Compound from Kembang Bulan [*Tithonia diversifolia* (HEMSLEY) A GRAY] Leaves. *International Jurnal of Pharmacy and Pharmaceutical Sciences*, 7 (6) : 298–301.
- Wallace, T. M., Levy, J. C. and Matthews, D. R. (2004). Use and abuse of HOMA modeling. *Diabetes Care*, 27 (6) : 1487–1495.
- Yashpal, S., MK, G., Nikhil, T. and Kumar, M. R. (2013). A Study of Insulin Resistance by HOMA-IR and its Cut-off Value to Identify Metabolic Syndrome in Urban Indian Adolescents. *Journal of Clinical Research in Pediatric Endocrinology*, 5 (4) : 245–251.
- Yazdi, H. B., Hojati, V., Shiravi, A., Hosseinian, S., Vaezi, G. and Hadjzadeh, M.-A.-R. (2019). Liver Dysfunction and Oxidative Stress in Streptozotocin-Induced Diabetic Rats: Protective Role of *Artemisia turanica*. *Journal of Pharmacopuncture*, 22 (2) : 109–114.
- Yin, O. Q. P., Tomlinson, B. and Chow, M. S. S. (2006). Variability in renal clearance of substrates for renal transporters in Chinese subjects. *Journal of Clinical Pharmacology*, 46 (2).
- Ylikallio, E., Tyynismaa, H., Tsutsui, H., Ide, T. and Suomalainen, A. (2010). High mitochondrial DNA copy number has detrimental effects in mice. *Human Molecular Genetics*, 19 (13) : 2695–2705.
- Zhao, G., Li, X., Chen, W., Xi, Z. and Sun, L. (2012). Three new sesquiterpenes from *Tithonia diversifolia* and their anti-hyperglycemic activity. *Fitoterapia*, 83 (8) : 1590–1597.
- Zhou, G., Myers, R., Li, Y., Chen, Y., Shen, X., Fenyk-Melody, J., *et al.* (2001). Role of AMP-activated protein kinase in mechanism of metformin action. *Journal of Clinical Investigation*, 108 (8).
- Zuo, C. S., Sung, Y.-H., Simonson, D. C., Habecker, E. and Wang, J. (2012). Reduced T2* Values in Soleus Muscle of Patients with Type 2 Diabetes Mellitus. *PLoS ONE*, 7 (11) : 49337.