



## DAFTAR PUSTAKA

- Akash, B. A. (2013) ‘Thermochemical Liquefaction of Coal’, *Int. J. of Thermal & Environmental Engineering*, 5(1), pp. 51–60. doi: 10.5383/ijtee.05.01.006.
- Berkowitz, N., Calderon, J. and Liron, A. (1988) ‘Some observations respecting reaction paths in coal liquefaction3. Time and temperature effects in coal solubilization’, *Fuel*, 67(8), pp. 1139–1142. doi: 10.1016/0016-2361(88)90384-5.
- Bodzek, D. and Marzec, A. (1981) ‘Molecular components of coal and coal structure’, *Fuel*, 60(1), pp. 47–51. doi: 10.1016/0016-2361(81)90030-2.
- BPPT (2019) *Indonesia Energy Outlook 2019: The Impact of Increased Utilization of New and Renewable Energy on the National Economy*. Edited by A. Sugiyono et al. Jakarta: PPIPE BPPT.
- BPPT (2020) *Indonesia Energy Outlook 2020 - Special Edition Dampak Pandemi COVID-19 terhadap Sektor Energi di Indonesia*, PPIPE BPPT. Edited by Adiarso, H. Edi, and S. Agus. Jakarta: PPIPE BPPT. Available at: [https://www.researchgate.net/publication/343903321\\_OUTLOOK\\_ENERGI\\_IN\\_DONESIA\\_2020\\_Dampak\\_Pandemi\\_COVID-19\\_terhadap\\_Sektor\\_Energi\\_di\\_Indonesia](https://www.researchgate.net/publication/343903321_OUTLOOK_ENERGI_IN_DONESIA_2020_Dampak_Pandemi_COVID-19_terhadap_Sektor_Energi_di_Indonesia).
- Chang, S. et al. (2016) ‘Clean Coal Technologies in China: Current Status and Future Perspectives’, *Engineering*. Elsevier LTD on behalf of Chinese Academy of Engineering and Higher Education Press Limited Company, 2(4), pp. 447–459. doi: 10.1016/J.ENG.2016.04.015.
- Channiwala, S. A. and Parikh, P. P. (2002) ‘A unified correlation for estimating HHV of solid, liquid and gaseous fuels’, *Fuel*, 81(8), pp. 1051–1063. doi: 10.1016/S0016-2361(01)00131-4.
- Chawla, B. and Davis, B. H. (1989) ‘Effect of temperature and solvent on coal extraction under mild conditions’, *Fuel Processing Technology*, 23(2), pp. 133–148. doi: 10.1016/0378-3820(89)90073-8.
- Chen, C., Gao, J. and Yan, Y. (1998) ‘Role of Noncovalent Bonding in Swelling of Coal’, *Energy & Fuels*, 12(6), pp. 1328–1334. doi: 10.1021/ef970065w.
- Chen, Y., Mastalerz, M. and Schimmelmann, A. (2012) ‘Characterization of chemical functional groups in macerals across different coal ranks via micro-FTIR spectroscopy’, *International Journal of Coal Geology*. Elsevier B.V., 104, pp. 22–33. doi: 10.1016/j.coal.2012.09.001.
- Choudhary, S. et al. (2016) ‘REVIEW ON DIFFERENT TECHNOLOGIES FOR COAL LIQUEFACTION’, *International Research Journal of Engineering and Technology*, 3(4), pp. 2704–2708.
- Cupera, J. (2009) *Mathematical models of dissolution*.
- Figueroa, D. C. (2012) *Modelling of Solvent Extraction of Coal*.



Giralt, J., Fabregat, A. and Giralt, F. (1988) 'Kinetics of the Solvolysis of a Catalan Lignite with Tetralin and Anthracene Oil', *Industrial and Engineering Chemistry Research*, 27(7), pp. 1110–1114. doi: 10.1021/ie00079a004.

Giri, C. C. and Sharma, D. K. (2000a) 'Kinetic studies and shrinking core model on solvolytic extraction of coal', *Fuel Processing Technology*, 68(2), pp. 97–109. doi: 10.1016/S0378-3820(00)00112-0.

Giri, C. C. and Sharma, D. K. (2000b) 'Mass-transfer studies of solvent extraction of coals in N-methyl-2-pyrrolidone', *Fuel*, 79(5), pp. 577–585. doi: 10.1016/S0016-2361(99)00152-0.

Hao, P. et al. (2018) 'Effect of solvent and atmosphere on product distribution, hydrogen consumption and coal structural change during preheating stage in direct coal liquefaction', *Fuel*. Elsevier, 211(September 2017), pp. 783–788. doi: 10.1016/j.fuel.2017.09.122.

He, W. et al. (2017) 'Behavior of radicals during solvent extraction of three low rank bituminous coals', *Fuel Processing Technology*. Elsevier B.V., 156, pp. 221–227. doi: 10.1016/j.fuproc.2016.10.029.

He, W. et al. (2021) 'Interactions between Coal and Solvent during the Solvent Extraction of Coal in View of Free Radicals', *ACS Omega*. doi: 10.1021/acsomega.1c04260.

Huang, H. et al. (1999) 'Thermogravimetric and Rock-Eval studies of coal properties and coal rank', *Energy and Fuels*, 13(2), pp. 396–400. doi: 10.1021/ef980088q.

Iino, M. et al. (1988) 'Extraction of coals with CS<sub>2</sub>-N-methyl-2-pyrrolidinone mixed solvent at room temperature. Effect of coal rank and synergism of the mixed solvent', *Fuel*, 67(12), pp. 1639–1647. doi: 10.1016/0016-2361(88)90208-6.

Ikeda, K. et al. (2000) 'Kinetic evaluation of progress in coal liquefaction in the 1 t/d PSU for the NEDOL process', *Fuel*, 79(3–4), pp. 373–378. doi: 10.1016/S0016-2361(99)00172-6.

Jiang, H. et al. (2015) 'Isothermal Stage Kinetics of Direct Coal Liquefaction for Shenhua Shendong Bituminous Coal', *Energy and Fuels*, 29(11), pp. 7526–7531. doi: 10.1021/acs.energyfuels.5b01484.

Jin, L. et al. (2014) 'Direct liquefaction behaviors of Bulianta coal and its macerals', *Fuel Processing Technology*. Elsevier B.V., 128, pp. 232–237. doi: 10.1016/j.fuproc.2014.07.033.

Kabe, T. et al. (2004) 'Chemical and Macromolecular Structure of Coal', in *Coal and Coal-related Compounds:structure, reactivity and catalytic reactions*. 1st editio. Tokyo: Kodansha, pp. 81–126. doi: 10.1016/S0167-2991(04)80007-7.

Kashimura, N., Takanohashi, T. and Saito, I. (2006) 'Upgrading the Solvent Used for the Thermal Extraction of Sub-Bituminous Coal', *Energy & Fuels*, 20(5), pp. 2063–2066. doi: 10.1021/ef0601014.

Kim, H. et al. (2018) 'Correlation between solubility parameters and recovery of phenolic compounds from fast pyrolysis bio-oil by diesel extraction', *Carbon*



Resources      Conversion.      Elsevier,      1(3),      pp.      238–244.      doi:  
10.1016/j.crcon.2018.08.004.

Kim, K. *et al.* (2016) ‘Coal and solvent properties and their correlation with extraction yield under mild conditions’, *Korean Journal of Chemical Engineering*, 33(7), pp. 2142–2159. doi: 10.1007/s11814-016-0062-1.

Kim, S. Do *et al.* (2008) ‘Production of low ash coal by thermal extraction with N-methyl-2-pyrrolidinone’, *Korean Journal of Chemical Engineering*, 25(4), pp. 758–763. doi: 10.1007/s11814-008-0124-0.

Koyano, K., Takanohashi, T. and Saito, I. (2011) ‘Estimation of the extraction yield of coals by a simple analysis’, *Energy and Fuels*, 25(6), pp. 2565–2571. doi: 10.1021/ef200151u.

Li, C., Takanohashi, T. and Yoshida, T. (2004) ‘Effect of acid treatment on thermal extraction yield in ashless coal production’, *Fuel*, 83, pp. 727–732. doi: 10.1016/j.fuel.2003.06.002.

Li, X. *et al.* (2008) ‘Role of iron-based catalyst and hydrogen transfer in direct coal liquefaction’, *Energy and Fuels*, 22(2), pp. 1126–1129. doi: 10.1021/ef7006062.

Masaki, K. *et al.* (2004) ‘The Effects of Pretreatment and the Addition of Polar Compounds on the Production of “HyperCoal” from Subbituminous Coals’, *Energy & Fuels*, 18(4), pp. 995–1000. doi: 10.1021/ef049970o.

Mathews, J. P., Burgess-Clifford, C. and Painter, P. (2015) ‘Interactions of Illinois No. 6 bituminous coal with solvents: A review of solvent swelling and extraction literature’, *Energy and Fuels*, 29(3), pp. 1279–1294. doi: 10.1021/ef502548x.

Mayo, F. R. *et al.* (1988) ‘Extractions and reactions of coals below 100 °C. 3. Comparison of extraction products of Illinois No. 6 coal’, *Fuel*, 67(5), pp. 607–611. doi: 10.1016/0016-2361(88)90287-6.

Meshram, P. *et al.* (2015) ‘Demineralization of low grade coal - A review’, *Renewable and Sustainable Energy Reviews*. Elsevier, 41, pp. 745–761. doi: 10.1016/j.rser.2014.08.072.

Miura, K. *et al.* (2002) ‘Estimation of hydrogen bond distributions formed between coal and polar solvents using in situ IR technique’, *Energy and Fuels*, 16(1), pp. 23–31. doi: 10.1021/ef0101457.

Muangthong-On, T. *et al.* (2017) ‘Effect of Solvent on the Degradative Solvent Extraction of Low Rank Coal’, *Energy and Fuels*, 31(11), pp. 11954–11962. doi: 10.1021/acs.energyfuels.7b02352.

Norinaga, K., Kuniya, M. and Iino, M. (2002) ‘Effect of associative interaction on the dynamic viscoelastic property of coal concentrated solution’, *Energy and Fuels*, 16(1), pp. 62–68. doi: 10.1021/ef010154f.

Önal, Y. and Akol, S. (2003) ‘Influence of pretreatment on solvent-swelling and extraction of some Turkish lignites’, *Fuel*, 82(11), pp. 1297–1304. doi: 10.1016/S0016-2361(03)00033-4.



Pande, S. and Sharma, D. K. (2001) ‘Studies of Kinetics of Diffusion of N-Methyl-2-pyrrolidone (NMP), Ethylenediamine (EDA), and NMP+EDA (1:1, vol/vol) Mixed Solvent System in Chinakuri Coal by Solvent Swelling Techniques Shailaja’, *Energy & Fuels*, 15(5), pp. 1063–1068.

Pande, S. and Sharma, D. K. (2002) ‘Ethylenediamine-assisted solvent extraction of coal in N-methyl-2-pyrrolidone: Synergistic effect of ethylenediamine on extraction of coal in N-methyl-2-pyrrolidone’, *Energy and Fuels*, 16(1), pp. 194–204. doi: 10.1021/ef0001742.

Rudling, J. (1986) ‘Simple Model Based on Solubility Parameters for Liquid Desorption of Organic Solvents Adsorbed on Activated Carbon’, *Journal of Chromatography*, 362, pp. 175–185. doi: 10.1016/s0021-9673(01)86966-1.

Sakimoto, N., Koyano, K. and Takanohashi, T. (2013) ‘Relationship between Extraction Yield of Coal by Polar Solvent and Oxygen Functionalities in Coals’, *Energy & Fuels*, 27(11), pp. 6594–6597. doi: 10.1021/ef4016262.

Sanada, Y. and Honda, H. (1962) ‘Solvent Extraction of Coal—Relation between the Yield of Extract and the Solubility Parameter of the Solvent’, *Bulletin of the Chemical Society of Japan*, 35(8), pp. 1358–1360. doi: 10.1246/bcsj.35.1358.

Schobert, H. H. (1989) ‘The geochemistry of coal: I. The classification and origin of coal’, *Journal of Chemical Education*, 66(3), pp. 242–244. doi: 10.1021/ed066p242.

Sehume, T. Z. (2019) *Solvent extraction of a bituminous coal using a sweet sorghum bagasse derived solvent*. The North-West University.

Sharma, D. K. and Dhawan, H. (2018) ‘Separative Refining of Coals through Solvolytic Extraction under Milder Conditions: A Review’, *Industrial and Engineering Chemistry Research*, 57(25), pp. 8361–8380. doi: 10.1021/acs.iecr.8b00345.

Shi, L. et al. (2013) ‘Pyrolysis behavior and bonding information of coal — A TGA study’, *Fuel Processing Technology*. Elsevier B.V., 108, pp. 125–132. doi: 10.1016/j.fuproc.2012.06.023.

Shi, Z. et al. (2017) ‘Effect of hydrothermal treatment on structure and liquefaction behavior of Baiyinhua coal’, *Fuel Processing Technology*. Elsevier, 167(August), pp. 648–654. doi: 10.1016/j.fuproc.2017.08.015.

Shui, H. et al. (2013) ‘Thermal dissolution of Shenfu coal in different solvents’, *Fuel*. Elsevier Ltd, 108, pp. 385–390. doi: 10.1016/j.fuel.2012.11.005.

Shui, H., Wang, Z. and Wang, G. (2006) ‘Effect of hydrothermal treatment on the extraction of coal in the CS2/NMP mixed solvent’, *Fuel*, 85(12–13), pp. 1798–1802. doi: 10.1016/j.fuel.2006.02.005.

Siffert, B. et al. (1994) ‘Experimental determination of the electron donor and acceptor numbers of oxides by zetametry in organic media’, *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 92(1–2), pp. 107–111. doi: 10.1016/0927-7757(94)02788-9.

Solum, M. S., Pugmire, R. J. and Grant, D. M. (1989) ‘Carbon-13 solid-state NMR of



Argonne-premium coals', *Energy & Fuels*, 3(2), pp. 187–193. doi: 10.1021/ef00014a012.

Sönmez, Ö. et al. (2018) 'Influence of the addition of various ionic liquids on coal extraction with NMP', *Fuel*, 212(February 2017), pp. 12–18. doi: 10.1016/j.fuel.2017.10.017.

Sun, X. (2005) 'The investigation of chemical structure of coal macerals via transmitted-light FT-IR microspectroscopy', *Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy*, 62(1–3), pp. 557–564. doi: 10.1016/j.saa.2005.01.020.

Szeliga, J. and Marzec, A. (1983) 'Swelling of coal in relation to solvent electron-donor numbers', *Fuel*, 62(10), pp. 1229–1231. doi: 10.1016/0016-2361(83)90070-4.

Vasireddy, S. et al. (2011) 'Clean liquid fuels from direct coal liquefaction: chemistry, catalysis, technological status and challenges', *Energy Environ. Sci.*, 4(2), pp. 311–345. doi: 10.1039/C0EE00097C.

Wang, Z. et al. (2008) 'Study on the hydrothermal treatment of Shenhua coal', *Fuel*, 87(4–5), pp. 527–533. doi: 10.1016/j.fuel.2007.03.017.

Wender, I. (1976) 'Catalytic Synthesis of Chemicals from Coal', *Catalysis Reviews*, 14(1), pp. 97–129. doi: 10.1080/03602457608073408.

White, C. M. et al. (1997) 'Practical Notes on the Use of N -Methyl-2-pyrrolidinone as a Solvent for Extraction of Coal and Coal-Related Materials', 5(12), pp. 1105–1106.

Yan, J. et al. (2014) 'Effects of organic solvent treatment on the chemical structure and pyrolysis reactivity of brown coal', *Fuel*. Elsevier Ltd, 128, pp. 39–45. doi: 10.1016/j.fuel.2014.03.001.

Zhang, H.-X. (2016) 'Determination of Aromatic Structures of Bituminous Coal Using Sequential Oxidation', *Industrial & Engineering Chemistry Research*, 55(10), pp. 2798–2805. doi: 10.1021/acs.iecr.5b04899.

Zhang, L. et al. (2017) 'Mechanistic influences of different solvents on microwave-assisted extraction of Shenfu low-rank coal', *Fuel Processing Technology*. Elsevier B.V., 166, pp. 276–281. doi: 10.1016/j.fuproc.2017.06.016.