

DAFTAR PUSTAKA

- Armstrong, V. J., Muzylak, M., Sunter, A., Zaman, G., Saxon, L. K., Price, J. S. dan Lanyon, L. E. 2007. 'Wnt/ β -catenin signaling is a component of osteoblastic bone cell early responses to load-bearing and requires estrogen receptor α ', *Journal of Biological Chemistry*, 282(28), pp. 20715–20727. doi: 10.1074/jbc.M703224200.
- Burton, M.S., 2013. Diagnosis and Treatment of Adolescent Idiopathic Scoliosis. *Pediatr. Ann.* 42, e233–e237. doi:10.3928/00904481-20131022-09
- Burwell, R.G., Clark, E.M., Dangerfield, P.H., Moulton, A., 2016. Adolescent idiopathic scoliosis (AIS): a multifactorial cascade concept for pathogenesis and embryonic origin. *Scoliosis Spinal Disord.* 11, 8. doi:10.1186/s13013-016-0063-1
- Burwell, R.G., Dangerfield, P.H., Moulton, A., Grivas, T.B., 2011. Adolescent idiopathic scoliosis (AIS), environment, exposome and epigenetics: a molecular perspective of postnatal normal spinal growth and the etiopathogenesis of AIS with consideration of a network approach and possible implications for medical therapy. *Scoliosis* 6, 26. doi:10.1186/1748-7161-6-26
- Cao, Y., Min, J., Zhang, Q., Li, H., Li, H., 2016. Associations of LBX1 gene and adolescent idiopathic scoliosis susceptibility: a meta-analysis based on 34,626 subjects. *BMC Musculoskelet. Disord.* 17, 309. doi:10.1186/s12891-016-1139-

- Cilli, K., Tezeren, G., Taş, T., Bulut, O., Oztürk, H., Oztemur, Z., Unsaldi, T., 2009. [School screening for scoliosis in Sivas, Turkey]. *Acta Orthop. Traumatol. Turc.* 43, 426–430. doi:10.3944/AOTT.2009.426
- Daruwalla, J.S., Balasubramaniam, P., Chay, S.O., Rajan, U., Lee, H.P., 1985. Idiopathic scoliosis. Prevalence and ethnic distribution in Singapore schoolchildren. *J. Bone Joint Surg. Br.* 67, 182–184.
- Esposito, T., Uccello, R., Caliendo, R., Di Martino, G. F., Gironi Carnevale, U. A., Cuomo, S., Ronca, D. dan Varriale, B. 2009. ‘Estrogen receptor polymorphism, estrogen content and idiopathic scoliosis in human: A possible genetic linkage’, *Journal of Steroid Biochemistry and Molecular Biology*, 116(1–2), pp. 56–60. doi: 10.1016/j.jsbmb.2009.04.010
- Fan, Y.-H., Song, Y.-Q., Chan, D., Takahashi, Y., Ikegawa, S., Matsumoto, M., Kou, I., Cheah, K.S.E., Sham, P., Cheung, K.M.C., Luk, K.D.K., 2012. SNP rs11190870 near LBX1 is associated with adolescent idiopathic scoliosis in southern Chinese. *J. Hum. Genet.* 57, 244–246. doi:10.1038/jhg.2012.11
- Grauers, A., Einarsdottir, E., Gerdhem, P., 2016. Genetics and pathogenesis of idiopathic scoliosis. *Scoliosis Spinal Disord.* 11, 45. doi:10.1186/s13013-016-0105-8
- Gross, M.K., Dottori, M., Goulding, M., 2012. Lbx1 specifies somatosensory association interneurons in the dorsal spinal cord. *Neuron* 34. doi:10.1016/S0896- 6273(02)00690-6
- Guo L, Yamashita H, Kou I, et al. Functional Investigation of a Non-coding Variant

Associated with Adolescent Idiopathic Scoliosis in Zebrafish: Elevated
Expression of the Ladybird Homeobox Gene Causes Body Axis Deformation.

PLoS Genet. 2016;12(1):1-28. doi:10.1371/journal.pgen.1005802

Herkowitz HN, Garfin SR, Eismont FJ, Bell GR. *Rothman - Simeone : The Spine 6th
Edition.* 6th ed. Philadelphia: Elseviers; 2011

Jagla K, Dollé P, Mattei MG, et al. Mouse Lbx1 and human LBX1 define a novel
mammalian homeobox gene family related to the Drosophila lady bird genes.
Mech Dev. 1995;53(3):345-356.
<http://www.ncbi.nlm.nih.gov/pubmed/8645601>. Accessed June 12, 2019.

Jiang H, Qiu X, Dai J, et al. Association of rs11190870 near LBX1 with adolescent
idiopathic scoliosis susceptibility in a Han Chinese population. *Eur Spine J.*
2013;22(2):282-286. doi:10.1007/s00586-012-2532-4

Kamtsiuris, P., Atzpodien, K., Ellert, U., Schlack, R., Schlaud, M., 2007. Prevalence
of somatic diseases in German children and adolescents. Results of the German
Health Interview and Examination Survey for Children and Adolescents
(KiGGS). *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*
50, 686–700. doi:10.1007

Kesling, K.L., Reinker, K.A., 1997. Scoliosis in twins. A meta-analysis of the literature
and report of six cases. *Spine Phila Pa* 1976 22. doi:10.1097/00007632-
199709010-00014

Konieczny, M.R., Senyurt, H., Krauspe, R., 2013. Epidemiology of adolescent
idiopathic scoliosis. *J. Child. Orthop.* 7, 3–9. doi:10.1007/s11832-012-0457-4

- Kudo D, Miyakoshi N, Hongo M, et al. LBX1 mRNA Expression in Paravertebral Muscles of Patients with Adolescent Idiopathic Scoliosis : A Preliminary Study. *Akita Univ.* 2013;(40):151-156.
- Kusumi, K., Dunwoodie, S.L., 2010. The genetics and development of scoliosis. Springer, New York.
- Leboeuf, D., Letellier, K., Alos, N., Edery, P. dan Moldovan, F. 2009. ‘Do estrogens impact adolescent idiopathic scoliosis?’, *Trends in Endocrinology and Metabolism*, 20(4), pp. 147–152. doi: 10.1016/j.tem.2008.12.004.
- Letellier, K., Azeddine, B., Parent, S., Labelle, H., Rompré, P. H., Moreau, A. dan Moldovan, F. 2008. ‘Estrogen cross-talk with the melatonin signaling pathway in human osteoblasts derived from adolescent idiopathic scoliosis patients’, *Journal of Pineal Research*, 45(4), pp. 383–393. doi: 10.1111/j.1600-079X.2008.00603.
- Liang, J., Xing, D., Li, Z., Chua, S., Li, S., 2014. Association Between rs11190870 Polymorphism Near LBX1 and Susceptibility to Adolescent Idiopathic Scoliosis in East Asian Population: A Genetic Meta-Analysis. *Spine*. doi:10.1097/BRS.0000000000000303
- Lonstein, J.E., Carlson, J.M., 1984. The prediction of curve progression in untreated idiopathic scoliosis during growth. *J Bone Jt. Surg Am* 66. doi:10.2106/00004623-198466070-00013
- Londono, D., Kou, I., Johnson, T.A., Sharma, S., Ogura, Y., Tsunoda, T., Takahashi, A., Matsumoto, M., Herring, J.A., Lam, T.-P., Wang, X., Tam, E.M.S., Song,

Y.-Q., Fan, Y.-H., Chan, D., Cheah, K.S.E., Qiu, X., Jiang, H., Huang, D., Japanese Scoliosis Clinical Research Group, TSRHC IS Clinical Group, null, International Consortium for Scoliosis Genetics, null, Su, P., Sham, P., Cheung, K.M.C., Luk, K.D.K., Gordon, D. Qiu, Y., Cheng, J., Tang, N., Ikegawa, S., Wise, C.A., 2014. A meta-analysis identifies adolescent idiopathic scoliosis association with LBX1 locus in multiple ethnic groups. *J. Med. Genet.* 51, 401–406. doi:10.1136/jmedgenet-2013-102067

Nery, L.S., Halpern, R., Nery, P.C., Nehme, K.P., Tetelbom Stein, A., 2010. Prevalence of scoliosis among school students in a town in southern Brazil. *Sao Paulo Med. J.* 128, 69–73. doi:10.1590/S1516-31802010000200005

Netter, Frank H. 1906-1991, *Atlas of Human Anatomy*. Philadelphia, PA: Elsevier, 2019.

Reamy, B. V dan Slakey, J. B. 2001. ‘Adolescent idiopathic scoliosis: review and current concepts.’, *American family physician*, 64(1), pp. 111–6. doi: 10.1016/S0140-6736(94)90572-X

Schlösser TPC, Van Der Heijden GJMG, Versteeg AL, Castelein RM. How “idiopathic” is adolescent idiopathic scoliosis? A systematic review on associated abnormalities. *PLoS One*. 2014;9(5):1-10. doi:10.1371/journal.pone.0097461

Sharma, S., Gao, X., Londono, D., Devroy, S.E., Mauldin, K.N., Frankel, J.T., Brandon, J.M., Zhang, D., Li, Q.-Z., Dobbs, M.B., Gurnett, C.A., Grant, S.F.A., Hakonarson, H., Dormans, J.P., Herring, J.A., Gordon, D., Wise, C.A., 2011.

Genome-wide association studies of adolescent idiopathic scoliosis suggest candidate susceptibility genes. *Hum. Mol. Genet.* 20, 1456–1466. doi:10.1093/hmg/ddq571

Skibinska, I., Tomaszewski, M., Andrusiewicz, M., Urbaniak, P., Czarnecka-Klos, R., Shadi, M., Kotwicki, T. dan Kotwicka, M. 2016. ‘Expression of estrogen receptor coactivator proline-, glutamic acid- and leucine-rich protein 1 within paraspinal muscles in adolescents with idiopathic scoliosis’, *PLoS ONE*, 11(4), pp. 1–13. doi: 10.1371/journal.pone.0152286.

Sieber, M.A., Storm, R., Martinez-de-la-Torre, M., Müller, T., Wende, H., Reuter, K., Vasyutina, E., Birchmeier, C., 2007. Lbx1 Acts as a Selector Gene in the Fate Determination of Somatosensory and Viscerosensory Relay Neurons in the Hindbrain. *J. Neurosci.* 27, 4902–4909. doi:10.1523/JNEUROSCI

Solomon, L., Warwick, D., Nayagam, S., Solomon, L., 2014. Apley and Solomon’s concise system of orthopaedics and trauma.

Soucacos, P.N., Soucacos, P.K., Zacharis, K.C., Beris, A.E., Xenakis, T.A., 1997. School-screening for scoliosis. A prospective epidemiological study in northwestern and central Greece. *J. Bone Joint Surg. Am.* 79, 1498–1503.

Takahashi Y, Kou I, Takahashi A, et al. A genome-wide association study identifies common variants near LBX1 associated with adolescent idiopathic scoliosis. *Nat Genet.* 2011;43(12):1237-1240. doi:10.1038/ng.974

Vaccarro AR. *Spine Core Knowledge in Orthopaedics*. Philadelphia: Mosby; 2005

Wang, W.J., Yeung, H.Y., Chu, W.C.-W., Tang, N.L.-S., Lee, K.M., Qiu, Y., Burwell,

R.G., Cheng, J.C.Y., 2011. Top theories for the etiopathogenesis of adolescent idiopathic scoliosis. *J. Pediatr. Orthop.* 31, S14-27.
doi:10.1097/BPO.0b013e3181f73c12

Weinstein SL, Zavala DC, Ponseti I V. Idiopathic scoliosis. Long-term follow-up and prognosis in untreated patients. *J Bone Jt Surg - Ser A.* 1981;63(5):702-712.
doi:10.2106/00004623-198163050-00003

Weinstein, S.L., Dolan, L.A., Cheng, J.C., Danielsson, A., Morcuende, J.A., 2008. Adolescent idiopathic scoliosis. *Lancet* 371.
doi:10.1016/S0140-6736(08)60658- 3

Zhu, Z., Xu, L., Qiu, Y., 2015b. Current progress in genetic research of adolescent idiopathic scoliosis. *Ann. Transl. Med.* 3, S19.
doi:10.3978/j.issn.2305- 5839.2015.02.04