

DAFTAR PUSTAKA

- Ak, O., Batirel, A., Ozer, S. and Çolakoğlu, S. (2011) 'Nosocomial infections and risk factors in the intensive care unit of a teaching and research hospital: A prospective cohort study', *Med. Sci. Monit.*, 17(5): 29–34. doi: 10.12659/MSM.881750.
- Al-Hasan, M. N., Lahr, B. D., Eckel-Passow, J. E. and Baddour, L. M. (2009) 'Antimicrobial resistance trends of Escherichia coli bloodstream isolates: A population-based study, 1998-2007', *J. Antimicrob. Chemother.*, 64(1): 169–174. doi: 10.1093/jac/dkp162.
- Al-Otaibi, F. E., Bukhari, E. E., Badr, M. and Alrabiaa, A. A. (2016) 'Prevalence and risk factors of Gram-negative bacilli causing blood stream infection in patients with malignancy', *Saudi Med. J.*, 37(9): 979–984. doi: 10.15537/smj.2016.9.14211.
- Albrecht, S. J., Fishman, N. O., Kitchen, J., Nachamkin, I., Bilker, W. B., Hoegg, C., Samel, C., Barbagallo, S., Arentzen, J. and Lautenbach, E. (2006) 'Reemergence of gram-negative health care-associated bloodstream infections', *Arch. Intern. Med.*, 166(12): 1289–1294. doi: 10.1001/archinte.166.12.1289.
- Alpern, E. R. (2008) 'In: Pediatric Emergency Medicine', *Elsevier Inc*, (Bacteremia): 530–534.
- Artini I, D. (2014) 'Ketidaktepatan Metode Microdilution Broth Otomatis Dalam Uji Sensitivitas 3 Jenis Antibiotik Pada Infeksi Acinetobacter Baumannii', *Program Studi Ilmu Kedokteran Klinik FKKMK Universitas Gadjah Mada*.
- Ataee, R. A., Mehrabi-Tavana, A., Hosseini, S. M. J., Moridi, K. and Zadeegan, M. G. (2012) 'A method for antibiotic susceptibility testing: Applicable and accurate', *Jundishapur J. Microbiol.*, 5(1): 341–345. doi: 10.5812/kowsar.20083645.2374.
- Bai, Y., Zheng, Z., Du, M., Yao, H., Liu, Y. and Suo, J. (2019) 'Bloodstream

- Infection and Its Clinical Characteristics and Relevant Factors Associated with Interventional Therapy in a Large Tertiary Hospital: A Six Years Surveillance Study', *BioMed Res. Inter.* 2019. doi: 10.1155/2019/8190475.
- Baldwin, C. D., Howe, G. B., Sampath, R., Blyn, L. B., Matthews, H., Harpin, V., Hall, T. A., *et al*(2009) 'Usefulness of multilocus polymerase chain reaction followed by electrospray ionization mass spectrometry to identify a diverse panel of bacterial isolates', *Diag. Microbiol. Infect. Dis.*. Elsevier B.V., 63(4): 403–408. doi: 10.1016/j.diagmicrobio.2008.12.012.
- Bauer, A., Kirby, W., Sherris, J. and Turck, M. (1966) 'Antibiotic susceptibility testing by a standardized single disk method', *Am J Clin Pathol*, 45: 493–496. doi: 10.1308/rcsann.2013.95.7.532.
- Bobenchik, A. M., Deak, E., Hindler, J. A., Charlton, C. L. and Humphries, R. M. (2017) 'Performance of vitek 2 for antimicrobial susceptibility testing of acinetobacter baumannii, pseudomonas aeruginosa, and stentrophomonas maltophilia with vitek 2 (2009 FDA) and CLSI M100S 26th edition breakpoints', *J. Clin. Microbiol.*, 55(2): 450–456. doi: 10.1128/JCM.01859-16.
- Buetti, N., Atkinson, A., Marschall, J. and Kronenberg, A. (2017) 'Incidence of bloodstream infections: A nationwide surveillance of acute care hospitals in Switzerland 2008-2014', *BMJ Open*, 7(3): 1–5. doi: 10.1136/bmjopen-2016-013665.
- Carmeli, Y., Troillet, N., Eliopoulos, G. M. and Samore, M. H. (1999) 'Emergence of antibiotic-resistant Pseudomonas aeruginosa: Comparison of risks associated with different antipseudomonal agents', *Antimicrob. Agents Chemother.*, 43(6): 1379–1382. doi: 10.1128/aac.43.6.1379.
- Carroll, K. C., Jorgensen, J. H. and Pfaller, M. A. (2015) *Manual of Clinical Microbiology*. Edited by L Landry Louise. ASM Press.
- CDC - Center for Disease Control and Prevention (2016) 'Bloodstream Infection Event (Central Line-Associated Bloodstream Infection and Non-central line-associated Bloodstream Infection)', *Published on-line: <http://www.cdc.gov/nhsn/>*, (January): 1–32. Available at: Centers for

Disease Control (CDC)/National Healthcare Safety Network (NHSN). Bloodstream Infection Event (Central Line-Associated Bloodstream Infection and Non-central line-associated Bloodstream Infection. Available at: <http://www.cdc.gov/nhsn/PDFs/pscM>.

Centers for Disease Control and Prevention (2015) 'National Center for Health Statistics. 2015. Deaths: final data for 2015.', *Natl. Vital Stat. Reports*, 6(6). doi: 10.1136/vr.h753.

Chan, L., Nor, M. B. M. and Arini, N. (2012) 'Guide to antimicrobial Therapy in the Adult ICU', *Malaysian Soc. Intens. Care*, p. 92. Available at: www.msic.org.my.

Chandrasekaran, S., Abbott, A., Campeau, S., Zimmer, B. L., Weinstein, M., Thrupp, L., *et al*(2018) 'Direct-from-blood-culture disk diffusion to determine antimicrobial susceptibility of gram-negative bacteria: Preliminary report from the clinical and Laboratory Standards Institute Methods Development and Standardization Working Group', *J. Clin. Microbiol.*, 56(3): 1–10. doi: 10.1128/JCM.01678-17.

CLSI (2020) *CLSI M100-ED29 : 2019 Performance Standards for Antimicrobial Susceptibility Testing , 29th Edition, CLSI*.

Coorevits, L., Boelens, J. and Claeys, G. (2015) 'Direct susceptibility testing by disk diffusion on clinical samples: a rapid and accurate tool for antibiotic stewardship', *Europe. J. Clin. Microbiol. Infect. Dis.*, 34(6): 1207–1212. doi: 10.1007/s10096-015-2349-2.

Diekema, D. J., Beekmann, S. E., Chapin, K. C., Morel, K. A., Munson, E. and Doern, G. V. (2003) 'Epidemiology and outcome of nosocomial and community-onset bloodstream infection', *J. Clin. Microbiol.*, 41(8): 3655–3660. doi: 10.1128/JCM.41.8.3655-3660.2003.

Drusano, G. L., Preston, S. L., Fowler, C., Corrado, M., Weisinger, B. and Kahn, J. (2004) 'Relationship between fluoroquinolone area under the curve:minimum inhibitory concentration ratio and the probability of eradication of the infecting pathogen, in patients with nosocomial pneumonia', *J. Infect. Dis.*, 189(9): 1590–1597. doi: 10.1086/383320.

- Engelkirk, P. G. and Duben-Engelkirk, J. L. (2003) 'Laboratory Diagnosis of Infectious Diseases: Essentials of Diagnostic Mycology', *Molecular Microbiology*.
- Erdem, I., Kucukercan, M. and Ceran, N. (2003) 'In vitro activity of combination therapy with cefepime, piperacillin-tazobactam, or meropenem with ciprofloxacin against multidrug-resistant *Pseudomonas aeruginosa* strains', *Chemother.*, 49(6): 294–297. doi: 10.1159/000074529.
- EUCAST (2020) *Reading guide; EUCAST disk diffusion method for antimicrobial susceptibility testing, Version 3.0 April*. Available at: <http://www.eucast.org>.
- Everett, M. J., Jin, Y. F., Ricci, V. and Piddock, L. J. V. (1996) 'Contributions of individual mechanisms to fluoroquinolone resistance in 36 *Escherichia coli* strains isolated from humans and animals', *Antimicrob. Agents Chemother.*, 40(10): 2380–2386. doi: 10.1128/aac.40.10.2380.
- Garcia, R. A., Spitzer, E. D., Beaudry, J., Beck, C., Diblasi, R., Gilleeny-Blabac, M., *et al* (2015) 'Multidisciplinary team review of best practices for collection and handling of blood cultures to determine effective interventions for increasing the yield of true-positive bacteremias, reducing contamination, and eliminating false-positive central line-a', *Am. J. Infect. Control*. Elsevier Inc, 43(11): 1222–1237. doi: 10.1016/j.ajic.2015.06.030.
- Garnacho-Montero, J., Garcia-Garmendia, J. L., Barrero-Almodovar, A., Jimenez-Jimenez, F. J., Perez-Paredes, C. and Ortiz-Leyba, C. (2003) 'Impact of adequate empirical antibiotic therapy on the outcome of patients admitted to the intensive care unit with sepsis', *Crit. Care Med.*, 31(12): 2742–2751. doi: 10.1097/01.CCM.0000098031.24329.10.
- Gashe, F., Mulisa, E., Mekonnen, M. and Zeleke, G. (2018) 'Antimicrobial Resistance Profile of Different Clinical Isolates against Third-Generation Cephalosporins', *J. Pharm.*, 2018: 1–7. doi: 10.1155/2018/5070742.
- Giamarellou, H. and Kanellakopoulou, K. (2008) 'Current Therapies for *Pseudomonas Aeruginosa*', *Crit. Care Clin.*, 24(2): 261–278. doi: 10.1016/j.ccc.2007.12.004.

- Goto, M. and Al-Hasan, M. N. (2013) 'Overall burden of bloodstream infection and nosocomial bloodstream infection in North America and Europe', *Clin. Microbiol. Infect.*, 19(6): 501–509. doi: 10.1111/1469-0691.12195.
- Hsu, L.-Y., Apisarnthanarak, A., Khan, E. and Suwantararat, N. (2017) 'Carbapenem-Resistant *Acinetobacter baumannii* and Enterobacteriaceae in South and Southeast Asia', *Clin. Microbiol.*, 30(1): 433–456. doi: 10.1007/978-1-62703-185-1_25.
- Ibrahim, E. H., Sherman, G., Ward, S., Fraser, V. J. and Kollef, M. H. (2000) 'The influence of inadequate antimicrobial treatment of bloodstream infections on patient outcomes in the ICU setting', *Chest*, 118(1): 146–155. doi: 10.1378/chest.118.1.146.
- Jorgensen, J. H. and Ferraro, M. J. (2009) 'Antimicrobial susceptibility testing: A review of general principles and contemporary practices', *Clin. Infect. Dis.*, 49(11): 1749–1755. doi: 10.1086/647952.
- Jorgensen, J. H., Maher, L. A. and Howell, A. W. (1991) 'Activity of meropenem against antibiotic-resistant or infrequently encountered gram-negative bacilli', *Antimicrob. Agents Chemother.*, 35(11): 2410–2414. doi: 10.1128/AAC.35.11.2410.
- Kang, C. I., Chung, D. R., Ko, K. S., Peck, K. R., Song, J. H. and Korean Network for Study of Infectious Diseases (2012) 'Risk factors for infection and treatment outcome of extended-spectrum β -lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* bacteremia in patients with hematologic malignancy.', *Ann. hematomol.*, 91(1): 115–121. doi: 10.1007/s00277-011-1247-7.
- Kang, C. I., Kim, S. H., Kim, H. Bin, Park, S. W., Choe, Y. J., Oh, M. D., Kim, E. C. and Choe, K. W. (2003) 'Pseudomonas aeruginosa bacteremia: Risk factors for mortality and influence of delayed receipt of effective antimicrobial therapy on clinical outcome', *Clin. Infect. Dis.*, 37(6): 745–751. doi: 10.1086/377200.
- Kim, E. S., Chong, Y. P., Park, S. J., Kim, M. N., Kim, S. H., Lee, S. O., Choi, S. H., Woo, J. H., Jeong, J. Y. and Kim, Y. S. (2017) 'Detection and genetic

- features of MCR-1-producing plasmid in human *Escherichia coli* infection in South Korea', *Diagn. Microbiol. Infect. Dis.*, 89(2): 158–160. doi: 10.1016/j.diagmicrobio.2017.06.020.
- Kola, A., Maciejewski, O., Sohr, D., Ziesing, S. and Gastmeier, P. (2007) 'Clinical impact of infections caused by ESBL-producing *E. coli*', *Scandinavian J. Infect. Dis.*, 39: 975–982. doi: 10.1080/00365540701466140.
- Kontula, K. S. K., Skogberg, K., Ollgren, J., Järvinen, A. and Lyytikäinen, O. (2018) 'The outcome and timing of death of 17,767 nosocomial bloodstream infections in acute care hospitals in Finland during 1999–2014', *Eur. J. Clin. Microbiol. Infect. Dis.*, 37(5): 945–952. doi: 10.1007/s10096-018-3211-0.
- Kumar, M., Shergill, S. P. S., Tandel, K., Sahai, K. and Gupta, R. M. (2019) 'Direct antimicrobial susceptibility testing from positive blood culture bottles in laboratories lacking automated antimicrobial susceptibility testing systems', *Med. J. Armed Forces India. Elsevier*, 75(4): 450–457. doi: 10.1016/j.mjafi.2018.08.010.
- Kutob, L. F., Justo, J. A., Bookstaver, P. B., Kohn, J., Albrecht, H. and Al-Hasan, M. N. (2016) 'Effectiveness of oral antibiotics for definitive therapy of Gram-negative bloodstream infections', *Int. J. Antimicrob. Agents. Elsevier B.V.*, 48(5): 498–503. doi: 10.1016/j.ijantimicag.2016.07.013.
- Lalitha, M. K. (2017) *Manual on Antimicrobial Susceptibility Testing. Departement of Microbiology Christian Medical College, Vellore, Tamil Nadu* (2017): 3–47. doi: 10.1002/9780470752579.ch15.
- Lamy, B., Dargère, S., Arendrup, M. C., Parienti, J. J. and Tattevin, P. (2016) 'How to optimize the use of blood cultures for the diagnosis of bloodstream infections? A state-of-the art', *Front. Microbiol.*, 7(MAY): 1–13. doi: 10.3389/fmicb.2016.00697.
- Laupland, K. B. (2013) 'Incidence of bloodstream infection: A review of population-based studies', *Clin. Microbiol. Infect.*, 19(6): 492–500. doi: 10.1111/1469-0691.12144.
- Li, X. Z., Plésiat, P. and Nikaido, H. (2015) 'The challenge of efflux-mediated antibiotic resistance in Gram-negative bacteria', *Clin. Microbiol.*, 28(2):

337–418. doi: 10.1128/CMR.00117-14.

- Liang, S. Y. and Kumar, A. (2015) ‘Empiric Antimicrobial Therapy in Severe Sepsis and Septic Shock: Optimizing Pathogen Clearance’, *Curr. Infect. Dis. Rep.*, 17(7). doi: 10.1007/s11908-015-0493-6.
- Linsenmeyer, K., Gupta, K., Strymish, J. M., Dhanani, M., Brecher, S. M. and Breu, A. C. (2016) ‘Culture if spikes? Indications and yield of blood cultures in hospitalized medical patients’, *J. Hosp. Med.*, 11(5): 336–340. doi: 10.1002/jhm.2541.
- Livermore, M. D.,(2012) ‘Current Epidemiology and Growing Resistance of Gram- Negative Pathogens’, *Korean J. Intern Med.*, 27: 128–142. doi: <http://dx.doi.org/10.3904/kjim.2012.27.2.128> ORIGIN
- Lodise, T. P., Patel, N., Kwa, A., Graves, J., Furuno, J. P., Graffunder, E., Lomaestro, B. and McGregor, J. C. (2007) ‘Predictors of 30-day mortality among patients with *Pseudomonas aeruginosa* bloodstream infections: Impact of delayed appropriate antibiotic selection’, *Antimicrob. Agents Chemother.*, 51(10): 3510–3515. doi: 10.1128/AAC.00338-07.
- Marchello, C. S., Dale, A. P., Pisharody, S., Rubach, M. P. and Crump, J. A. (2020) ‘A Systematic Review and Meta-analysis of the Prevalence of Community-Onset Bloodstream Infections among Hospitalized Patients in Africa and Asia’, *Antimicrob. Agents Chemother.*, 64(1): 1–16. doi: 10.1128/AAC.01974-19.
- Marshall, W. F. and Blair, J. E. (1999) ‘The cephalosporins’, *Mayo Clinic Proceedings*, 74(2): 187–195. doi: 10.4065/74.2.187.
- Mauro, M. V., Dodaro, S., Perugini, D., Greco, F. and Giraldi, C. (2017) ‘Introduction of Gram negative microorganisms rapid identification and direct susceptibility testing to reduce turnaround time in blood cultures’, *Microbiol. Medica*, 32(4): 145–148. doi: 10.4081/mm.2017.6963.
- McNamara, J. F., Righi, E., Wright, H., Hartel, G. F., Harris, P. N. A. and Paterson, D. L. (2018) ‘Long-term morbidity and mortality following bloodstream infection: A systematic literature review’, *J. Infect.* Elsevier Ltd, 77(1): 1–8. doi: 10.1016/j.jinf.2018.03.005.

- Munford, R. S. (2006) 'Severe sepsis and septic shock: The role of gram-negative bacteremia', *Ann. Rev. Pathol.*, 1: 467–496. doi: 10.1146/annurev.pathol.1.110304.100200.
- Pan, H. W., Li, W., Li, R. G., Li, Y., Zhang, Y. and Sun, E. H. (2018) 'Simple sample preparation method for direct microbial identification and susceptibility testing from positive blood cultures', *Front. Microbiol.*, 9(3): 1–9. doi: 10.3389/fmicb.2018.00481.
- Permenkes RI (2011) 'Peraturan Menteri Kesehatan Republik Indonesia Nomor 2406/MENKES/PER/XII', *Pedoman Umum Penggunaan Antibiotik*: 34–44.
- Perry, J. D. (2017) 'A Decade of Development of Chromogenic Culture Media for Clinical Microbiology in an Era of Molecular Diagnostics', *Clin. Microbiol. Rev.* 30(2): 449–479. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/28122803>.
- Peters, R. P. H., Van Agtmael, M. A., Gierveld, S., Danner, S. A., Groeneveld, A. B. J., Vandenbroucke-Grauls, C. M. J. E. and Savelkoul, P. H. M. (2007) 'Quantitative detection of *Staphylococcus aureus* and *Enterococcus faecalis* DNA in blood to diagnose bacteremia in patients in the intensive care unit', *J. Clin. Microbiol.*, 45(11): 3641–3646. doi: 10.1128/JCM.01056-07.
- Pincus, D. H. (2010) 'Microbial identification using the bioMérieux VITEK® 2 system', *Encycl. Rapid Microbiol. Methods*: 1–32.
- Retamar, P., Portillo, M. M., López-Prieto, M. D., Rodríguez-López, F., De Cueto, M., García, M. V., Gómez, M. J., Del Arco, A., Muñoz, A., Sánchez-Porto, A., Torres-Tortosa, M., *et al*(2012) 'Impact of inadequate empirical therapy on the mortality of patients with bloodstream infections: A propensity score-based analysis', *Antimicrob. Agents Chemother.*, 56(1): 472–478. doi: 10.1128/AAC.00462-11.
- Rhomberg, P. R., Fritsche, T. R., Sader, H. S. and Jones, R. N. (2005) 'Comparative antimicrobial potency of meropenem tested against gram-negative bacilli: Report from the MYSTIC Surveillance Program in the United States (2004)', *J. Chemother.*, 17(5): 459–469. doi: 10.1179/joc.2005.17.5.459.

- Ruppé, É., Woerther, P. L. and Barbier, F. (2015) 'Mechanisms of antimicrobial resistance in Gram-negative bacilli', *Ann. Intensive Care*. Springer Paris, 5(1). doi: 10.1186/s13613-015-0061-0.
- Ryan, K. J., Nafees, A., W. Lawrence, D., Michael, L., C. George, R., Paul, P., L. Barth, R. and Charles R, S. (2018) '*Sherris Medical Microbiology*, Seventh Edition': 431–458.
- Shanthraju, L. and Devi, G. (2016) 'Evaluation of direct sensitivity testing as a method for early initiation of treatment in Gram-negative sepsis', *J. Acad. Clin. Microbiol.*, p. 131. doi: 10.4103/0972-1282.194953.
- Stefaniuk, E., Mrówka, A. and Hryniewicz, W. (2005) 'Susceptibility testing and resistance phenotypes detection in bacterial pathogens using the VITEK 2 system', *Polish J. Microbiol.*, 54(4): 311–316.
- Stryjewski, M. E. and Boucher, H. W. (2009) 'Gram-negative bloodstream infections.', *Int. J. Antimicrob. Agents*. Elsevier B.V., 34 Suppl 4: S21–S25. doi: 10.1016/s0924-8579(09)70561-8.
- Timbrook, T. T., Morton, J. B., Mcconeghy, K. W., Caffrey, A. R., Mylonakis, E. and LaPlante, K. L. (2017) 'The effect of molecular rapid diagnostic testing on clinical outcomes in bloodstream infections: A systematic review and meta-analysis', *Clin. Infect. Dis.*, 64(1): 15–23. doi: 10.1093/cid/ciw649.
- U.S. Department of Health and Human Services Food and Drug Administration (2009) '*Guidance for Industry and FDA. Class II Special Controls Guidance Document: Antimicrobial Susceptibility Test (AST) Systems*': 1–42. Available at: <https://www.fda.gov>.
- Uslan, D. Z. (2007) 'Age and Sex-Associated Trends in Bloodstream Infection', *Arch. Intern. Med.*, 167(8), p. 834. doi: 10.1001/archinte.167.8.834.
- Viera, A. J. and Garrett, J. M. (2005) 'Anthony J. Viera, MD; Joanne M. Garrett, PhD (2005). Understanding interobserver agreement: the kappa statistic. *Fam Med* 2005;37(5):360-63.', *Fam. Med.*, 37(5): 360–3. Available at: http://www1.cs.columbia.edu/~julia/courses/CS6998/Interrater_agreement.Kappa_statistic.pdf.
- Wang, X. and Quinn, P. J. (2010) 'Lipopolysaccharides of Gram-Negative

Bacteria', *Curr. Top.. Membr. Trans.*, 17(C): 79–151. doi: 10.1016/S0070-2161(08)60309-3.

Wootton, M. (2013) 'BSAC Methods for Antimicrobial Susceptibility Testing', *Bsac*, 44(May): 1–87. Available at: http://bsac.org.uk/wp-content/uploads/2012/02/Version-12-Apr-2013_final.pdf.

Yakoob, Rahila ; Bhat, G. (2017) 'Comparison of Direct Antibiotic Susceptibility Testing with Standard Testing in Blood Culture', *J. Pure Appl. Microbiol.*, 1(1): 2289–2296. doi: 10.23959/sfahrj-1000006.

Yakoob, Rahila ; Bhat, G. (2018) 'Research Article', *Comparision of Direct Antibiotic Susceptibility Testing with Standard Testing in Blood Culture*, 1(1): 2289–2296. doi: 10.23959/sfahrj-1000002.