

LIST OF CONTENT

	Page
List of Tables	ix
List of Figures	x
List of Appendicies	xiv
Abstract	xv
Intisari	xvi
 CHAPTER I. GENERAL INTRODUCTION	
1. Background	1
2. Research Objectives	5
3. Research Benefit	5
4. Research Novelty	6
 CHAPTER II. LITERATURE REVIEW	
1. Systematics of Shallot	7
2. Fungal Foliar Diseases of Shallot	
2.1. Purple blotch	8
2.2. Stemphylium leaf blight	14
2.3. Anthracnose	16
3. Plant Associated Bacteria	
3.1. Description	17
3.2. Variability of endophytic bacteria	18
3.3. Plant Associated Bacteria as biocontrol agent	19
4. Molecular Identification	
4.1. <i>Alternaria porri</i> identification	20
4.2. Plant Associated Bacteria identification	22
5. Biological Control	
5.1. Antagonism	23
5.2. Systemic induce resistance	27
6. Theoretical Research Framework	30
7. Hypothesis	31

CHAPTER III. TOPIC 1. IDENTIFICATION OF PRPLE BLOTCH	
PATHOGEN OF SHALLOT BY PCR USING SPECIFIC PRIMER	
FOR ALTERNARIA GENUS	
1. Introduction	33
2. Materials and Methods	
2.1. Diseased leaf samples collection	34
2.2. Purple blotch pathogen isolation	35
2.3. Isolates fingerprinting by ERIC-PCR and BOX-PCR	35
2.4. Molecular identification using Alternaria specific primers	36
2.5. Colony morphology characterization	37
2.6. Pathogenicity test of the isolates	37
3. Results and Discussion	
3.1. Diseased leaf samples collection	37
3.2. Purple blotch pathogen isolation	39
3.3. Isolates fingerprinting by ERIC-PCR and BOX-PCR	39
3.4. Molecular identification using Alternaria specific primers	41
3.5. Colony morphology characterization	45
3.6. Pathogenicity test of the isolates	47
CHAPTER IV. TOPIC 2. EXPLORATION AND CHARACTERIZATION OF	
PLANT ASSOCIATED BACTERIA FROM SHALLOT LEAVES	
FOR FUNGAL FOLIAR DISEASES CONTROL ON SHALLOT	
1. Introduction	49
2. Materials and Methods	
2.1. Plant Associated Bacteria isolation	50
2.2. Antagonistic test of Plant Associated Bacteria against	
<i>Alternaria porri</i> in vitro	51
2.3. Selection of antagonist and non-antagonist Plant Associated	
Bacteria	52
2.4. Hypersensitive response (HR) test of plant associated	
bacteria isolates	53
2.5. Identification of selected bacteria isolates	54
2.6. Compatibility growth between antagonistic Plant Associated	
Bacteria combination in vitro.....	59
2.7. Growth inhibition of <i>A. porri</i> by antagonistic Plant Associated	
Bacteria as single isolate and as combination isolates in	
vitro	59

2.8. Antagonist test of six chosen PAB isolates against <i>A. porri</i> (KP10), <i>A. alternata</i> (B1), <i>Stemphylium</i> sp. (B2, B14, B17, T4, GS2, and GS4), and <i>Colletotrichum</i> sp. in vitro	60
2.9. Extracellular hydrolytic chitinase and glucanase assay	60
2.10. Antagonist assay of Cell Free Supernatant (CFS), SDS-CFS and heat treatment-CFS	61
2.11. Characterization of antagonist metabolite in CFS by using GC-MS method	61
3. Results and Discussion	
3.1. Plant Associated Bacteria (PAB) isolation	62
3.2. Antagonistic test of Plant Associated Bacteria against <i>Alternaria porri</i> in vitro	65
3.3. Selection of antagonist and non-antagonist PAB.....	68
3.4. Hypersensitive Response (HR) test	72
3.5. Identification of selected bacteria isolates	73
3.6. Compatibility growth between antagonistic Plant Associated Bacteria combination in vitro	78
3.7. Growth inhibition of <i>A. porri</i> by antagonistic Plant Associated Bacteria as single isolate and as combination isolates in vitro	79
3.8. Antagonist test of six chosen PAB isolates against <i>A. porri</i> (KP10), <i>A. alternata</i> (B1), <i>Stemphylium</i> sp. (B2, B14, B17, T4, GS2, and GS4), <i>Colletotrichum</i> sp. in vitro	80
3.9. Extracellular hydrolytic chitinase and glucanase assay	81
3.10. Antagonist assay of Cell Free Supernatant (CFS), SDS-CFS and Hot treatment-CFS	83
3.11. Characterization of antagonist metabolite in CFS by using GC-MS method	85
 CHAPTER V. TOPIC 3. BACILLUS SPECIES AND <i>Pseudomonas aeruginosa</i> AS POTENTIAL AGENTS OF INDUCE RESISTANCE ON SHALLOT AGAINST <i>Alternaria porri</i>	
1. Introduction	91
2. Materials and Methods	
2.1. Experimental design	92
2.2. Experimental plant preparation	93
2.3. Plant Associated Bacteria (PAB) preparation and inoculation	93
2.4. Pathogen inoculation	93

2.5. Analysis of total phenolic compounds	94
2.6. Peroxidase activity assessment	94
2.7. Salicylic Acid analysis	95
3. Results and Discussion	
3.1. Disease incidence	95
3.2. Total phenolic (Ph) compounds content of leaves	97
3.3. Peroxidase activity	99
3.4. Salicylic Acid content accumulation	103
CHAPTER VI. TOPIC 4. PERFORMANCE OF INDIGENOUS PLANT ASSOCIATED BACTERIA IN CONTROLLING FUNGAL FOLIAR DISEASES ON SHALLOT	
1. Introduction	108
2. Materials and Methods	
2.1. Experimental design	109
2.2. Experiment 1: bulb dipping plus soil drenching method	110
2.3. Experiment 2: foliar spraying method	110
2.4. Plant Associated Bacteria preparation	111
2.5. Experimental plant preparation and crops maintenance	111
2.6. Parameter assessed and data analysis	112
3. Results and Discussion	
3.1. Disease symptom and disease incidence	112
3.2. Germination percentage	118
3.3. Vegetative growth parameter: plant height and number of leaves	119
3.4. Shallot yield	120
CHAPTER VII. GENERAL DISCUSSION	123
CHAPTER VIII. CONCLUSSION AND SUGGESTION	127
SUMMARY	129
RINGKASAN	141
REFERENCES	153
APPENDICIES	184

LIST OF TABLE

Table	page
3.1. Table Survey location for samples collection	34
3.2. List of primers used in this study	36
3.3. Description of colony morphology of collected isolates	39
3.4. Characteristic of colony morphology of 41 fungal isolates from shallot leaves showing purple blotch symptom	47
4.1. Survey location and sampling of shallot crops for bacteria isolation	64
4.2. Number of antagonist and non antagonist Plant Associated Bacteria isolated from Bantul and Nganjuk Regencies at dry and wet season	67
4.3. Information of the six antagonist isolates and two non-antagonist isolates (isolate 24 & 35) selected for further experiment	73
4.4. Morphological and biochemical characteristic of eight selected isolates of Plant Associated Bacteria from shallot	74
4.5. Compatibility test of the six antagonist PAB isolates of shallot	78
4.6. ANOVA of inhibition percentage of nine fungal isolates from shallot purple blotch symptom by six indigenous antagonistic bacteria isolates of shallot	82
4.7. Assessment of bacterial CFS antagonist test against <i>A. porri in vitro</i> at seven days incubation in well method assay	83
4.8. Fungal growth diameter (mm) at seven days incubation on bacterial CFS antagonist test against <i>A. porri in vitro</i> in pour plate method assay	83
4.9. Compounds identified in ethyl acetate extract of Cell Free Supernatant (CFS) of Isolate 60 bacteria (<i>B. methylotrophicus</i>) by GC-MS	88
4.10. Compounds identified in ethyl acetate extract of Cell Free Supernatant (CFS) of Isolate 9 bacteria (<i>B. subtilis</i>) by GC-MS	89

LIST OF FIGURES

Figure	Page
1.1. Shallot production, harvest area, and Productivity in Indonesia from 2008 – 2016 (BPS - Statistics Indonesia and Directorate General of Horticulture/ Badan Pusat Statistik dan Direktorat Jenderal Hortikultura)	1
2.1. Shallot bulb	7
2.2. Multiplier onion bulb	7
2.3. Onion bulb	7
2.4. Purple blotch symptom: (a) on onion leaves (Black <i>et al.</i> , 2012); (b) on shallot leaves (Survey on CIAT-UGM experimental field, 2013); (c) on the neck of shallot bulb (Survey on Bantul, 2014)	8
2.5. Spore of <i>Alternaria porri</i>	9
2.6. Dark brown to black sporulation on leaves caused by <i>Stemphylium</i> sp. (Black, <i>et al.</i> , 2012)	14
2.7. Spore of <i>Stemphylium vesicarium</i>	15
2.8. Symptom of anthracnose disease on shallot	16
2.9. Conidia of <i>Colletotrichum</i> sp.	17
3.1. Purple blotch symptom on shallot leaves	38
3.2. In-situ conidia resembling <i>Alternaria</i> sp. observed from shallot crops in CIAT-UGM experimental field and in Donotirto villages, Bantul Regency, Special Region of Yogyakarta. Bar represents 100 μ m	38
3.3. In-situ conidia resembling <i>Stemphylium</i> sp. observed from shallot crops in Tirtohargo and Gading Sari villages, Bantul Regency, Special Region of Yogyakarta, Bar represents 100 μ m	38
3.4. Dendogram showing relatedness of 13 fungal isolates from shallot leaves having purple blotch symptom based on rep-PCR using BOX A1R primer	40
3.5. Dendogram showing relatedness of 13 fungal isolates from shallot leaves having purple blotch symptom based on rep-PCR using ERIC2 primer	40
3.6. Dendogram showing relatedness of 13 fungal isolates from shallot leaves having purple blotch symptom based on accumulation of ERIC2 and BOX PCR Analysis	40

Figure	Page
3.7. PCR product of 13 fungal isolates from shallot having purple blotch symptom and 1 isolate of <i>Fusarium</i> sp. as control on 1 % agarose gel	41
3.8. Phylogenetic tree constructed by using Neighbor Joining Tree statistical method through 1000 replication of Bootstrap test method and Maximum Composite Likelihood model based on sequences of ITS1 and ITS2 region using Dir1ITSAlt/Inv1ITSAlt primers specific for <i>Alternaria</i> genus	42
3.9. Pattern of nucleotide sequence comparison of 10 published sequences and 13 collected isolate sequences generated by Multalin version 5.4.1 program.....	42
3.10. Colony morphology of 13 fungal isolates of shallot from upper & reverse side	46
3.11. PCR product of fungal isolates re-isolated from shallot leaves on pathogenicity test on 1% agarose gel	48
4.1. Antagonist test of Plant Associated Bacteria against <i>Alternaria porri</i> on PDA.	52
4.2. Abundance of cultured-able Bacteria isolated from farmers shallot crops in Bantul and Nganjuk Districts at dry and wet season from healthy and diseased shallot plants	63
4.3. Representative of 14 colony morphology under light microscope at 100x magnification	66
4.4. BOX-PCR result showing variation in band pattern of 53 isolates in figures (A), (B), (C), and (D)	69
4.5. Grouping of 53 isolates based on BOX-PCR and analysis with NTsys program.....	70
4.6. Analysis of 26 isolates representing each group in the 1 st , 2 nd , & 3 rd analysis	71
4.7. Hypersensitive reaction of isolate 7, isolate 8, and isolate 28 that causing necrosis on tobacco leaf at the sixth day after inoculation	72
4.8. Phylogenetic tree constructed based on 16S rRNA gene fragment sequence of the eight indigenous isolates and with 18 published sequences.....	77
4.9. Inhibition of <i>A. porri</i> by six isolates and two combinations of isolates in PDA plate at seven days after treatment	79
4.10. Light microscopy figures of disrupted hyphae on the antagonist zone of <i>Bacillus</i> species and <i>A. porri</i> at seven days after treatment, (a) normal hyphae; (b) hyphae showing excessive vacuole; (c) cells malformation. Bar representing 100 μ m	80

Figure	Page
4.11. Inhibition of <i>A. porri</i> at day-7 after treatment by CFS of <i>B. methylotrophicus</i> (isolate 60) and <i>B. subtilis</i> (isolate 9): (A) isolate 60 CFS-heat treatment; (B) isolate 60 CFS-SDS treatment; (C) isolate 9 CFS-heat treatment; (D) isolate 9 CFS-SDS treatment; (E) control	85
4.12. Chromatogram of ethyl acetate extract of Cell Free Supernatant of 48-hour culture on Nutrient Broth of bacteria isolate 60 (<i>B. methylotrophicus</i>) identified through GC-MS	86
4.13. The Gas Chromatogram of ethyl acetate extract of Cell Free Supernatant of 48-hour culture on Nutrient Broth of bacteria isolate 9 (<i>Bacillus subtilis</i>) identified through GC-MS	87
5.1. Disease incidence on shallot at day-5 post inoculation; treatment having same notation in white letter was not significantly different in DMRT 5%	96
5.2. Total phenol concentration in shallot leaves treated with Plant Associated Bacteria <i>Bacillus methylotrophicus</i> , <i>B. subtilis</i> , <i>Pseudomonas aeruginosa</i> , <i>B. aerophilus</i> and <i>B. tequilensis</i> , and inoculated with <i>Alternaria porri</i>	98
5.3. Peroxidase activity in shallot leaves treated with Plant Associated Bacteria <i>Bacillus methylotrophicus</i> , <i>B. subtilis</i> , <i>Pseudomonas aeruginosa</i> , <i>B. aerophilus</i> and <i>B. tequilensis</i> , and inoculated with <i>Alternaria porri</i> assessed at 1, 3, and 5 days after <i>A. porri</i> inoculation	102
5.4. Salicylic acid concentration in shallot leaves assessed at day-5 post <i>Alternaria porri</i> inoculation	103
6.1. Shallot plant showing anthracnose disease caused by <i>Colletotrichum gloeosporioides</i>	113
6.2. Anthracnose disease incidence curve of shallot crops treated with PAB	115
6.3. Area Under Disease Progress Curve (AUDPC) of disease incidence of shallot crops treated with PAB	116
6.4. Germination of shallot bulb after introduced with PAB isolates ...	119
6.5. Plant height (A) and number of leaves (B) of shallot plants treated with PAB in bulb dipping plus soil drenching method experiment	121
6.6. Plant height (A) and number of leaves (B) of shallot plants treated with PAB in foliar spraying method experiment	121

Figure	Page
6.7. Fresh weight of shallot bulb and leaves and dry weight of shallot bulb (Kg m ²) in bulb dipping plus soil drenching method experiment	122
6.8. Fresh weight of shallot bulb and leaves and dry weight of shallot bulb (Kg m ²) in foliar spraying method experiment	122