

## DAFTAR PUSTAKA

- Andersson O, Hellström-Westas L, Andersson D, *et al.* (2011). Effect of delayed versus early umbilical cord clamping on neonatal outcomes and iron status at 4 months: a randomised controlled trial. *British Medical Journal*. pp.343:1–12.
- Atladdottir H & Thorsdottir I. (2000). Energy intake and growth of infants in Iceland—a population with high frequency of breast-feeding and high birth weight. *European Journal of Clinical Nutrition*. pp.54:695–701.
- Bekri, S., Gual, P., Anty, R., *et al.* (2006). Increased Adipose Tissue Expression of Hepcidin in Severe Obesity Is Independent From Diabetes and NASH. *Gastroenterology*, 131(3), pp.788-796.
- Bothwell TH. (2000). Iron requirements in pregnancy and strategies to meet them. *The American Journal of Clinical Nutrition*; 72: 257S–264S.
- Calvo EB, Galindo AC, Aspres NB. (1992). Iron status in exclusively breast-fed infants. *The Journal of Pediatrics*. ;90:375–379.
- Centers for Disease Control and Prevention. (1998). Recommendations to prevent and control iron deficiency in the United States. *MMWR Recommendations and Reports*.;47(RR-3):1–29.
- Ceriani Cernadas JM, Carroli G, Pellegrini L, *et al.* ( 2010). The effect of early and delayed umbilical cord clamping on ferritin levels in term infants at six months of life: A randomized controlled trial. *Archivos Argentinos De Pediatría Journals*. pp.108:201–208.
- Chantry CJ, Howard CR, Auinger P. (2007). Full breastfeeding duration and risk for iron deficiency in U.S. infants. *Breastfeed Medical Journal*.pp.2:63–73.
- Dallman PR. (1989). Review of iron metabolism. In: Filer LJ Jr, editor. *Dietary iron: Birth to two years*. New York: Raven Press:1-17
- De Benoist B, Egli I, Cogswell M. (2008). *Worldwide prevalence of anaemia 1993–2005*. Geneva: World Health Organization.
- Departemen Kesehatan. (2013). *Laporan Riset Kesehatan Dasar (Riskesdas) 2010 Bidang Biomedis*. Jakarta: Badan Litbangkes, Depkes RI.
- Departemen Kesehatan. (2001). *Survey Kesehatan Nasional. Laporan*. Depkes RI Jakarta.

- Desforges, J. and Oski, F. (1993). Iron Deficiency in Infancy and Childhood. *New England Journal of Medicine*. pp.190-193.
- Dewey KG, Cohen RJ, Rivera LL, *et al.* (1998). Effects of age of introduction of complementary foods on iron status of breast-fed infants in Honduras. *The American Journal of Clinical Nutrition*.pp.67:878–84.
- Douglas, SE. (2011). *Maternal and Infant Iron Status and First Year Illness. Dietics and Nutrition*. The Graduate Faculty of the University of Kansas.
- Earl R, Woteki CE, eds. (1993). *Iron Deficiency Anemia: Recommended Guidelines for the Prevention, Detection, and Management Among U.S. Children and Women of Childbearing Age*. Washington, DC: The National Academies Press.
- Eskeland B, Baerheim A, Ulvik R, Hunskaar S. (2002). Influence of mild infections on iron status parameters in women of reproductive age. *Scandinavian Journal of Primary Health Care*. pp.20: 50–6
- Ferrucci L, Semba RD, Guralnik JM, *et al.* (2010). Proinflammatory state, hepcidin, and anemia in older persons. *Blood Journal*. pp.115: 3810-6.
- Finch C.A., Bellotti V., Stray S., Lipschitz D.A., Cook J.D., Pippard M.J., Huebers H.A. (1986). Plasma ferritin determination as a diagnostic tool. *The Western Journal of Medicine*. pp.145: 657–663.
- Fomon, S., Nelson, S. and Ziegler, E. (2000). Retention of iron by infants. *Annual Review of Nutrition*, 20(1), pp.273-290.
- Gale, E., Torrance, J. & Bothwell, T. (1963). The quantitative estimation of total iron stores in human bone marrow. *Journal of Clinical Investigation*. pp.1076–1082.
- Galesloot TE, Vermeulen SH, Geurts-Moespot AJ, *et al.* (2011). Serum hepcidin: reference ranges and biochemical correlates in the general population. *Blood Journal*; 117(25): e218-e225.
- Gambling, L., Lang, C. and McArdle, H. (2011). Fetal regulation of iron transport during pregnancy. *The American Journal of Clinical Nutrition*. pp.1903S-1907S.
- Garcia-Casal, M.N., Peña-Rosas, J.P. & Pasricha, S.-R. (2014) Rethinking ferritin cutoffs for iron deficiency and overload. *The Lancet Haematology Journal*. 1, e92–e94.
- Georgieff, M., Wewerka, S., Nelson, C. and de Regnier, R. (2002). Iron status at 9 months of infants with low iron stores at birth. *The Journal of Pediatrics*, 141(3), pp.405-409. 6]

- Guyatt G.H., Oxman A.D., Ali M., Willan A., McIlroy W., Patterson C. *et al.* (1992). Laboratory diagnosis of iron-deficiency anemia: an overview. *The Journal of General Internal Medicine*. pp.7:145–153.
- Hagnevik K, Faxelius G, Irestedt L, *et al.* (1984). Catecholamine surge and metabolic adaptation in the newborn after vaginal delivery and caesarean section. *Acta Paediatrica Scandinavica Journal*. pp.73:602–9.
- Haschke F, Vanura H, Male C, Owen G, Pietschnig B, Schuster E, *et al.* (1993). Iron nutrition and growth of breast- and formula-fed infants during the first 9 months of life. *Journal of Pediatric Gastroenterology and Nutrition*. pp.16(2):151e6.
- Hay G, Refsum H, Whitelaw A, Melbye EL, Haug E, Borch-Johnsen B. *et al.* (2007). Predictors of serum ferritin and serum soluble transferrin receptor in newborns and their associations with iron status during the first 2 years of life. *The American Journal of Clinical Nutrition*. pp.86:64–73.
- Hentze MW, Muckenthaler MU, Galy B, Camaschella C. (2010). Two to tango: regulation of mammalian iron metabolism. *The Journal Cell*. pp.142: 24-38.
- Ihyauddin, Z., (2014). Korelasi Antara Kadar Feritin Serum Ibu Melahirkan dengan Kadar Feritin Serum Bayi Baru Lahir di RS PKU Muhammadiyah Bantul. Skripsi. Fakultas Kedokteran Universitas Gadjah Mada.
- Innis SM, Nelson CM, Wadsworth LD, MacLaren IA, Lwanga D. *et al.* (1997). Incidence of iron-deficiency anaemia and depleted iron stores among nine-month-old infants in Vancouver, Canada. *Canadian Journal of Public Health*. pp.88(2):80e4.
- Irestedt L, Dahlin I, Hertzberg T, *et al.* (1989). Adenosine concentration in umbilical cord blood of newborn infants after vaginal delivery and cesarean section. *The Journal of Pediatric Research*. pp.26:106–8.
- Jonsdottir OH, Thorsdottir I, Hibberd PL, *et al.* (2012). Timing of the introduction of complementary foods in infancy: a randomized controlled trial. *Journal of Pediatrics*. pp.130:1038–45.
- Kassebaum NJ, Jasrasaria R, Naghavi M, *et al.* (2014). A systematic analysis of global anemia burden from 1990 to 2010. *Blood Journal*. 123: 615–24.
- Koehler K, Braun H, Achtzehn S, *et al.* (2012). Iron status in elite young athletes: gender-dependent influences of diet and exercise. *European Journal of Applied Physiology*. pp.112: 513–23.
- Koller O. (1982). The clinical significance of hemodilution during pregnancy. *Obstretical and Gynecological Survey*. pp.37 649–652

- Kratz A., Ferraro M., Sluss P.M., Lewandrowski K.B. *et al.* (2004). Case records of the Massachusetts General Hospital. Weekly clinicopathological exercises. Laboratory reference values. The New England Journal of Medicine. pp.351 1548–1563.
- Kulaksiz H, Theilig F, Bachmann S, *et al.* (2005). The iron-regulatory peptide hormone hepcidin: expression and cellular localization in the mammalian kidney. *The Journal of Endocrinology*; 184: 361–70
- Lapillonne, A. and Becquet, O. (2017). Recommandations pour les apports en fer pendant les six premiers mois de vie. *Archives de Pédiatrie Journal*, 24(5), pp.5S40-5S44.
- Letsky E. A. (1995). Erythropoiesis in pregnancy. *The Journal of Perinatal Meicine*. 23 39–45
- Marques RF, Taddei JA, Lopez FA, Braga JAP. (2014). Breastfeeding exclusively and iron deficiency anemia during the first 6 months of age. *Revista Da Associacao Medica Brasileria*. pp. 60:18–22.
- Maryani, A.M. (2015). Korelasi Antara Kadar Feritin Serum Ibu Hamil Aterm dengan Kadar Feritin Serum Bayi Usia Tiga Bulan di RS PKU Muhammadiyah Bantul. Skripsi. Fakultas Kedokteran Universitas Gadjah Mada.
- McLean E, Cogswell M, Egli I, Wojdyla D, de Benoist B. (2009). Worldwide prevalence of anaemia, WHO Vitamin and Mineral Nutrition Information System, 1993-2005. *The Journal of Public Health Nutrition*. 12: 444-54.
- Mei Z, Cogswell ME, Looker AC, Pfeiffer CM, Cusick SE, Lacher DA, *et al.* (2011). Assessment of iron status in US pregnant women from the National Health and Nutrition Examination Survey (NHANES), 1999-2006. *The American Journal of Clinical Nutrition*. 93:1312-20.
- Merle U, Fein E, Gehrke SG, Stremmel W, Kulaksiz H. (2007). The iron regulatory peptide hepcidin is expressed in the heart and regulated by hypoxia and inflammation. *Journal of Endocrinology*; 148: 2663–68.
- Milman N, Ibsen KK, Christensen JM. (1987). Serum ferritin and iron status in mothers and newborn infants. *Acta Obstetricia et Gynecologica Scandinavia*; 66: 205-11.
- Milman, N. (2011). Iron in pregnancy – how do we secure an appropriate iron status in mother and child?. *Annals of Nutrition & Metabolism*: Page: 1-12.

- Moschovis PP, Banajeh S, MacLeod WB, *et al.* (2013). Childhood anemia at high altitude: risk factors for poor outcomes in severe pneumonia. *The Journal of Pediatrics*. 132:e1156–62.
- Nybo M, Friis-Hansen L, Felding P, Milman N. (2007). Higher prevalence of anaemia among pregnant immigrant women compared to pregnant ethnic Danish women. *Annals Hematology Journal*; 86: 647–651.
- Olaya, G., Lawson, M. and Fewtrell, M. (2017). Iron Status at Age 6 Months in Colombian Infants Exclusively Breast-fed for 4 to 5 Versus 6 Months. *Journal of Pediatric Gastroenterology and Nutrition*, 64(3), pp.465-471.
- Pacheco L., Costantine M. M, Hankins G. D. V. (2013). “Physiologic changes during pregnancy,” in *Clinical Pharmacology During Pregnancy* ed. Mattison D. R., editor. (San Diego: Academic Press; )
- Paiva, A., Rondó, P., Pagliusi, R., Latorre, M., Cardoso, M. and Gondim, S. (2007). Relationship between the iron status of pregnant women and their newborns. *Revista de Saúde Pública*, 41(3), pp.321-327.
- Pasricha SR, Drakesmith H, Black J, Hipgrave D, Biggs BA. (2013). Control of iron deficiency anemia in low- and middle-income countries. *Blood Journal*; 121: 2607-17.
- Pavord S., Myers B., Robinson S., Allard S., Strong J., Oppenheimer C. (2012). UK guidelines on the management of iron deficiency in pregnancy. *The British Journal of Haematology*. 156: 588-600.
- Peeling P, Sim M, Badenhorst CE, *et al.* (2014). Iron status and the acute post-exercise hepcidin response in athletes. *PLoS One*; 9: e93002.
- Pigeon C, Ilyin G, Courselaud B, *et al.* (2001). A new mouse liver-specific gene, encoding a protein homologous to human antimicrobial peptide hepcidin, is overexpressed during iron overload. *The Journal of Biological Chemistry*; 276: 7811–19.
- Pisacane A, De Vizia B, Valiante A, Vaccaro F, Russo M, Grillo G, *et al.* (1995). Iron status in breast-fed infants. *The Journal of Pediatrics*;127(3):429e31.
- Pizarro, F., Yip, R., Dallman, P., Olivares, M., Hertrampf, E. and Walter, T. (1991). Iron status with different infant feeding regimens: Relevance to screening and prevention of iron deficiency. *The Journal of Pediatrics*, 118(5), pp.687-692.
- Reeves JD, Yip R, Kiley VA, Dallman PR. (1984). Iron deficiency in infants: the influence of mild antecedent infection. *The Journal of Pediatrics*; 105: 874–9

- Rios, E, Lipschitz, DA, Cook, JD & Smith, NJ. (1975). Relationship of Maternal and infant iron stores as Assessed by Determination of plasma Ferritin. *Pediatrics: Official Journal of the American Academy of Pediatrics*, Vol.55, No 5. pp 694.
- Schifman RB, James E, Thomasson A, Evers JM. (1987). Red blood zinc protoporphyring testing for iron deficiency anemia. *The American Journal of Obstetrics and Gynecology*; 157: 304-7.
- Scholl TO. (2005). Iron status during pregnancy: setting the stage for mother and infant. *The American Journal of Clinical Nutrition*. 81:1218S-1222S.
- Scholl, T.S. (2011). Maternal iron status: relation to fetal growth, length of gestation, and iron endowment of the neonate. *Nutrition Reviews*; 69(1): S23-S29.
- Scott-Emuakpor AB, Okola AA, Omene JA, Ukpe SI. (1985). Normal haematological values of the African neonate. *Blut Journal*; 11-18.
- Siddappa, A., Rao, R., Long, J., Widness, J. and Georgieff, M. (2007). The Assessment of Newborn Iron Stores at Birth: A Review of the Literature and Standards for Ferritin Concentrations. *Neonatology Journal*, 92(2), pp.73-82.
- Siimes MA, Salmenpera L, Perheentups J. (1984). Exclusive breast-feeding for 9 months: risk of iron deficiency. *The Journal of Pediatrics*. 104:196-99.
- Soedjatmiko. (2007). Pengaruh defisiensi besi terhadap kecerdasan dan perilaku anak, Pendekatan praktis pucat: masalah kesehatan yang terabaikan pada bayi dan anak: Jakarta, Departemen Ilmu Kesehatan Anak FKUI, hal. 72-86.
- Steinbicker AU, Muckenthaler MU. (2013). Out of balance—systemic iron homeostasis in iron-related disorders. *The Journal of Nutrients*; 5: 3034-61.
- Stevens GA, Finucane MM, De-Regil LM, Paciorek CJ, Flaxman SR, Branca F *et al.* (2013). Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995-2011: a systematic analysis of population-representative data. *Lancet Global Health Journal*. 1:E16-E25.
- Stotzfus RJ, Dreyfuss ML. (2011). Guidelines for the Use of Iron Supplements to Prevent and Treat Iron Deficiency Anemia, INACG, USA.
- Sweet D., Savage G., Tubman T. *et al.* (2001). Study of maternal influences on fetal iron status at term using cord blood transferrin receptors. *Archives of Disease in Childhood - Fetal and Neonatal Edition*, 84(1), pp.40F-43.

- Taylor M., Holland C., Spencer R., Jackson J., O'connor G. and O'donnell J. (1997). Haematological Reference Ranges For Schoolchildren. *Clinical And Laboratory Haematology Journal*, 19(1), Pp.1-15.
- Thorsdottir I., Gunnarsson B., Atladottir H., Michaelsen K. and Palsson G. (2003). Iron status at 12 months of age — effects of body size, growth and diet in a population with high birth weight. *European Journal of Clinical Nutrition*, 57(4), pp.505-513.
- Traglia M., Girelli D., Biino G., Campostrini N., *et al.* (2011). Association of HFE and Tmprss6 genetic variants with iron and erythrocyte parameters is only in part dependent on serum hepcidin concentrations. *Journal of Medical Genetics*, 48(9), pp.629-634.
- Van Rheenen PF, de Moor LT, Eschbach S, Brabin BJ. (2008). A cohort study of haemoglobin and zinc protoporphyrin levels in term Zambian infants: Effects of iron stores at birth, complementary food and placental malaria. *Eur J Clin Nutr.* 62: 1379–1387.
- WHO. (2005), *Worldwide prevalence of anaemia 1993-2005*, Who Press, Geneva, Switzerland.
- WHO. (2001). *Iron Deficiency Anaemia. Assessment, prevention and control. A guide for programme managers*. Accessed January 8, 2018.
- Worwood M. (1997). The laboratory assessment of iron status—an update. *Internation Journal of Clinical Chemistr.*259:3–23
- Wu, AC, Lesperance, L & Bernstein, H. (2002). Screening for Iron Deficiency. *Pediatrics in Review*, vol. 23, no.5, pp. 171-178.
- Yao AC, Moinian M, Lind J. (1969). Distribution of blood between infant and placenta after birth. *Lancet Journal.*2:871–873.
- Yip R, Dallman PR. (1988). The roles of inflammation and iron deficiency as causes of anemia. *The American Journal of Clinical Nutrition*; 48: 1295–1300