

DAFTAR PUSTAKA

- Afzal, Z., Pogge, E., & Boomersshine, V. (2017). Evaluation of a Pharmacist and Nurse Practitioner Smoking Cessation Program. *Journal of Pharmacy Practice*, 30(4), 406–411. <https://doi.org/10.1177/0897190016659221>
- Agomo, C., Udoh, A., Kpokiri, E., & Osuku-Opio, J. (2018). Community pharmacists' contribution to public health: Assessing the global evidence base. *Clinical Pharmacist*, 10(4). <https://doi.org/10.1211/CP.2018.20204556>
- Armour, C. L., Reddel, H. K., LeMay, K. S., Saini, B., Smith, L. D., Bosnic-Anticevich, S. Z., Song, Y. J. C., Alles, M. C., Burton, D. L., Emmerton, L., Stewart, K., & Krass, I. (2013). Feasibility and effectiveness of an evidence-based asthma service in Australian community pharmacies: a pragmatic cluster randomized trial. *The Journal of Asthma: Official Journal of the Association for the Care of Asthma*, 50(3), 302–309. <https://doi.org/10.3109/02770903.2012.754463>
- Australian Government. (2018). *Australian Government Response To the Review of Pharmacy Remuneration and Regulation* (Issue May).
- Barnett, A., Ding, H., Hay, K. E., Yang, I. A., Bowman, R. V., Fong, K. M., & Marshall, H. M. (2020). The effectiveness of smartphone applications to aid smoking cessation: A meta-analysis. *Clinical EHealth*, 3, 69–81. <https://doi.org/10.1016/j.ceh.2020.09.001>
- Beaupre, L. A., Hammal, F., DeSutter, C., Stiegelmar, R. E., Masson, E., & Finegan, B. A. (2019). Impact of a standardized referral to a community pharmacist-led smoking cessation program before elective joint replacement surgery. *Tobacco Induced Diseases*, 17, 14. <https://doi.org/10.18332/tid/101600>
- Beaupre, L. A., Hammal, F., Stiegelmar, R., Masson, E., & Finegan, B. (2020). A community-based pharmacist-led smoking cessation program, before elective total joint replacement surgery, markedly enhances smoking cessation rates. *Tobacco Induced Diseases*, 18, 78. <https://doi.org/10.18332/tid/126405>
- Bouchet-Benezech, B., Champanet, B., & Rouzaud, P. (2018). Smoking cessation at the pharmacy: feasibility and benefits based on a French observational study with six-month follow-up. *Substance Abuse and Rehabilitation*, 9, 31–42. <https://doi.org/10.2147/SAR.S152186>
- Breen, R. J., Frandsen, M., & Ferguson, S. G. (2021). Incentives for smoking cessation in a rural pharmacy setting: The Tobacco Free Communities program. *The Australian Journal of Rural Health*, 29(3), 455–463.

<https://doi.org/10.1111/ajr.12724>

- Caponnetto, P., DiPiazza, J., Aiello, M. R., & Polosa, R. (2017). Training pharmacists in the stage-of-change model of smoking cessation and motivational interviewing: A randomized controlled trial. *Health Psychology Open*, 4(2). <https://doi.org/10.1177/2055102917736429>
- Condinho, M., Ramalhinho, I., & Sinogas, C. (2021). Smoking Cessation at the Community Pharmacy: Determinants of Success from a Real-Life Practice. *Pharmacy (Basel, Switzerland)*, 9(3). <https://doi.org/10.3390/pharmacy9030143>
- Curry, S., Grothaus, L., & McBride, C. (1997). *Reasons for quitting: intrinsic and extrinsic motivation for smoking cessation in a population-bases sample of smokers*. 22(6), 727.
- Deeks, L. S., Kosari, S., Develin, A., Peterson, G. M., & Naunton, M. (2019). Smoking cessation and the general practice pharmacist. *Journal of Smoking Cessation*, 14(3), 186–189. <https://doi.org/10.1017/jsc.2019.3>
- Dent, L. A., Harris, K. J., & Noonan, C. W. (2007). Tobacco interventions delivered by pharmacists: a summary and systematic review. *Pharmacotherapy*, 27(7), 1040–1051. <https://doi.org/10.1592/phco.27.7.1040>
- Doll, R., PeBoreham, J., & Sutherland, I. (2004). Mortality in relation to smoking: 50 years observations on male British doctors. *BMJ*, 328(7455), 1519.
- El Hajj, M. S., Kheir, N., Al Mulla, A. M., Al-Badriyeh, D., Al Kaddour, A., Mahfoud, Z. R., Salehi, M., & Fanous, N. (2015). Assessing the effectiveness of a pharmacist-delivered smoking cessation program in the State of Qatar: Study protocol for a randomized controlled trial. *Trials*, 16(1), 1–12. <https://doi.org/10.1186/s13063-015-0570-z>
- El Hajj, M. S., Kheir, N., Al Mulla, A. M., Shami, R., Fanous, N., & Mahfoud, Z. R. (2017). Effectiveness of a pharmacist-delivered smoking cessation program in the State of Qatar: a randomized controlled trial. *BMC Public Health*, 17(1), 215. <https://doi.org/10.1186/s12889-017-4103-4>
- Eriksen, M., & Mackay, J. (2015). The tobacco atlas, Fifth Edition. *Atlanta: The American Cancer Society*.
- Fai, S. C., Yen, G. K., & Malik, N. (2016). Quit rates at 6 months in a pharmacist-led smoking cessation service in Malaysia. *Canadian Pharmacists Journal : CPJ = Revue Des Pharmaciens Du Canada : RPC*, 149(5), 303–312. <https://doi.org/10.1177/1715163516662894>
- Farley, A., Tearne, S., Taskila, T., Williams, R. H., Mackaskill, S., Etter, J.-F., &

- Aveyard, P. (2017). A mixed methods feasibility study of nicotine-assisted smoking reduction programmes delivered by community pharmacists - The RedPharm study. *BMC Public Health*, 17(1). <https://doi.org/10.1186/s12889-017-4116-z>
- Fiore, M. C., Jaén, C., Baker, T. B., Bailey, W. C., Benowitz, N. L., Curry, S. J., Dorfman, S. F., Froelicher, E., & Goldstein, M. (2008). A Clinical Practice Guideline for Treating Tobacco Use and Dependence: 2008 Update. In *U.S. Department of Health and Human Services* (Vol. 35).
- Gong, J., Baker, C. L., Zou, K. H., Bruno, M., Jumadilova, Z., Lawrence, D., Wilson, B., & Ewel, C. (2016). A pragmatic randomized trial comparing telephone-based enhanced pharmacy care and usual care to support smoking cessation. *Journal of Managed Care and Specialty Pharmacy*, 22(12), 1417–1425. <https://doi.org/10.18553/jmcp.2016.22.12.1417>
- Hartmann-Boyce, J., Hong, B., Livingstone-Banks, J., Wheat, H., & Fanshawe, T. R. (2019). Additional behavioural support as an adjunct to pharmacotherapy for smoking cessation. *Cochrane Database of Systematic Reviews*, 2019(6). <https://doi.org/10.1002/14651858.CD009670.pub4>
- Hartmann-Boyce, Jamie, Stead, L. F., Cahill, K., & Lancaster, T. (2013). Efficacy of interventions to combat tobacco addiction: Cochrane update of 2012 reviews. *Addiction (Abingdon, England)*, 108(10), 1711–1721. <https://doi.org/10.1111/add.12291>
- Herbec, A., Brown, J., Shahab, L., West, R., & Raupach, T. (2019). Pragmatic randomised trial of a smartphone app (NRT2Quit) to improve effectiveness of nicotine replacement therapy in a quit attempt by improving medication adherence: results of a prematurely terminated study. *Trials*, 20(1 CC-Tobacco Addiction), 547. <https://doi.org/10.1186/s13063-019-3645-4>
- Hollands, G. J., Naughton, F., Farley, A., Lindson, N., & Aveyard, P. (2019). Interventions to increase adherence to medications for tobacco dependence. *Cochrane Database of Systematic Reviews*, 2019(8). <https://doi.org/10.1002/14651858.CD009164.pub3>
- Kawachi, I., Colditz, G. A., Stampfer, J., Willett, W. C., Manson, J. E., Rosner, B., Speizer, F. E., & Hennekens, C. H. (1994). Smoking Cessation and Time Course of Decreased Risks of Coronary Heart Disease in Middle-Aged Women. *Archives of Internal Medicine*, 154(2), 169–175. <https://doi.org/10.1001/archinte.1994.00420020075009>
- Komalasari, D., & Helmi, A. (2000). Faktor-faktor penyebab perilaku merokok pada remaja. *Jurnal Psikologi*, 28, 37–47.

- Law, M., & Tang, J. L. (1995). An analysis of the effectiveness of interventions intended to help people stop smoking. *Archives of Internal Medicine*, 155(18), 1933–1941.
- Le Foll, B., Melihan-Cheinin, P., Rostoker, G., & Lagrue, G. (2005). Smoking cessation guidelines: Evidence-based recommendations of the French Health Products Safety Agency. *AFSSAPS*, 20, 431–441.
- Madurasinghe, V. W., Sohanpal, R., James, W., Steed, L., Eldridge, S., Taylor, S., Griffiths, C., & Walton, R. (2017). Smoking treatment optimisation in pharmacies (STOP): a cluster randomised pilot trial of a training intervention. *Pilot Feasibility Stud*, 3 CC-T, 1. <https://doi.org/10.1186/s40814-016-0120-9>
- Manfredi, C., Crittenden, K. S., R, W., J, E., Y, C., & C, S. (1999). Evaluation of a motivational smoking cessation intervention for women in Public Health Clinics. *Preventive Med*, 28, 51–60.
- McRobbie, H., & McEwen, A. (2005). *Helping smokers to stop: advice for pharmacists in England*. April, 9. <https://www.ncsct.co.uk/usr/pub/helping-smokers-stop-guidance-for-pharmacist-in-england.pdf>
- Mitchell, E. A., & Milerad, J. (2006). Smoking and the sudden infant death syndrome. *Reviews on Environmental Health*, 21(2), 81–103. <https://doi.org/10.1515/reveh.2006.21.2.81>
- O'Reilly, E., Frederick, E., & Palmer, E. (2019). Models for pharmacist-delivered tobacco cessation services: a systematic review. *Journal of the American Pharmacists Association : JAPhA*, 59(5), 742–752. <https://doi.org/10.1016/j.japh.2019.05.022>
- Onda, M., Kuwanoe, T., Hashimoto, A., Horiguchi, M., Domichi, M., & Sakane, N. (2019). Pharmacist-Delivered Smoking Cessation Program in Community Pharmacy (The FINE Program) in Japan—The Development of a Training Course and a Feasibility Study. *Journal of Pharmacy Practice*. <https://doi.org/10.1177/0897190019889745>
- Oskamp, S., & Schultz, W. (1998). *Applied social psychology*. Prenticehall.
- Page, M. J. et al. (2021). The PRISMA 2020 statement: An updated for reporting systematic reviews. *The BMJ*, 372. doi: 10.1136/bmj.n71.
- Palavra, I. R., Franelić, I. P., Milanović, S. M., & Puljić, K. (2013). [Passive smoking--active killer]. In *Lijecnicki vjesnik* (Vol. 135, Issues 11–12, pp. 326–329).
- Prochaska, J. O., & DiClemente, C. C. (1982). Transtheoretical therapy toward a more integrative model of change. In *Psychotherapy: Theory, Research*,

Practice (Vol. 19, pp. 276–288).

- Raw, M., & McNeill, A. (1994). The prevention of tobacco-related disease. *Addiction* (Abingdon, England), 89(11), 1505–1509. <https://doi.org/10.1111/j.1360-0443.1994.tb03750.x>
- Saba, M., Diep, J., Saini, B., & Dhippayom, T. (2014). Meta-analysis of the effectiveness of smoking cessation interventions in community pharmacy. *Journal of Clinical Pharmacy and Therapeutics*, 39(3), 240–247. <https://doi.org/10.1111/jcpt.12131>
- Saputra, A. M., & Sary, N. M. (2013). Konseling Model Transteoritik dalam Perubahan Perilaku Merokok pada Remaja. *Jurnal Kesehatan Masyarakat Nasional*, 8(4), 152–157. <https://media.neliti.com/media/publications/39540-ID-konseling-model-transteoritik-dalam-perubahan-perilaku-merokok-pada-remaja.pdf>
- Service, U. D. o. H. a. H. (2014). Preventing Tobacco Use among Youth and Young Adults: We Can Make the Next Generation Tobacco-Free. A Report of the Surgeon General. *Atlanta*.
- Sheridan, R. (1992). Health psychology: challenging the biomedical model. In *John Willey & Sons Inc.*
- Sinclair, H. K., Silcock, J., Bond, C. M., Lennox, A. S., & Winfield, A. J. (1999). The cost-effectiveness of intensive pharmaceutical intervention in assisting people to stop smoking. *International Journal of Pharmacy Practice*, 7(2), 107–112. <https://doi.org/10.1111/j.2042-7174.1999.tb00957.x>
- Sohanpal, R., Jumbe, S., James, W.-Y., Steed, L., Yau, T., Rivas, C., Madurasinghe, V., Houlihan, C., Berdunisov, V., Taylor, M., & al., et. (2019). Evaluating the effectiveness and cost-effectiveness of the Smoking Treatment Optimisation in Pharmacies (STOP) intervention: protocol for a cluster randomised controlled trial. *Trials*, 20(1 CC-Tobacco Addiction). <https://doi.org/10.1186/s13063-019-3368-6>
- Susanna, D., Budi, H., & Hendra, F. (2003). Penentuan kadar nikotin dalam asap rokok. *Jurnal Kesehatan*, 7(9), 47.
- Taylor, B. V., Oudit, G. Y., Kalman, P. G., & Liu, P. (1998). Clinical and pathophysiological effects of active and passive smoking on the cardiovascular system. *The Canadian Journal of Cardiology*, 14(9), 1129–1139.
- United States Public Health Service Office of the Surgeon General, & N. C. for C. D. P. and H. P. (US) O. on S. and H. (2020). *Smoking Cessation: A Report of the Surgeon General*. US Department of Health and Human Services.

- Walker, N., Smith, B., Barnes, J., Verbiest, M., Kurdziel, T., Parag, V., Pokhrel, S., & Bullen, C. (2019). Cytisine versus varenicline for smoking cessation for Māori (the indigenous people of New Zealand) and their extended family: protocol for a randomized non-inferiority trial. *Addiction (Abingdon, England)*, 114(2), 344–352. <https://doi.org/10.1111/add.14449>
- Walker, N., Smith, B., Barnes, J., Verbiest, M., Parag, V., Pokhrel, S., Wharakura, M.-K., Lees, T., Cubillos Gutierrez, H., Jones, B., & Bullen, C. (2021). Cytisine versus varenicline for smoking cessation in New Zealand indigenous Māori: a randomized controlled trial. *Addiction (Abingdon, England)*, 116(10), 2847–2858. <https://doi.org/10.1111/add.15489>
- Whincup, P. H., Gilg, J. A., Emberson, J. R., Jarvis, M. J., Feyerabend, C., Bryant, A., Walker, M., & Cook, D. G. (2004). Passive smoking and risk of coronary heart disease and stroke: Prospective study with cotinine measurement. *British Medical Journal*, 329(7459), 200–204. <https://doi.org/10.1136/bmj.38146.427188.55>
- WHO. (2015). *WHO global report on trends in prevalence of tobacco smoking*. https://apps.who.int/iris/bitstream/handle/10665/156262/9789241564922_eng.pdf;jsessionid=18318DE7B3C3CE50967CF3B71C8B4B5D?sequence=1
- Wolf, T., Gepstein, L., Dror, U., Hayam, G., Shofti, R., Zaretzky, A., Uretzky, G., Oron, U., & Ben-Haim, S. A. (2001). Relationship of current and past smoking to mortality and morbidity in patients with left ventricular dysfunction. *Journal of the American College of Cardiology*, 37(6), 1677–1682. [https://doi.org/10.1016/S0735-1097\(01\)01195-0](https://doi.org/10.1016/S0735-1097(01)01195-0)
- World Health Organization. (2014). *Toolkit for delivering the 5A's and 5R's brief tobacco interventions in primary care*. 17.
- World Health Organization. (2017). *Prevalence of Tobacco Smoking*. World Health Organization.
- World Health Organization. (2018). Heart disease and stroke are the commonest ways by which tobacco kills people. *World Health Organization*, 1(1), 1–2. http://www.searo.who.int/tobacco/data/ino_rtc_reports
- World Health Organization. (2019). Country profile Indonesia WHO report on the global tobacco epidemic 2019. *World Health Organization*, 1–10. https://www.who.int/tobacco/surveillance/policy/country_profile/idn.pdf
- Zwar, N. (2008). Smoking cessation--what works? *Australian Family Physician*, 37(1–2), 10–14.
- Zwar, N., Richmond, R., Borland, R., Stillman, S., Bennett, M., & Litt, J. (2005). Smoking cessation guidelines for Australian general practice. *Australian*

Family Physician, 34, 461–466.